

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 54 (2008)
Heft: 1-2

Artikel: Metastable embedding, 2-equivalence and generic rigidity of flag manifolds
Autor: Golver, Henry
DOI: <https://doi.org/10.5169/seals-109903>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

34

METASTABLE EMBEDDING, 2-EQUIVALENCE AND GENERIC RIGIDITY OF FLAG MANIFOLDS

by Henry GLOVER

CONJECTURE 34.1. *Any 2-equivalent manifolds embed in the same metastable dimension. I.e., let M^n and N^n be two simply connected closed differentiable manifolds such that their 2-localizations are homotopy equivalent. If M^n embeds in \mathbf{R}^{n+k} , $k \geq (n+3)/2$, then N^n embeds in euclidean space of the same dimension, cf. [7].*

R. Rigdon [15] proved this result in the case that there exists a global map, $f: M \rightarrow N$ realizing this 2-equivalence, e.g., an odd covering. Glover, Mislin [8] and independently Bendersky [1] proved an analogous result for immersing manifolds in euclidean space. Glover, Mislin [9] proved an analogous result for the number of linearly independent tangent vector fields on a smooth manifold. Although the embedding result would just be a technical generalization of Rigdon's result it still seems interesting and would apply to such situations as the Hilton, Roitberg criminal H -manifolds [11], or manifolds made by Zabrodsky mixing [16].

CONJECTURE 34.2. *All complex flag manifolds are generically rigid. I.e., given a simply connected space X of finite type, let $\mathcal{G}(X)$ denote the (Mislin) genus of X , the set of all homotopy types $[Y]$, of simply connected, finite type spaces Y , such that the p -localization of X is homotopy equivalent to the p -localization of Y , for all primes p . We say that a simply connected, finite type X is generically rigid or generically trivial if $\mathcal{G}(X) = \{[X]\}$, the single homotopy type. A complex flag manifold is any space G/H , where $G = U(n)$ and $H = U(n_1) \times U(n_2) \times \cdots \times U(n_k)$, with $\sum_{i=1}^k n_i = n$.*

See [9] for cases when Conjecture 34.2 has been proved. These include complex Grassmann manifolds and complete flag manifolds $U(n)/T^n$, where

$T^n = \prod_{i=1}^n U(1)$. Note that Papadima has proved this result in the context of G any compact Lie group and H its maximal torus [14].

A survey of the Mislin genus is given in [13]. Many simply connected spaces of finite type fail to be generically trivial. First examples are $|\mathcal{G}(\mathbf{HP}^n)| = 2^k$, where k is the number of primes p , such that $2 \leq p \leq 2n-1$.

This conjecture began with the author's question to Albrecht Dold in 1973 of why we didn't know more manifolds with the *fixed point property* (every self map has a fixed point). The obvious ones at that point were the real, complex projective spaces of even dimension and all quaternionic projective spaces (except \mathbf{HP}^1) as shown by the Lefschetz fixed-point formula. Dold suggested the Grassmann manifold of complex 2-planes (through the origin) in 5-dimensional complex space, $U(5)/(U(2) \times U(3))$. This was correct as seen by applying the Lefschetz fixed-point formula to the integral cohomology ring

$$H^*(U(p+q)/(U(p) \times U(q)); \mathbf{Z}) = \mathbf{Z}[c, \bar{c}]/\{c\bar{c} = 1\},$$

showing there were only Adams maps, $c_i \mapsto \lambda^i c_i$ for $i = 1, 2, \dots, p$, in this case $p = 2$, $q = 3$. Here c is the total Chern class of the canonical p -plane bundle over this Grassmann manifold and \bar{c} the total Chern class of the canonical q -plane bundle. In [4] it is shown that this result is true in general for $p \gg q$. This result led to the independent proofs by Stephen Brewster (OSU PhD thesis 1978) [2] and Mike Hoffman [12] that the only cohomology ring endomorphisms of Grassmann manifolds $U(p+q)/(U(p) \times U(q))$ were given by Adams maps when $p \neq q$, and $\lambda \neq 0$, and $c_i \mapsto \bar{c}_i$, $i = 1, 2, \dots, p$, when $p = q$.

The results in [5] give a conjecture for all the integral cohomology ring endomorphisms of the general complex flag manifold and as a consequence give the conjecture that all the rational cohomology ring automorphisms are given by Adams maps, and actions of the Weyl group N/H , where N is the normalizer of $H = \prod_{i=1}^k U(n_i)$, $\sum_{i=1}^k n_i = n$, in $G = U(n)$. It is this conjecture, proved in special cases, that gives the results in [10] and would prove Conjecture 34.2. Another consequence of the cohomology ring endomorphism conjecture would be a complete classification of which complex flag manifolds have the fixed point property (cf. [6]). There are a number of other applications of the cohomology ring endomorphism and automorphism theorems, e.g., by S. Papadima [14] to isometry invariant geodesics, and P. Gilkey [3] to the classification of Hermitian Riemannian manifolds.

REFERENCES

- [1] BENDERSKY, M. A functor which localizes the higher homotopy groups of an arbitrary CW complex. In: *Localization in Group Theory and Homotopy Theory*, 13–21. Lecture Notes in Mathematics 418. Springer-Verlag, 1974.
- [2] BREWSTER, S. Automorphisms of the cohomology ring of finite Grassmann manifolds. Ph.D. Dissertation, Ohio State University, Columbus, 1978.
- [3] GILKEY, P. Bundles over projective spaces and algebraic curvature tensors. *J. Geom.* 71 (2001), 54–67.
- [4] GLOVER, H. and W. HOMER. Endomorphisms of the cohomology ring of finite Grassmann manifolds. In: *Proceedings of the Evanston Conference on the Geometrical Applications of Homotopy Theory (1977)*, 170–193. Lecture Notes in Mathematics 657. Springer-Verlag, 1978.
- [5] GLOVER, H. and W. HOMER. Self-maps of flag manifolds. *Trans. Amer. Math. Soc.* 267 (1981), 423–434.
- [6] GLOVER, H. and W. HOMER. Fixed points on flag manifolds. *Pacific J. Math.* 101 (1982), 303–306.
- [7] GLOVER, H. and G. MISLIN. Metastable embedding and 2-localization. In: *Localization in Group Theory and Homotopy Theory*, 48–57. Lecture Notes in Mathematics 418. Springer-Verlag, 1974.
- [8] GLOVER, H. and G. MISLIN. Immersion in the metastable range and 2-localization. *Proc. Amer. Math. Soc.* 43 (1974), 443–448.
- [9] GLOVER, H. and G. MISLIN. Vector fields on 2-equivalent manifolds. In: *Proceedings of the Conference on Homotopy Theory (1974), Evanston*, 29–45. Mexican Math. Soc., 1977.
- [10] GLOVER, H. and G. MISLIN. On the genus of generalized flag manifolds. *L'Enseignement Math.* (2) 27 (1981), 211–219.
- [11] HILTON, P. and J. ROITBERG. On principal S^3 -bundles over spheres. *Ann. of Math.* (2) 90 (1969), 91–107.
- [12] HOFFMAN, M. Endomorphisms of the cohomology of complex Grassmannians. *Trans. Amer. Math. Soc.* 281 (1984), 745–760.
- [13] MCGIBBON, C. The Mislin genus of a space. In: *The Hilton Symposium 1993: Topics in Topology and Group Theory*, 75–102. CRM Proceedings and Lecture Notes 6. Amer. Math. Soc., 1994.
- [14] PAPADIMA, S. Rigidity properties of compact Lie groups modulo maximal tori. *Math. Ann.* 275 (1986), 637–652.
- [15] RIGDON, R. p -equivalences and embeddings of manifolds. *J. London Math. Soc.* (2) 11 (1975), 233–244.
- [16] ZABRODSKY, A. *Hopf Spaces*. Mathematics Studies 22. North-Holland, Amsterdam, 1976.

Henry Glover

The Ohio-State University
 231 West 18th Avenue
 43210 Columbus, OH
 USA
e-mail: glover@math.ohio-state.edu