

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 54 (2008)
Heft: 1-2

Artikel: Self-similar contracting groups
Autor: Grigorchuk, Rostislav I. / Nekrashevych, Volodymyr
DOI: <https://doi.org/10.5169/seals-109904>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 06.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

35

SELF-SIMILAR CONTRACTING GROUPS

by Rostislav I. GRIGORCHUK and Volodymyr NEKRASHEVYCH

Let X be a finite set (alphabet) and let X^* be the free monoid generated by it. We imagine X^* as a rooted tree with the root equal to the empty word and a word v connected to every word of the form vx for $x \in X$.

DEFINITION 35.1. A *self-similar group* is a group G acting faithfully on X^* such that for every $g \in G$ and $x \in X$ there exist $h \in G$ and $y \in X$ such that

$$g(xv) = yh(v)$$

for all $v \in X^*$.

It follows that for every $g \in G$ and $u \in X^*$ there exists $h \in G$ such that

$$g(uv) = g(u)h(v)$$

for all $v \in X^*$. The element h is denoted $g|_v$ and is called the *restriction of g in v* .

DEFINITION 35.2. A self-similar group G acting on X^* is called *contracting* if there exists a finite set $N \subset G$ such that for every $g \in G$ there exists $n \in \mathbf{N}$ such that $g|_v \in N$ for all $v \in X^*$ of length $|v| \geq n$.

CONJECTURE 35.3. *Finitely generated contracting groups have solvable conjugacy problem.*

CONJECTURE 35.4. *Finitely generated contracting groups have solvable membership problem.*

REMARK 35.5. It is known that the word problem in contracting groups is solvable in polynomial time.

The next three conjectures are ordered by their strength (the last one is the strongest).

CONJECTURE 35.6. *Contracting groups have no non-abelian free subgroups.*

CONJECTURE 35.7. *Contracting groups are amenable.*

CONJECTURE 35.8. *A simple random walk on a contracting group has zero entropy.*

CONJECTURE 35.9. *The group generated by the transformations a and b of $\{0, 1\}^*$ defined by*

$$a(0w) = 1w, \quad a(1w) = 0a(w), \quad b(0w) = 0b(w), \quad b(1w) = 1a(w)$$

is amenable.

REMARK 35.10. This group is not contracting; however, it is known (due to a result of S. Sidki [4]) that this group does not contain a free subgroup. The graphs of the action of this group on the boundary of the tree X^* have intermediate growth.

The next conjecture was suggested by Zoran Šunić.

CONJECTURE 35.11. *The group H_k generated by the transformations a_{ij} of $\{1, 2, \dots, k\}^*$, for $1 \leq i < j \leq k$, defined by*

$$a_{ij}(iw) = jw, \quad a_{ij}(jw) = iw, \quad a_{ij}(kw) = ka_{ij}(w) \quad \text{for } k \neq i, j$$

is non-amenable for $k \geq 4$.

REMARK 35.12. The group H_k models the “Hanoi tower game” on k pegs. The graph of its action on the n th level of the tree $\{1, \dots, k\}^*$ coincides with the graph of the game with n discs [3]. It is also not contracting, but the graphs of its action on the boundary of the tree are of intermediate growth. The group H_3 is amenable and the graphs of the action on the boundary have polynomial growth.

Following Furstenberg we say that a group is *Tychonoff* if it has a fixed ray for any affine action on a convex cone with compact base [1]. A definition of branch groups can be found in [2]. Every proper quotient of a branch group is virtually abelian.

CONJECTURE 35.13. *A branch group G is Tychonoff if and only if G is indicable⁴⁾ and every proper non-trivial quotient is Tychonoff.*

ADDED IN PROOF. Conjecture 35.6 is now a theorem by the second author. See V. NEKRASHEVYCH. ‘Free subgroups in groups acting on rooted trees’. Preprint arXiv: math.GR/0802.2554 (2008).

REFERENCES

- [1] GRIGORCHUK, R. I. On Tychonoff groups. In: *Geometry and Cohomology in Group Theory (Durham, 1994)*, 170–187. London Math. Soc. Lecture Note Ser. 252. Cambridge Univ. Press, Cambridge, 1998.
- [2] —— Just infinite branch groups. In: *New Horizons in pro- p Groups (M. du Sautoy, D. Segal and A. Shalev, eds.)*, 121–179. Progress in Mathematics 184. Birkhäuser Verlag, Basel, 2000.
- [3] GRIGORCHUK, R. I. and Z. ŠUNIĆ. Asymptotic aspects of Schreier graphs and Hanoi Towers groups *C. R. Acad. Sci. Paris Sér. I Math.* 342 (2006), 545–550.
- [4] SIDKI, S. Finite automata of polynomial growth do not generate a free group. *Geom. Dedicata* 108 (2004), 193–204.

Rostislav I. Grigorchuk

Mathematics Department
Texas A&M University
College Station, TX 77843-3368
USA
e-mail: grigorch@math.tamu.edu

Volodymyr Nekrashevych

Mathematics Department
Texas A&M University
College Station, TX 77843-3368
USA
e-mail: nek rash@math.tamu.edu

⁴⁾ An infinite group is called *indicable* if it admits a homomorphism onto \mathbf{Z} .