Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	54 (2008)
Heft:	1-2
Artikel:	The genus of a group
Autor:	Gruenberg, Karl
DOI:	https://doi.org/10.5169/seals-109905

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

## Download PDF: 06.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# 36

### THE GENUS OF A GROUP

### by Karl GRUENBERG

Localization methods arise in infinite group theory and also, in a seemingly different incarnation, in integral representations of finite groups. Is there a common generalization?

Let  $\pi$  be a finite set of primes. A group P is  $\pi$ -local if  $x \mapsto x^m$  is bijective for all integers m coprime to  $\pi$ . Every (abstract, discrete) group Ghas (essentially) a unique  $\pi$ -localization  $\phi_{\pi}: G \to G_{\pi}$  (meaning  $G_{\pi}$  is  $\pi$ -local and any homomorphism from G to a  $\pi$ -local group factors uniquely through  $\phi_{\pi}$ ). Guido Mislin and Peter Hilton began the study of localizations of finitely generated nilpotent groups that led Guido to introduce the *genus* of a finitely generated nilpotent group [2]. With this as a starting point, we make the following definition (it coincides with Guido's for finitely generated nilpotent groups whose centre has finite index): the *genus*  $\mathcal{G}(G)$  of G is all isomorphism classes [H] of groups H such that H is finitely generated and residually of finite exponent and  $H_{\pi} \simeq G_{\pi}$  for all finite sets  $\pi$ ; write  $H \lor G$ . (If all structural requirements on H were dropped, then  $\mathcal{G}(G)$  would be infinite for all G, which would not be a satisfactory situation.)

Guido's paper [2] is concerned with finitely generated nilpotent groups whose centre has finite index. What happens for finitely generated abelian-by-finite groups? This question was successfully investigated by Niamh O'Sullivan ([3], [4], [5]). Her techniques involve the module version of genus.

Recall that if Q is a finite group and A, B are  $\mathbb{Z}Q$ -lattices, then  $A \vee B$ (same genus) means that  $A_{\pi} \simeq B_{\pi}$  for all finite sets  $\pi$ . A pointed lattice is a pair (A, x) where  $x \in H^2(Q, A)$  and  $(A, x) \vee (B, y)$  means there exists a Q-map  $f: A \rightarrow B$  with finite cokernel of order prime to |Q| (such maps exist if, and only if,  $A \vee B$ ) and  $f_*(x) = y$ . Let  $\mathcal{G}(A, x)$  denote all isomorphism classes [B, y] such that  $(A, x) \vee (B, y)$ .

#### K. GRUENBERG

Let G be a finitely generated abelian-by-finite group, choose m so that  $A := \langle g^m | g \in G \rangle$  is free abelian of finite index in G and write Q = G/A. Let x be the cohomology class of the resulting extension.

(1) There is a well defined surjective map  $\theta$ :  $\mathcal{G}(A, x) \twoheadrightarrow \mathcal{G}(G)$  and  $\mathcal{G}(A, x)$  is finite.

(2) There is an explicitly defined subgroup J of AutQ that acts on  $\mathcal{G}(A, x)$ and  $\theta$  induces a bijection  $\mathcal{G}(A, x)/J \xrightarrow{\sim} \mathcal{G}(G)$ .

The point here is that the left hand side of (2) is better suited for calculations than is the right hand side: cf. O'Sullivan's papers for explicit examples, including new derivations of some of Guido's results.

What is the natural level of generalization for this point of view? For example, do the basic connexions that we have outlined carry over to the class of polycyclic-by-finite groups? A relevant fact here is that if G is such a group, then the number of isomorphism classes of polycyclic-by-finite groups H in the genus of G is known to be finite [1].

When Guido wrote his 1974 paper he added some problems at the end; so he clearly thought there was unfinished business here<sup>5</sup>). I rather think that this is still true today.

### REFERENCES

- [1] GRUNEWALD, F. J., P. F. PICKEL and D. SEGAL. Polycyclic groups with isomorphic quotients. Ann. of Math. (2) 111 (1980), 155–195.
- [2] MISLIN, G. Nilpotent groups with finite commutator subgroups. In: Localization in group theory and related topics, 103–120. Lecture Notes in Mathematics 418. Springer-Verlag, 1974.
- [3] O'SULLIVAN, N. The genus and localization of finitely generated (torsion-free abelian)-by-finite groups. *Math. Proc. Cambridge Philos. Soc. 128* (2000), 257–268.
- [4] Genus and localization of virtually nilpotent groups. In: Crystallographic Groups and Their Generalizations: Workshop Leuven (1999), 253–262. Contemp. Math. 262. Amer. Math. Soc., 2000.
- [5] Genus and cancellation. Comm. Algebra 28 (2000), 3387–3400.

<sup>&</sup>lt;sup>5</sup>) His Problem 1 is solved in C. Casacuberta, C. Cassidy, D. Scevenels, 'On genus and embeddings of torsion-free nilpotent groups of class two', *Manuscripta Math.* 92 (1997) 463–475; his Problem 2 is solved in [5].