

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 54 (2008)  
**Heft:** 1-2

**Artikel:** Alperin's weight conjecture  
**Autor:** Kessar, Radha / Linckelmann, Markus  
**DOI:** <https://doi.org/10.5169/seals-109911>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 27.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## 42

### ALPERIN'S WEIGHT CONJECTURE

by Radha KESSAR and Markus LINCKELMANN

Let  $p$  be a prime number.

CONJECTURE 42.1. *Let  $G$  be a finite group and let  $P$  be a Sylow  $p$ -subgroup of  $G$ .*

(i) *The number of conjugacy classes of  $p'$ -elements of  $G$  is greater than or equal to the number of conjugacy classes of  $N_G(P)/P$ .*

(ii) *If  $P$  is abelian, then the number of conjugacy classes of  $G$  is greater than or equal to the number of conjugacy classes of  $N_G(P)$ .*

The above inequalities would follow from Alperin's weight conjecture [1] which we describe now.

Let  $k$  be an algebraically closed field of characteristic  $p$ . For a finite group  $H$  denote by  $l(kH)$  the number of isomorphism classes of simple  $kH$ -modules, and by  $w(kH)$  the number of isomorphism classes of simple projective  $kH$ -modules. The weight conjecture predicts the following

CONJECTURE 42.2. *Let  $G$  be a finite group. Then*

$$l(kG) = \sum_{Q \in \mathcal{I}} w(kN_G(Q)/Q),$$

where  $\mathcal{I}$  denotes a set of representatives of  $G$ -conjugacy classes of  $p$ -subgroups of  $G$ .

Conjecture 42.2 comes in a block-wise version as well. The reformulation of this conjecture in terms of alternating chains in [13] paved the way for many extensions (see for instance [9], [10], [11], [16]). Despite having been verified for many families of finite groups, including finite  $p$ -solvable groups,

symmetric groups, and in some cases for finite groups of Lie type and some sporadic simple groups (see for instance [2], [3], [4], [5], [6], [8], [12]), a true understanding of Conjecture 42.2 or indeed of Conjecture 42.1 remains elusive.

In its original form stated above, Alperin's weight conjecture is a numerical equality interpreting the number of simple modules of a finite group or a  $p$ -block in terms of the involved  $p$ -local structure. In subsequent years structural approaches to this and related conjectures in terms of linear source modules [7], fusion category algebras [14], and cohomological invariants of functors over certain finite categories [15], have emerged.

We briefly explain how 42.1 would follow from 42.2. By a theorem of Brauer,  $l(kG)$  is equal to the number of conjugacy classes of  $p'$ -elements in  $G$ , and the summand for  $Q = P$  on the right side of the equality 42.2 is equal to the number of conjugacy classes of  $N_G(P)/P$ . Thus 42.2 implies the inequality 42.1 (i). The inequality 42.1 (ii) follows from 42.1 (i) applied to centralizers of  $p$ -elements.

## REFERENCES

- [1] ALPERIN, J.L. Weights for finite groups. In: *The Arcata Conference on Representations of Finite Groups*, 369–379. Proc. Sympos. Pure Math. 47. Amer. Math. Soc., Providence, R.I., 1987.
- [2] ALPERIN, J.L. and P. FONG. Weights for symmetric and general linear groups. *J. Algebra* 131 (1990), 2–22.
- [3] AN, J. 2-weights for general linear groups. *J. Algebra* 149 (1992), 500–527.
- [4] ——— Weights for classical groups. *Trans. Amer. Math. Soc.* 342 (1994), 1–42.
- [5] ——— The Alperin and Uno conjectures for the Fischer simple group  $\text{Fi}_{22}$ . *Comm. Algebra* 33 (2005), 1529–1557.
- [6] AN, J. and R. A. WILSON. The Alperin weight conjecture and Uno's conjecture for the Baby Monster  $\mathbf{B}$ ,  $p$  odd. *London Math. Soc. J. Comput. Math.* 7 (2004), 120–166.
- [7] BOLTJE, R. Alperin's weight conjecture in terms of linear source modules and trivial source modules. In: *Modular Representation Theory of Finite Groups (Charlottesville, VA, 1998)*, 147–155. Walter de Gruyter, Berlin, 2001.
- [8] CABANES, M. and M. ENGUEHARD. On fusion in unipotent blocks. *Bull. London Math. Soc.* 31 (1999), 143–148.
- [9] DADE, E.C. Counting characters in blocks, I. *Invent. Math.* 109 (1992), 187–210.
- [10] ——— Counting characters in blocks, II. *J. Reine Angew. Math.* 448 (1994), 97–190.

- [11] ——— Counting characters in blocks, II.9. In: *Representation Theory of Finite Groups*, 45–59. Ohio State Univ. Math. Res. Inst. Publ. 6. Walter de Gruyter, Berlin, 1997.
- [12] ISAACS, I.M. and G. NAVARRO. Weights and vertices for characters of  $\pi$ -separable groups. *J. Algebra* 177 (1995), 339–366.
- [13] KNÖRR R. and G.R. ROBINSON. Some remarks on a conjecture of Alperin. *J. London Math. Soc. (2)* 39 (1989), 48–60.
- [14] LINCKELMANN, M. Fusion category algebras. *J. Algebra* 277 (2004), 222–235.
- [15] ——— Alperin’s weight conjecture in terms of equivariant Bredon cohomology. *Math. Z.* 250 (2005), 495–513.
- [16] ROBINSON, G.R. Local structure, vertices and Alperin’s conjecture. *Proc. London Math. Soc. (3)* 72 (1996), 312–330.

Radha Kessar, Markus Linckelmann

Department of Mathematical Sciences

University of Aberdeen

Meston Building

Aberdeen AB24 3UE

United Kingdom

*e-mail*: kessar@maths.abdn.ac.uk, m.linckelmann@maths.abdn.ac.uk