Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 54 (2008)
Heft: 1-2

Artikel: Relative completions of linear groups
Autor: Knudson, Kevin P.
DOI: https://doi.org/10.5169/seals-109912

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 15.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

L'Enseignement Mathématique (2) 54 (2008), 123–125

43

RELATIVE COMPLETIONS OF LINEAR GROUPS

by Kevin P. KNUDSON

Here is a question that I've thought about a lot, but I can't seem to solve. The classical Malcev completion of a group is well known. It has a universal mapping property that allows one to generalize the definition as follows. Let k be a field and let G be a group. The *unipotent k-completion* of G is a prounipotent k-group $\mathcal{U}$ that is universal among such groups admitting a map from G. The Malcev completion is the case $k = \mathbf{Q}$.

One possible problem with this construction is that it might be trivial; that is, the group $\mathcal{U}$ may consist of a single element. This happens, for example, when $H_1(G,k) = 0$. To get around this, there is a generalization (due to Deligne) called the *relative completion*. The set-up is the following. Suppose G is a discrete group and that $\rho\colon G \to S$ is a representation of G in a semisimple algebraic k-group S. Assume that the image of ρ is Zariski dense. The *completion of G relative to ρ* is a proalgebraic k-group $\mathcal{G}$ that is an extension of S by a prounipotent k-group $\mathcal{U}$:

$$1 \longrightarrow \mathcal{U} \longrightarrow \mathcal{G} \longrightarrow S \longrightarrow 1\,,$$

along with a lift $\tilde{\rho}\colon G \to \mathcal{G}$ of ρ. The group $\mathcal{G}$ should satisfy the obvious universal mapping property. If S is the trivial group, then this reduces to the unipotent completion. Full details about this construction may be found in [1], [2].

Consider the group $G = \mathrm{SL}_n(k[t])$ with the map $\rho\colon \mathrm{SL}_n(k[t]) \to \mathrm{SL}_n(k)$ induced by setting $t = 0$.

QUESTION 43.1. *What is the completion of G relative to ρ ?*

There is an obvious guess, namely the group $\mathrm{SL}_n(k[[T]])$, and this turns out to be correct sometimes.

I proved this when k is a number field or a finite field, and $n \geq 3$ [2]. The proof goes like this. Let K be the kernel of ρ; this is the *congruence subgroup of the ideal* (t). Filter K by powers of (t): $K^i = \{A \in K : A \equiv I \mod t^i\}$. Then it is easy to see that for each i, $K^i/K^{i+1} \cong \mathfrak{sl}_n(k)$. Moreover, the filtration $K^\bullet$ turns out to be the lower central series in this case, and so it follows that the unipotent k-completion of K is $\varprojlim K/K^i = \ker\{\mathrm{SL}_n(k[[T]]) \xrightarrow{T=0} \mathrm{SL}_n(k)\}$. General properties of the relative completion (e.g., it is always a *split* extension) then imply that the correct answer is $\mathrm{SL}_n(k[[T]])$.

This approach fails for other fields though. Here's why. Denote the lower central series of K by $\Gamma^\bullet$. For any field, there is a short exact sequence

$$1 \longrightarrow K^2/\Gamma^2 \longrightarrow H_1(K, \mathbf{Z}) \longrightarrow K/K^2 \longrightarrow 1\,.$$

The last group is $\mathfrak{sl}_n(k)$, and most of the time, the kernel K^2/Γ^2 surjects onto the module $\Omega^1_{k/\mathbf{Z}}$ [4]. Recall that this is the k-module generated by symbols df, where the f range over k, subject to the relations $d(fg) = f\,dg + g\,df$ for $f, g \in k$, and $dr = 0$ for $r \in \mathbf{Z}$ (here, we mean the image of r under the map $\mathbf{Z} \to k$). For finite fields and number fields, this is no obstruction since it is easily seen that $\Omega^1_{k/\mathbf{Z}} = 0$, but for $k = \mathbf{C}$, for example, we see that K^2/Γ^2 is very large. So $K^\bullet$ differs wildly from $\Gamma^\bullet$ and it is therefore not easy to compute the unipotent completion of K.

Still, I conjecture that $\mathrm{SL}_n(k[[T]])$ is the correct answer all the time. In fact, I make the following, more ambitious, conjecture.

CONJECTURE 43.2. *Let k be a field and let C be a smooth affine curve over k. Denote the coordinate ring of C by A and assume that C has a k-rational point with associated maximal ideal $\mathfrak{m} \subset A$. Let ρ: $\mathrm{SL}_n(A) \to \mathrm{SL}_n(k)$ be induced by the isomorphism $A/\mathfrak{m} \to k$. Finally, let $\widehat{A}$ be the $\mathfrak{m}$-adic completion of A. Then the completion of $\mathrm{SL}_n(A)$ relative to ρ is the group $\mathrm{SL}_n(\widehat{A})$.*

I proved [2] that this is true if we replace A by the localization of A at $\mathfrak{m}$. And, not surprisingly, it is true when k is a number field [3].

REFERENCES

[1] HAIN, R.M. Completions of mapping class groups and the cycle $C - C^{-}$. In: *Mapping Class Groups and Moduli Spaces of Riemann Surfaces (Göttingen, 1991 and Seattle, WA, 1991)*, 75–105. Contemp. Math. *150*. Amer. Math. Soc., 1993.

[2] KNUDSON, K.P. Relative completions and the cohomology of linear groups over local rings. *J. London Math. Soc. (2) 65* (2002), 183–203.

[3] —— Relative completions and K_2 of curves. Preprint arXiv: math.KT/0502553 (2005).

[4] KRUSEMEYER, M.I. Fundamental groups, algebraic K-theory, and a problem of Abhyankar. *Invent. Math. 19* (1973), 15–47.

Kevin P. Knudson

Department of Mathematics and Statistics
Mississippi State University
Mississippi State, MS 39762
USA
e-mail: knudson@math.msstate.edu