

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 54 (2008)
Heft: 1-2

Artikel: A vanishing conjecture for the homology of congruence subgroups
Autor: Arlettaz, Dominique
DOI: <https://doi.org/10.5169/seals-109871>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 06.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

2

A VANISHING CONJECTURE FOR THE HOMOLOGY OF CONGRUENCE SUBGROUPS

by Dominique ARLETTAZ

For any prime number p , let $\Gamma_{n,p}$ denote the *congruence subgroup* of $\mathrm{SL}_n(\mathbf{Z})$ of level p , i.e., the kernel of the surjective homomorphism $f_p: \mathrm{SL}_n(\mathbf{Z}) \rightarrow \mathrm{SL}_n(\mathbf{F}_p)$ induced by the reduction mod p , and let us write $\Gamma_p = \varinjlim_n \Gamma_{n,p}$, where the limit is defined by upper left inclusions. Notice that the groups $\Gamma_{n,p}$ are not homology stable with \mathbf{Z} -coefficients (see [2]).

If p is odd, then the group Γ_p is torsion-free. Therefore, it is of particular interest to detect torsion classes in the integral (co)homology of Γ_p . It turns out that $H^*(\Gamma_p; \mathbf{Z})$ contains 2-torsion elements in arbitrarily large dimensions (see Corollary 1.10 of [1]). Groups like this are called groups with “very strange torsion” by S. Weintraub in 1986.

However, vanishing results for the (co)homology of Γ_p are also extremely useful. Let us propose the following

CONJECTURE 2.1. *For an odd integer n and an odd prime p , the homology group $H_n(\Gamma_p; \mathbf{Z})$ contains no q -torsion if q is a sufficiently large prime (in comparison to n), $q \neq p$.*

As far as I know, this problem is not solved, but one should notice its relationship with the study of the Dwyer–Friedlander map

$$\varphi_{\mathbf{Z}}: (K_n(\mathbf{Z}))_q \rightarrow K_n^{\text{ét}}(\mathbf{Z}[\frac{1}{q}])$$

relating the q -torsion of algebraic K -theory to étale K -theory. This map is known to be surjective and it is conjecturally an isomorphism (this is a version of the Quillen–Lichtenbaum conjecture, see [3], Theorem 8.7 and Remark 8.8).

For $q \neq p$, the Dwyer–Friedlander map and the reduction mod p induce the commutative diagram

$$\begin{array}{ccc} (K_n(\mathbf{Z}))_q & \xrightarrow{\varphi_{\mathbf{Z}}} & K_n^{\text{ét}}(\mathbf{Z}[\frac{1}{q}]) \\ \downarrow (f_p)_* & & \downarrow \\ (K_n(\mathbf{F}_p))_q & \xrightarrow{\varphi_{\mathbf{F}_p}} & K_n^{\text{ét}}(\mathbf{F}_p) \end{array}$$

The map $\varphi_{\mathbf{F}_p}$ is an isomorphism, since the K -theory of finite fields is completely known. If we define $A_n = \ker \varphi_{\mathbf{Z}}$ and $B_n = \ker(f_p)_*$, this implies that A_n is contained in B_n .

On the other hand, one can show by using Postnikov decompositions that B_n is a direct summand of $(H_n(\Gamma_p; \mathbf{Z}))_q$ for large enough primes $q \neq p$ (see [1], Introduction and Theorem 2.1).

Consequently, the proof of the above conjecture for n odd and q a large enough prime would imply the vanishing of B_n and therefore the vanishing of A_n which provides the assertion that the Dwyer–Friedlander map $\varphi_{\mathbf{Z}}$ is an isomorphism.

REFERENCES

- [1] ARLETTAZ, D. Torsion classes in the cohomology of congruence subgroups. *Math. Proc. Cambridge Philos. Soc.* 105 (1989), 241–248.
- [2] CHARNEY, R. On the problem of homology stability for congruence subgroups. *Comm. Algebra* 12 (1984), 2081–2123.
- [3] DWYER, W. G. and E. M. FRIEDLANDER. Algebraic and étale K -theory. *Trans. Amer. Math. Soc.* 292 (1985), 247–280.

Dominique Arlettaz

Université de Lausanne
 Faculté des géosciences et de l’environnement
 CH-1015 Lausanne
 Switzerland
e-mail: Dominique.Arlettaz@unil.ch