

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 54 (2008)  
**Heft:** 1-2

**Artikel:**  $\mathbb{L}^p$ -homology of one-relator groups  
**Autor:** Linnell, Peter A.  
**DOI:** <https://doi.org/10.5169/seals-109918>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# 49

## $\ell^p$ -HOMOLOGY OF ONE-RELATOR GROUPS

by Peter A. LINNELL

Let  $G$  be a group and let

$$\dots \longrightarrow \mathbf{C}G^{e_{n+1}} \xrightarrow{d_n} \mathbf{C}G^{e_n} \xrightarrow{d_{n-1}} \dots \xrightarrow{d_1} \mathbf{C}G^{e_1} \xrightarrow{d_0} \mathbf{C}G \longrightarrow \mathbf{C} \longrightarrow 0$$

be a free  $\mathbf{C}G$ -resolution of  $\mathbf{C}$ . Let  $1 \leq p < \infty$  and for  $n \geq 0$ , let

$$\begin{aligned} d_*^n : \mathbf{C}G^{e_{n+1}} \otimes_{\mathbf{C}G} \ell^p(G) &\longrightarrow \mathbf{C}G^{e_n} \otimes_{\mathbf{C}G} \ell^p(G), \\ d_n^* : \mathrm{Hom}_{\mathbf{C}G}(\mathbf{C}G^{e_n}, \ell^p(G)) &\longrightarrow \mathrm{Hom}_{\mathbf{C}G}(\mathbf{C}G^{e_{n+1}}, \ell^p(G)) \end{aligned}$$

be the maps induced by  $d_n$ ; for convenience, we let  $d_*^{-1} = d_{-1}^* = 0$ . Then one has the usual homology and cohomology groups

$$\begin{aligned} H_n(G, \ell^p(G)) &= \ker d_*^{n-1} / \mathrm{im} \, d_*^n, \\ H^n(G, \ell^p(G)) &= \ker d_n^* / \mathrm{im} \, d_{n-1}^*, \end{aligned}$$

which we shall call the (unreduced)  $\ell^p$ -homology and cohomology groups of  $G$  respectively. In the case when all the  $e_n$  are finite,

$$\mathbf{C}G^{e_n} \otimes_{\mathbf{C}G} \ell^p(G) \cong \ell^p(G)^{e_n} \cong \mathrm{Hom}_{\mathbf{C}G}(\mathbf{C}G^{e_n}, \ell^p(G)),$$

so one can also define the reduced  $\ell^p$ -homology and cohomology groups of  $G$ :

$$\begin{aligned} \overline{H}_n(G, \ell^p(G)) &= \ker d_*^{n-1} / \overline{\mathrm{im} \, d_*^n}, \\ \overline{H}^n(G, \ell^p(G)) &= \ker d_n^* / \overline{\mathrm{im} \, d_{n-1}^*}, \end{aligned}$$

where  $\overline{\phantom{x}}$  indicates the closure in  $\ell^p(G)^{e_n}$ . The first  $\ell^p$ -cohomology groups (reduced and unreduced) have been studied extensively recently by Bekka, Bourdon, Martin, Valette and others. Also Kappos has interesting results on general reduced homology and cohomology groups.

Let us concentrate now on the case where  $G$  is a finitely generated torsion-free one-relator group. Let  $d$  denote the number of generators of  $G$ . Then we have a  $\mathbf{C}G$ -resolution of the form

$$0 \longrightarrow \mathbf{C}G \longrightarrow \mathbf{C}G^d \longrightarrow \mathbf{C}G \longrightarrow \mathbf{C} \longrightarrow 0.$$

From this it is clear that the homology groups  $H_n(G, \ell^p(G))$ ,  $H^n(G, \ell^p(G))$ ,  $\bar{H}_n(G, \ell^p(G))$  and  $\bar{H}^n(G, \ell^p(G))$  are all zero for  $n \geq 3$ . Warren Dicks and I determined the  $\ell^2$ -Betti numbers of such groups [2], Theorem 4.2; an immediate consequence of this is that  $H_2(G, \ell^p(G)) = \bar{H}_2(G, \ell^p(G)) = 0$  for  $p \leq 2$ . The proof of this depended on the results that a torsion-free one-relator group is left orderable, and that if  $H$  is a left orderable group,  $0 \neq \alpha \in \mathbf{C}H$  and  $0 \neq \theta \in \ell^2(H)$ , then  $\alpha\theta \neq 0$  [4], Theorem 2. However if  $p > 2$ , then we can have  $0 \neq \alpha \in \mathbf{C}H$  and  $0 \neq \theta \in \ell^p(H)$  with  $\alpha\theta = 0$ ; see [5] for information about this. Thus we have the following conjecture:

**CONJECTURE 49.1.** *Let  $G$  be a finitely generated torsion-free one-relator group and let  $1 \leq p < \infty$ . Then  $H_2(G, \ell^p(G)) = \bar{H}_2(G, \ell^p(G)) = 0$ .*

By Poincaré duality, this conjecture is true if  $G$  is a surface group, orientable or not. If Conjecture 49.1 is true, then it follows from Kappos's [3], Proposition 3.5 that  $\bar{H}^2(G, \ell^p(G)) = 0$  for all  $p > 1$ . The situation for unreduced cohomology is less clear. Let us consider the special case  $p = 2$ . Recall that  $\langle x, y \mid x^2y^2 = 1 \rangle$  is the Klein bottle group. Our next conjecture is:

**CONJECTURE 49.2.** *Let  $G$  be a finitely generated torsion-free one-relator group. Then  $H^2(G, \ell^2(G)) = 0$ , provided  $G$  has neither  $\mathbf{Z} \times \mathbf{Z}$  nor the Klein bottle group as a free factor.*

This conjecture is also true if  $G$  is a surface group, orientable or not. On the other hand if  $G$  is  $\mathbf{Z} \times \mathbf{Z}$  or the Klein bottle group, then  $H^2(G, \ell^2(G)) \neq 0$ , and then it follows from the Mayer–Vietoris sequence that  $H^2(G * \mathbf{Z}, \ell^2(G * \mathbf{Z})) \neq 0$ ; of course  $G * \mathbf{Z}$  is still a torsion-free one-relator group.

Finally we consider the first homology groups. A result of Guichardet [1], Theorem A shows that if  $G$  is an arbitrary infinite group, then the natural epimorphism  $H^1(G, \ell^2(G)) \rightarrow \bar{H}^1(G, \ell^2(G))$  is an isomorphism if and only if  $G$  is non-amenable. This leads to the following conjecture.

**CONJECTURE 49.3.** *Let  $G$  be a finitely generated torsion-free one-relator group that has neither  $\mathbf{Z} \times \mathbf{Z}$  nor the Klein bottle group as a free factor. Then the natural epimorphism  $H_1(G, \ell^2(G)) \rightarrow \overline{H}_1(G, \ell^2(G))$  is an isomorphism.*

I am very grateful to the referee for pointing out that my original formulations of Conjectures 49.2 and 49.3 were too easy to answer.

#### REFERENCES

- [1] BEKKA, M. E. B. and A. VALETTE. Group cohomology, harmonic functions and the first  $L^2$ -Betti number. *Potential Anal.* 6 (1997), 313–326.
- [2] DICKS, W. and P.A. LINNELL.  $L^2$ -Betti numbers of one-relator groups. *Math. Ann.* 337 (2007), 855–874.
- [3] KAPPOS, E.  $\ell^p$ -cohomology for groups of type  $FP_n$ . Preprint arXiv: math.FA/0511002 (2005–2006).
- [4] LINNELL, P.A. Zero divisors and  $L^2(G)$ . *C. R. Acad. Sci. Paris Sér. I Math.* 315 (1992), 49–53.
- [5] LINNELL, P.A. and M.J. PULS. Zero divisors and  $L^p(G)$ , II. *New York J. Math.* 7 (2001), 49–58.

Peter A. Linnell

Math Department  
 Virginia Tech  
 Blacksburg, VA 24061-0123  
 USA  
*e-mail:* linnell@math.vt.edu