

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 54 (2008)  
**Heft:** 1-2

**Artikel:** The Farrell-Jones Conjecture in algebraic K- and L-theory  
**Autor:** Lück, Wolfgang  
**DOI:** <https://doi.org/10.5169/seals-109919>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 16.07.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# 50

## THE FARRELL–JONES CONJECTURE IN ALGEBRAIC $K$ - AND $L$ -THEORY

by Wolfgang LÜCK

One of my favorite conjectures in geometric topology is the *Borel Conjecture*. It says that a homotopy equivalence between closed aspherical topological manifolds is homotopic to a homeomorphism. This implies that two closed aspherical topological manifolds are homeomorphic if and only if their fundamental groups are isomorphic. (This is not true if one considers smooth manifolds and replaces homeomorphic by diffeomorphic.) This is the topological analogue of Mostow rigidity.

Another prominent conjecture is the *Novikov Conjecture*. It asserts that the higher signatures of a closed oriented smooth manifold are homotopy invariants. It is motivated by the signature formula of Hirzebruch.

Furthermore the *Bass Conjecture* has gotten a lot of attention since it was formulated. It says that for a group  $G$  and an integral domain  $R$  of characteristic 0 the *Hattori–Stallings rank* of a finitely generated projective  $RG$ -module evaluated at an element  $g$  of  $G$  is non-trivial only if  $g$  has finite order  $|g|$  and  $|g| \cdot 1_R$  is not a unit in  $R$ .

Finally we mention the conjectures that for a torsionfree group  $G$  the *reduced projective class group*  $\tilde{K}_0(\mathbf{Z}G)$  and its *Whitehead group*  $\text{Wh}(G)$  vanish. The last two conjectures have equivalent geometric counterparts, namely: if the fundamental group  $G$  is finitely presented and torsionfree, then any *finitely dominated CW-complex* is homotopy equivalent to a finite CW-complex and any *h-cobordism* over a closed manifold of dimension  $\geq 5$  is trivial.

Also a lot of work was done about the *Kaplansky Conjecture* which predicts for an integral domain  $R$  and a torsionfree group  $G$  that 0 and 1 are the only idempotents in the group ring  $RG$ .

It turns out the Farrell–Jones Conjecture for algebraic  $K$ - and  $L$ -theory does imply all of the conjectures above and gives a good understanding of the algebraic  $K$ - and  $L$ -theory of group rings  $RG$  in terms of the algebraic  $K$ - and  $L$ -theory of the coefficient ring  $R$  and the homology of the group  $G$ . I will formulate it only for a torsionfree group  $G$  and a regular ring  $R$ . In this case the *Farrell–Jones Conjecture* predicts that the classical assembly maps

$$\begin{aligned} H_n(BG; \mathbf{K}_R) &\xrightarrow{\cong} K_n(RG); \\ H_n(BG; \mathbf{L}_R) &\xrightarrow{\cong} L_n(RG); \end{aligned}$$

are bijective for all  $n$ . The sources of the assembly maps above are homology theories such that  $H_n(\text{pt}; \mathbf{K}_R) = K_n(R)$  and  $H_n(\text{pt}; \mathbf{L}_R) = L_n(R)$  hold. In the  $L$ -theory case one does not need regular. The general formulation uses more elaborate *equivariant homology theories* and *classifying spaces of families*.

The original formulation of the *Farrell–Jones Conjecture* can be found in [3], 1.6 on page 257. The Farrell–Jones Conjecture is still open but known for a good class of groups. Its non-commutative counterpart is the *Baum–Connes Conjecture*. For more explanations, information about the status and references concerning the Farrell–Jones Conjecture we refer for instance to the survey article [4]. For very recent new developments we refer to [1] and [2].

#### REFERENCES

- [1] BARTELS, A., W. LÜCK and H. REICH. The  $K$ -theoretic Farrell–Jones conjecture for hyperbolic groups. *Invent. Math.* 172 (2008), 29–70.
- [2] BARTELS, A., W. LÜCK and H. REICH. On the Farrell–Jones conjecture and its applications. *J. Topology* 1 (2008), 57–86.
- [3] FARRELL, F. T. and L. E. JONES. Isomorphism conjectures in algebraic  $K$ -theory. *J. Amer. Math. Soc.* 6 (1993), 249–297.
- [4] LÜCK, W. and H. REICH. The Baum–Connes and the Farrell–Jones conjectures in  $K$ - and  $L$ -theory. In: *Handbook of  $K$ -theory*. Vol. 1, 2, 703–842. Springer-Verlag, Berlin, 2005.

W. Lück

Westfälische Wilhelms-Universität Münster  
 Einsteinstr. 62  
 D-48149 Münster  
 Germany  
*e-mail:* lueck@math.uni-muenster.de