

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 54 (2008)
Heft: 1-2

Artikel: The uniform Kazhdan property for $SL_n(\mathbb{Z})$, n3
Autor: Arzhantseva, Goulmira N.
DOI: <https://doi.org/10.5169/seals-109872>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 28.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

3

THE UNIFORM KAZHDAN PROPERTY FOR $SL_n(\mathbf{Z})$, $n \geq 3$

by Goul'nara N. ARZHANTSEVA

Let Γ be a discrete group, and let S be a finite subset of Γ . For a unitary representation π of Γ in a separable Hilbert space \mathcal{H} we define the number

$$K(\pi, \Gamma, S) = \inf_{0 \neq u \in \mathcal{H}} \max_{s \in S} \frac{\|\pi(s)u - u\|}{\|u\|}.$$

Then the *Kazhdan constant* of Γ with respect to S is defined as

$$K(\Gamma, S) = \inf_{\pi} K(\pi, \Gamma, S),$$

where the infimum is taken over unitary representations π having no invariant vectors. We also define the *uniform Kazhdan constant* of Γ as

$$K(\Gamma) = \inf_S K(\Gamma, S),$$

where the infimum is taken over all finite generating sets S of Γ .

A group Γ is said to have *Kazhdan property (T)* (or to be a *Kazhdan group*) if there exists a finite subset S of Γ with $K(\Gamma, S) > 0$. A group Γ is *uniform Kazhdan* if $K(\Gamma) > 0$.

Shortly after its introduction by David Kazhdan in the mid 60's, property (T) was used by Gregory Margulis to give a first explicit construction of infinite families of expander graphs of bounded degree. In particular, a major problem of practical application in the design of efficient communication networks was solved.

A classical example of a Kazhdan group is the group $SL_n(\mathbf{Z})$ for $n \geq 3$ (for more details and a general context of locally compact groups see the recent book [2]). Surprisingly, the following question is still open.

QUESTION 3.1. *Is the group $SL_n(\mathbf{Z})$, for $n \geq 3$, uniform Kazhdan?*

Infinite finitely generated uniform Kazhdan groups were discovered very recently, see [6] or [1]. However, these groups are neither finitely presented nor residually finite. The latter construction provides an infinite uniform Kazhdan group that weakly (see [4]) contains an infinite family of expanders in its Cayley graph.

An affirmative answer to the above question would give, in particular, the first example of a residually finite (and, in addition, finitely presented) infinite uniform Kazhdan group. It is crucial for applications: infinite families of expanders could be constructed independently of the choice of the group generating set.

A negative answer would be interesting as well. In that case, this classical group would belong to the class of non-uniform Kazhdan groups. First examples of such groups were obtained using Lie groups (see [3]). Then, all word hyperbolic groups were also shown to have zero uniform Kazhdan constant (see [5]).

REFERENCES

- [1] ARZHANTSEVA, G. and T. DELZANT. A random group with uniform Kazhdan property. *Preliminary version*.
- [2] BEKKA, B., P. DE LA HARPE and A. VALETTE. Kazhdan's Property (T). *New Mathematical Monographs II*. Cambridge University Press, Cambridge, 2008.
- [3] GELANDER, T. and A. ZUK. Dependence of Kazhdan constants on generating subsets. *Israel J. Math.* 129 (2002), 93–98.
- [4] GROMOV, M. Random walk in random groups. *Geom. Funct. Anal.* 13 (2003), 73–146.
- [5] OSIN, D. Kazhdan constants of hyperbolic groups. *Funktsional. Anal. i Prilozhen.* 36 (2002), 46–54; translation in *Funct. Anal. Appl.* 36 (2002), 290–297.
- [6] OSIN, D. and D. SONKIN. Uniform Kazhdan groups. Preprint arXiv: math.GR/0606012 (2006).

Goulnara N. Arzhantseva

Département de Mathématiques
Université de Genève
2-4, rue du Lièvre
CH-1211 Genève 4
Switzerland
e-mail: goulnara.arzhantseva@math.unige.ch