

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 54 (2008)  
**Heft:** 1-2

**Artikel:** Automatic permutations  
**Autor:** Bartholdi, Laurent  
**DOI:** <https://doi.org/10.5169/seals-109875>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 27.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## 6

### AUTOMATIC PERMUTATIONS

by Laurent BARTHOLDI

Finite state automata are wonderful little devices to produce interesting groups — for examples, see [1] or [2]. We fix a finite set  $X$ , called the *alphabet*. An *automaton*  $\mathcal{A}$  is given by a finite set  $Q$ , a map  $\tau: Q \times X \rightarrow X \times Q$ , and an element  $q \in Q$ .

It defines an *automatic transformation*, still written  $\mathcal{A}: X^* \rightarrow X^*$  on strings over  $X$ , by the following rule: given  $w = x_1 \dots x_n$ , set  $q_1 = q$ , and compute  $\tau(q_1, x_1) = (y_1, q_2)$ ,  $\tau(q_2, x_2) = (y_2, q_3)$ , etc. up to  $\tau(q_n, x_n) = (y_n, q_{n+1})$ . Then the image of  $w$  under  $\mathcal{A}$  is  $y_1 \dots y_n$ .

Part of the interest in these transformations is that they can be used to generate groups (and semigroups) with striking features, such as infinite finitely generated torsion groups, groups of intermediate word growth, of non-uniformly exponential word growth, etc.

It is easy to test whether an automatic transformation is the identity — assuming  $Q$  is minimal, this happens if  $\tau(*, x) = (x, *)$  for all possible  $*$ 's. It is also easy to test whether it is invertible — this happens if the composition  $x \rightarrow (q, x) \rightarrow \tau(q, x) = (y, *) \rightarrow y$  is a permutation of  $X$  for all  $q \in Q$ . It is finally easy to see that the product and inverse of automatic transformations are again automatic.

Therefore, in a group  $G$  generated by automatic invertible transformations, all elements of  $G$  are themselves represented by automatic transformations. It follows that the most basic decision problem — determining whether a word over a given generating set is trivial — is solvable for such groups.

What seems difficult to test, and is the object of this conjecture, is whether an automatic permutation  $\mathcal{A}$  has finite order; i.e. whether the following decision problem is solvable: “determine whether the cyclic subgroup generated by a given group element is finite or infinite”.

If the order of  $\mathcal{A}$  is finite, then this can be checked by computing powers of  $\mathcal{A}$  and seeing that one of them induces the identity transformation. But how can one determine that  $\mathcal{A}$  has infinite order? This would follow from

CONJECTURE 6.1. *If  $\mathcal{A}$  has finite order, then its order divides  $(\#X!)^{\#Q}$ .*

A sharper bound should actually be  $\exp(\text{Sym}(X))^{\#Q}$ . If  $X = \mathbf{Z}/d\mathbf{Z}$  and  $\sigma \in \text{Sym}(X)$  has order  $e$  and  $Q = \{0, \dots, q\}$ , the automaton defined by initial state  $q$  and

$$\tau(i, x) = \begin{cases} (x, 0) & \text{if } i = 0, \\ (x + 1, i - 1) & \text{if } i > 0 \text{ and } x = 0, \\ (x + 1, 0) & \text{if } i > 0 \text{ and } x \neq 0 \end{cases}$$

has order  $e^q$ .

#### REFERENCES

- [1] BARTHOLDI, L., R. GRIGORCHUK, and V. NEKRASHEVYCH. From fractal groups to fractal sets. In: *Fractals in Graz 2001: Analysis, Dynamics, Geometry, Stochastics*, Trends Math., Birkhäuser, Basel, 2003, 25–118.
- [2] BARTHOLDI, L., R. GRIGORCHUK and Z. ŠUNÍK. Branch groups. In: *Handbook of Algebra, vol. 3*. North-Holland, Amsterdam, 2003, 989–1112.

Laurent Bartholdi

EPFL  
 SB IMB MAD  
 CH-1015 Lausanne  
 Switzerland  
*e-mail*: laurent.bartholdi@gmail.com