**Zeitschrift:** L'Enseignement Mathématique
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique
**Band:** 54 (2008)
**Heft:** 1-2

**Artikel:** The extended quotient
**Autor:** Baum, Paul
**DOI:** https://doi.org/10.5169/seals-109877

**Nutzungsbedingungen**
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

**Conditions d'utilisation**
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

**Terms of use**
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 12.01.2026

**ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch**

*L'Enseignement Mathématique (2) 54 (2008), 23–26*

# 8

## THE EXTENDED QUOTIENT

by Paul BAUM

Let $\Gamma$ be a finite group acting on a (topological space $X$ or) an affine variety $X$. The quotient variety (or quotient topological space) $X/\Gamma$ is obtained by collapsing each orbit to a point. For $x \in X$, $\Gamma_x$ denotes the stabilizer group of $x$, that is

$$\Gamma_x = \{\gamma \in \Gamma \mid \gamma x = x\},$$

and $c(\Gamma_x)$ denotes the set of conjugacy classes of $\Gamma_x$.

The *extended quotient* is obtained by replacing the orbit of $x$ by $c(\Gamma_x)$. This is done as follows:

Set $\widetilde{X} = \{(\gamma, x) \in \Gamma \times X \mid \gamma x = x\}$, this is an affine variety and is a sub-variety of $\Gamma \times X$. Moreover, $\Gamma$ acts on $\widetilde{X}$ as follows

$$\begin{aligned} \Gamma \times \widetilde{X} &\longrightarrow \widetilde{X} \\ (g, (\gamma, x)) &\mapsto g(\gamma, x) = (g\gamma g^{-1}, gx) \quad \text{where } g \in \Gamma,\ (\gamma, x) \in \widetilde{X}. \end{aligned}$$

The extended quotient, denoted by $X//\Gamma$, is given by $\widetilde{X}/\Gamma$, the ordinary quotient of $\widetilde{X}$ by the above action of $\Gamma$.

The extended quotient is an affine variety (or a topological space). The evident projection $\widetilde{X} \to X$, $(\gamma, x) \mapsto x$ passes to quotient spaces to give a map $\rho\colon X//\Gamma \to X/\Gamma$. The map $\rho$ is the projection of the extended quotient onto the ordinary quotient.

Let $G$ be a reductive $p$-adic group (examples are $\mathrm{GL}(n, F)$ and $\mathrm{SL}(n, F)$, where $F$ is any finite extension of the $p$-adic numbers $\mathbf{Q}_p$). Let $V$ be a vector space over the complex numbers $\mathbf{C}$.

DEFINITION 8.1. A representation $\phi\colon G \to \mathrm{Aut}_{\mathbf{C}}(V)$ of $G$ is *smooth* if for every $v \in V$,

$$G_v = \{g \in G \mid \phi(g)v = v\}$$

is an open subgroup of $G$.

We will denote by $\widehat{G}$ the set of equivalence classes of smooth irreducible representations of $G$. One of the main problems in the representation theory of $p$-adic groups (which is closely related to the local Langlands conjecture) is to describe $\widehat{G}$.

The *Hecke algebra* of $G$, denoted by $\mathcal{H}G$, is the convolution algebra of all complex-valued locally-constant compactly-supported functions $f\colon G \to \mathbf{C}$. Then $\widehat{G}$ is in bijection with $\operatorname{Prim}\mathcal{H}G$, the set of primitive ideals in $\mathcal{H}G$. On $\operatorname{Prim}\mathcal{H}G$ there is the Jacobson topology. Hence we may consider each connected component of the primitive ideal space. Typically there will be countably many of these connected components.

Let $\mathbf{C}^\times$ denote the (complex) affine variety $\mathbf{C} - \{0\}$.

DEFINITION 8.2. A *complex torus* is a (complex) affine variety $T$ such that there exists an isomorphism of affine varieties

$$T \cong \mathbf{C}^\times \times \mathbf{C}^\times \times \cdots \times \mathbf{C}^\times .$$

J. Bernstein assigns to each $\alpha \in \pi_0(\operatorname{Prim}\mathcal{H}G)$ a complex torus $T_\alpha$ and a finite group $\Gamma_\alpha$ acting on $T_\alpha$.

He then forms the quotient variety $T_\alpha/\Gamma_\alpha$ and proves that there is a surjective map (the infinitesimal character)

$$\pi_\alpha\colon X_\alpha \twoheadrightarrow T_\alpha/\Gamma_\alpha .$$

The set $X_\alpha$ is the connected component of $\operatorname{Prim}\mathcal{H}G$ corresponding to $\alpha$. In Bernstein's work $X_\alpha$ is a set (i.e. is only a set) so $\pi_\alpha$ is a map of sets, which is surjective, finite-to-one and generically one-to-one.

CONJECTURE 8.3. *There is a certain resemblance between*

$$\begin{array}{ccc} T_\alpha//\Gamma_\alpha & & X_\alpha \\ {\scriptstyle \rho_\alpha}\downarrow & \textit{and} & \downarrow{\scriptstyle \pi_\alpha} \\ T_\alpha/\Gamma_\alpha & & T_\alpha/\Gamma_\alpha \end{array}$$

*Here $\rho_\alpha$ is (as above) the projection of the extended quotient onto the ordinary quotient.*

For the precise conjecture, see papers of A.-M. Aubert, P. Baum and R. Plymen [1] and [2], but we now explain what is meant by *resemblance*.

For each $\alpha \in \pi_0(\mathrm{Prim}\,\mathcal{H}G)$ there exists (conjecturally) a bijection

$$\nu_\alpha : T_\alpha//\Gamma_\alpha \longrightarrow X_\alpha$$

such that:

- In the possibly non-commutative diagram

$$\begin{tikzcd} T_\alpha//\Gamma_\alpha \arrow[r, "\nu_\alpha"] \arrow[d, "\rho_\alpha"'] & X_\alpha \arrow[d, "\pi_\alpha"] \\ T_\alpha/\Gamma_\alpha \arrow[r, "I"'] & T_\alpha/\Gamma_\alpha \end{tikzcd}$$

  the bijection $\nu_\alpha : T_\alpha//\Gamma_\alpha \longrightarrow X_\alpha$ is continuous, where $T_\alpha//\Gamma_\alpha$ has the Zariski topology, $X_\alpha$ the Jacobson topology, and the composition

$$\pi_\alpha \circ \nu_\alpha : T_\alpha//\Gamma_\alpha \longrightarrow T_\alpha/\Gamma_\alpha$$

  is a morphism of algebraic varieties.

- For each $\alpha \in \pi_0(\mathrm{Prim}\,\mathcal{H}G)$, there is an algebraic family

$$\theta_t : T_\alpha//\Gamma_\alpha \longrightarrow T_\alpha/\Gamma_\alpha$$

  of morphisms of algebraic varieties, with $t \in \mathbf{C}^\times$, such that $\theta_1 = \rho_\alpha$ and $\theta_{\sqrt{q}} = \pi_\alpha \circ \nu_\alpha$, where $q$ is the order of the residue field of the $p$-adic field $F$ over which $G$ is defined, and $\pi_\alpha$ is the infinitesimal character of Bernstein.

This conjecture is true for $\mathrm{GL}(n, F)$, where $n$ is any positive integer and $F$ is any finite extension of the $p$-adic numbers $\mathbf{Q}_p$, see [3].

## REFERENCES

[1] AUBERT, A.-M., P. BAUM and R. PLYMEN. The Hecke algebra of a reductive $p$-adic group: a geometric conjecture. In: *Noncommutative Geometry and Number Theory* (K. Consani and M. Marcolli, eds.) Aspects of Math. Vol. 37, Vieweg, Wiesbaden, 2006, 1–34.

[2] AUBERT, A.-M., P. BAUM and R. PLYMEN. Geometric structure in the representation theory of $p$-adic groups. *C. R. Acad. Sci. Paris. Sér. I Math. 345* (2007), 573–578.

[3] BRODZKI, J. and R. PLYMEN. Geometry of the smooth dual of $\mathrm{GL}(n)$. *C. R. Acad. Sci. Paris Sér. I Math. 331* (2000), 213–218.

Paul Frank Baum

Mathematics Department
206 McAllister Building
The Pennsylvania State University
University Park, PA 16802
USA
*e-mail:* baum@math.psu.edu