

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 54 (2008)
Heft: 3-4

Artikel: Normal surfaces in topologically finite 3-manifolds

Autor: Tillmann, Stephan

Bibliographie

DOI: <https://doi.org/10.5169/seals-109942>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 06.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

TABLE 1
Normal surface in the figure eight knot complement

solution	$\nu(\mu)$	$\nu(\lambda)$	slope
(2, 0, 0, 0, 0, 1)	1	4	-4
(0, 2, 0, 0, 0, 1)	-1	4	4
(0, 0, 1, 2, 0, 0)	-1	-4	-4
(0, 0, 1, 0, 2, 0)	1	-4	4

quadrilateral types in M' by r, r', r'' , where $r^{(i)}$ lifts to $p^{(i)}$. The Q -matching equation is $r + r' - 2r'' = 0$. It can be worked out from the triangulation or by observing that the induced involution on quadrilateral types in M is $(p \ q')(p' \ q)(p'' \ q'')$. Thus, $\dim PQ(\mathcal{T}') = 1$ and $PF(\mathcal{T}') = \emptyset$.

The boundary curve map is defined via the induced triangulation of the double cover of the Klein bottle; using the generators from the above section, one has: $\nu(\lambda) = -2r - 2r' + 4r'' = 0$ and $\nu(\mu) = -2r' + 2r''$. Generators λ', μ' can be chosen for $H_1(B_v) \cong \mathbf{Z} \oplus \mathbf{Z}_2$ such that the map $H_1(\tilde{B}_v) \rightarrow H_1(B_v)$ is given by $\lambda \rightarrow \lambda'$ and $\mu \rightarrow (\mu')^2 = 0$. The composition

$$Q(\mathcal{T}') \rightarrow \mathbf{Z}^2 \rightarrow H_1(\tilde{B}_v; \mathbf{R}) \rightarrow H_1(B_v; \mathbf{R})$$

is then

$$N \rightarrow (-\nu_N(\lambda), \nu_N(\mu)) = (0, \nu_N(\mu)) \rightarrow \nu_N(\mu)\lambda \rightarrow \nu_N(\mu)\lambda'.$$

Since $\nu(\mu) = -2r' + 2r''$, it follows that the map $\partial: Q(\mathcal{T}') \rightarrow H_1(B_v; \mathbf{R})$ is surjective. Its restriction to integral points in $Q(\mathcal{T}')$ has image of index two in $H_1(B_v; \mathbf{Z})$, which gives a subgroup of index four in $H_1(B_v)$.

REFERENCES

- [1] BACHMAN, D. A note on Kneser-Haken finiteness. *Proc. Amer. Math. Soc.* 132 (2004), 899–902
- [2] GORDON, C. MCA. *The Theory of Normal Surfaces*. Lecture notes typeset by Richard Kent, available at <http://www.math.brown.edu/~rkent/> (2001).
- [3] JACO, W. and U. OERTEL. An algorithm to decide if a 3-manifold is a Haken manifold. *Topology* 23 (1984), 195–209.
- [4] JACO, W. and J. H. RUBINSTEIN. PL equivariant surgery and invariant decompositions of 3-manifolds. *Adv. in Math.* 73 (1989), 149–191.
- [5] JACO, W. and J. H. RUBINSTEIN. 0-efficient triangulations of 3-manifolds. *J. Differential Geom.* 65 (2003), 61–168.

- [6] GABAI, D. Essential laminations and Kneser normal form. *J. Differential Geom.* 53 (1999), 517–574 .
- [7] HATCHER, A. *Algebraic Topology*. Cambridge University Press, Cambridge, 2002.
- [8] KANG, E. Normal surfaces in non-compact 3-manifolds. *J. Aust. Math. Soc.* 78 (2005), 305–321.
- [9] KANG, E. and J.H. RUBINSTEIN Ideal triangulations of 3-manifolds I: spun normal surface theory. *Geom. Topol. Monogr.* 7, Proceedings of the Casson Fest (2004), 235–265.
- [10] MATVEEV, S. *Algorithmic Topology and Classification of 3-Manifolds*. Algorithms and Computation in Mathematics 9. Springer-Verlag, Berlin, 2003.
- [11] THOMPSON, A. Thin position and the recognition problem for S^3 . *Math. Res. Lett.* 1 (1994), 613–630.
- [12] TOLLEFSON, J. L. Normal surface Q -theory. *Pacific J. Math.* 183 (1998), 359–374.
- [13] WALSH, G. Incompressible surfaces and spunnormal form. Preprint arXiv: math.GT/0503027 (2005).
- [14] WEEKS, J. *SnapPea*, available at www.geometrygames.org.

(Reçu le 14 juin 2007)

Stephan Tillmann

Department of Mathematics and Statistics
 The University of Melbourne
 VIC 3010,
 Australia
e-mail: tillmann@ms.unimelb.edu.au