Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 55 (2009)
Heft: 1-2

Artikel: The Big Picard theorem and other results on Riemann surfaces
Autor: Arés-Gastesi, Pablo / Venkataramana, T.N.
Bibliographie
DOI: https://doi.org/10.5169/seals-110098

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 09.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

One could now use some results on Kleinian groups to conclude that $\alpha_n = 1$ (see, for example [7, I.D.4 and II.C.6]) but we prefer to give the following elementary argument. A simple computation shows that

$$\gamma_n^{-1} \begin{bmatrix} 1 & 1/m \\ 0 & 1 \end{bmatrix} \gamma_n = \begin{bmatrix} 1 & \frac{1}{m\alpha_n} \\ 0 & 1 \end{bmatrix}, \qquad \gamma_n \begin{bmatrix} 1 & 1/m \\ 0 & 1 \end{bmatrix} \gamma_n^{-1} = \begin{bmatrix} 1 & \frac{\alpha_n}{m} \\ 0 & 1 \end{bmatrix}.$$

These two transformations are in $\Gamma \cap \mathcal{T}$, and so α_n and $1/\alpha_n$ must be (positive) integers; that is, $\alpha_n = 1$. But then γ_n will be in $\Gamma \cap \mathcal{T}$ and therefore $z_n = z_n'$, since $X = \mathbf{H}/(\Gamma \cap \mathcal{T})$.

This proposition gives us a second way to prove the Big Picard Theorem without using the modular function, as follows. As in the proof given above, the function $f\colon \mathbf{D}^* \to X = \widehat{\mathbf{C}} \backslash \{\infty, 0, 1\}$ lifts to a function $\widetilde{f}\colon \mathbf{D}^* \to \mathbf{D}^*$ that gives a commutative diagram:

$$\begin{tikzcd} & \mathbf{D}^* \arrow[d, "\rho"] \\ \mathbf{D}^* \arrow[ur, "\widetilde{f}"] \arrow[r, "f"'] & X. \end{tikzcd}$$

The function $\widetilde{f}$ has a removable singularity at the origin with $\widetilde{f}(0) = 0$; the problem is to determine the behaviour of the map ρ. By Proposition 4.5 there is an ϵ such that ρ is a finite covering from $\mathbf{D}_\epsilon^*$ onto its image. But then by the Casorati-Weierstrass theorem ρ cannot have an essential singularity at the origin, and neither can f.

REFERENCES

[1] AHLFORS, L. V. *Conformal Invariants: Topics in Geometric Function Theory.* McGraw-Hill, New York, 1973.

[2] —— , *Complex Analysis.* Mc-Graw Hill, New York, 1979.

[3] CONWAY, J. B. *Functions of One Complex Variable.* Graduate Texts in Mathematics *11.* Springer-Verlag, Berlin and New York, 1973.

[4] FARKAS, H. M. and I. KRA. *Riemann Surfaces,* 2nd ed. Graduate Texts in Mathematics *71.* Springer-Verlag, New York, Heidelberg and Berlin, 1992.

[5] HATCHER, A. *Algebraic Topology.* Cambridge University Press, Cambridge, 2002.

[6] KRANTZ, S. G. *Complex Analysis: the Geometric Viewpoint.* Carus Mathematical Monographs *23.* Mathematical Association of America, Washington DC, 1990.

[7] MASKIT, B. *Kleinian Groups.* Grundlehren der mathematischen Wissenschaften *287.* Springer-Verlag, Berlin, Heidelberg, 1988.

[8] NARASIMHAN, R. *Complex Analysis in One Variable.* Birkhäuser, Basel-Boston-Berlin, 1985.

[9] NEUENSCHWANDER, E. Studies in the history of complex function theory. I. The Casorati-Weierstrass theorem. *Historia Math. 5* (1978), 139–166.

[10] PICARD, É. *Œuvres de Ch.-É. Picard*, Tome I. Éditions du Centre National de la Recherche Scientifique, Paris, 1978.

[11] RAGHUNATHAN, M. S. *Discrete Subgroups of Lie Groups.* Ergebnisse der Mathematik und ihrer Grenzgebiete *68.* Springer-Verlag, Berlin, Heidelberg, 1972.

[12] REMMERT, R. *Classical Topics in Complex Function Theory.* Graduate Texts in Mathematics *172.* Springer-Verlag, New York, Heidelberg and Berlin, 1998.

[13] RUDIN, W. *Real and Complex Analysis.* McGraw-Hill, New York, 1966.

(Reçu le 28 juin 2007; version révisée reçue le 8 janvier 2008)

Pablo Arés-Gastesi
T. N. Venkataramana

School of Mathematics
Tata Institute of Fundamental Research
Mumbai 400005
India
e-mail: pablo@math.tifr.res.in, venky@math.tifr.res.in