

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 60 (2014)

Artikel: Moduli spaces in genus zero and inversion of power series
Autor: McMullen, Curtis T.
DOI: <https://doi.org/10.5169/seals-515844>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Moduli spaces in genus zero and inversion of power series

Curtis T. McMULLEN*

Abstract. This note shows, using elementary properties of ribbon trees, that the universal formula for the inversion of power series can be obtained by counting strata in the compactified moduli space $\overline{\mathcal{M}}_{0,n}$.

Mathematics Subject Classification (2010). 32G15.

Keywords. Moduli space, Euler characteristic, stable graphs.

Let $\mathcal{M}_{0,n}$ denote the moduli space of Riemann surfaces of genus 0 with n ordered marked points. Its Deligne–Mumford compactification $\overline{\mathcal{M}}_{0,n}$ is naturally partitioned into connected strata of the form

$$S \cong \mathcal{M}_{0,n_1} \times \cdots \times \mathcal{M}_{0,n_s},$$

indexed by the different topological types of stable curves with n marked points. The stable curves in the stratum above have s irreducible components and $s - 1$ nodes; thus $\sum n_i = n + 2s - 2$.

This note provides a short proof of the following result, which shows that the universal formula for inversion of power series is encoded in the stratification of moduli space.

Theorem 1. *The formal inverse of*

$$f(x) = x - \sum_2^{\infty} a_n x^n / n!$$

is given by

$$g(x) = x + \sum_2^{\infty} b_n x^n / n!,$$

*Research supported in part by the NSF.

where

$$b_n = \sum a_{n_1} \cdots a_{n_s} \times \left(\begin{array}{c} \text{the number of strata } S \subset \bar{\mathcal{M}}_{0,n+1} \\ \text{isomorphic to } \mathcal{M}_{0,n_1+1} \times \cdots \times \mathcal{M}_{0,n_s+1} \end{array} \right).$$

That is, $g(f(x)) = x$.

Here the coefficients of $f(x)$ and $g(x)$ are regarded as elements of the polynomial ring $\mathbb{Q}[a_2, a_3, \dots]$, and the sum is over all $s \geq 1$ and all multi-indices (n_1, \dots, n_s) with $n_i \geq 2$.

Using basic properties of the Euler characteristic, we obtain:

Corollary 2 (Getzler). *The generating functions*

$$f(x) = x - \sum_{n=2}^{\infty} \chi(\mathcal{M}_{0,n+1}) \frac{x^n}{n!} \quad \text{and} \quad g(x) = x + \sum_{n=2}^{\infty} \chi(\bar{\mathcal{M}}_{0,n+1}) \frac{x^n}{n!}$$

are formal inverses of one another.

It is easy to see that $a_n = \chi(\mathcal{M}_{0,n+1}) = (-1)^n(n-2)!$, using the fibration $\mathcal{M}_{0,n+1} \rightarrow \mathcal{M}_{0,n}$. Thus by formally inverting $f(x)$, one can readily compute

$$\langle \chi(\bar{\mathcal{M}}_{0,n}) \rangle_{n=3}^{\infty} = \langle 1, 2, 7, 34, 213, 1630, 14747, 153946, 1821473, \dots \rangle.$$

Corollary 2 is a consequence of [Ge1], Theorem 5.9, stated explicitly in [LZ], Remark 4.5.3. The development in [Ge1] uses operads and yields more information, such as Betti numbers for $\bar{\mathcal{M}}_{0,n}$. Theorem 1 shows that Corollary 2 holds for any generalized Euler characteristic on the Grothendieck ring of varieties over $\bar{\mathbb{Q}}$ (cf. [Bi]).

The proof of Theorem 1 will be based on simple properties of trees. Its aim is to provide an elementary entry point to the enumerative combinatorics of moduli spaces.

Trees. A *tree* τ is a finite, connected graph with no cycles; its vertices will be denoted $V(\tau)$. The degree function $d: V(\tau) \rightarrow \mathbb{N}$ gives the number of edges incident to each vertex. To each tree we associate the monomial

$$A(\tau) = \prod_{v \in V(\tau)} A_{d(v)-1}$$

in the polynomial ring $\mathbb{Z}[A_1, A_2, A_3, \dots]$, with the convention $A_0 = 1$.

A tree is *stable* if it has no vertices of degree 2. An *endpoint* of τ is a vertex with $d(v) = 1$. We say τ is *rooted* if it has a distinguished endpoint (the root). The number of endpoints of τ , other than its root, will be denoted $N(\tau)$.

We always assume τ has *at least* one edge, so $N(\tau) \geq 1$; and the tree with *just one edge* is considered stable.

A *ribbon tree* is a rooted stable tree equipped with a cyclic ordering of the edges incident to each vertex. A ribbon structure records the same information as a planar embedding $\tau \hookrightarrow \mathbb{R}^2$ up to isotopy.

A *marked tree* is a rooted stable tree equipped with a labeling of its endpoints by the integers $1, 2, \dots, N(\tau) + 1$. We require that the root be labeled 1.

Theorem 3. *The formal inverse of $F(x) = x - \sum_2^\infty A_n x^n$ is given by*

$$(1) \quad G(x) = \sum_{\text{ribbon } \tau} A(\tau) x^{N(\tau)}.$$

Here the sum is taken over all ribbon trees, up to isomorphism.




FIGURE 1

Three ribbon trees grafted together at their roots

Proof. Suppose we are given ribbon trees τ_1, \dots, τ_d with $d \geq 2$. We can then construct a new ribbon tree τ by identifying the roots of these trees with a single vertex w , and adding a new edge leading from w to the root of τ (see Figure 1). The ribbon structure at w is determined by the ordering of the trees (τ_i) , and by the condition that the root of τ lies between τ_d and τ_1 .

Conversely, any ribbon tree with $N(\tau) \geq 2$ is obtained by applying this construction to the subtrees (τ_1, \dots, τ_d) leading away from the edge adjacent to its root. Taking into account the vertex w of degree $d + 1$ where these trees are attached, we find:

$$A(\tau) x^{N(\tau)} = A_d \prod_{i=1}^d A(\tau_i) x^{N(\tau_i)}.$$

But the right hand side above is precisely one of the terms occurring in the expression $A_d G(x)^d$. Summing over all possible values for $d = d(w)$ we obtain

$$G(x) = x + \sum_{d=2}^{\infty} A_d G(x)^d,$$

where the first term accounts for the unique tree with $N(\tau) = 1$. Rearranging terms gives $F(G(x)) = x$. \square

Corollary 4. *The formal inverse of $f(x) = x - \sum_2^\infty a_n x^n / n!$ is given by*

$$(2) \quad g(x) = \sum_{\text{marked } \tau} a(\tau) \frac{x^{N(\tau)}}{N(\tau)!},$$

where $a(\tau) = \prod_{V(\tau)} a_{d(v)-1}$ and $a_0 = 1$.

Proof. The number of ribbon structures on a given stable rooted tree τ is given by $\prod (d(v) - 1)!$. The group $\text{Aut}(\tau)$ acts freely on the space of ribbon structures, so τ contributes $\prod (d(v) - 1)! / |\text{Aut}(\tau)|$ identical terms to equation (1) for $G(x)$. Similarly, τ contributes $N(\tau)! / |\text{Aut}(\tau)|$ terms to equation (2) for $g(x)$. Setting $A_n = a_n / n!$, we find $F(x) = f(x)$ and

$$G(x) = \sum_{\text{marked } \tau} \frac{\prod (d(v) - 1)!}{N(\tau)!} A(\tau) x^{N(\tau)} = g(x),$$

so $f(g(x)) = F(G(x)) = x$. \square

Remark 1. The same reasoning shows that (2) can be rewritten as

$$f^{-1}(x) = \sum_{\text{stable } \tau} \frac{N(\tau) + 1}{|\text{Aut}(\tau)|} a(\tau) x^{N(\tau)}.$$

For example, using the trees shown in Figure 2 we find

$$f^{-1}(x) = x + \frac{a_2}{2} x^2 + \frac{(a_3 + 3a_2^2)}{6} x^3 + \frac{(a_4 + 10a_2a_3 + 15a_2^3)}{24} x^4 + O(x^5).$$

For a quite different approach to Corollary 4, see [Ge2], Theorem 1.3.




FIGURE 2
The stable trees with $N(\tau) \leq 4$

Proof of Theorem 1. A stable curve $X \in \bar{\mathcal{M}}_{0,n+1}$ of genus zero determines a marked tree $t(X)$ whose interior vertices correspond to the irreducible components of X , and whose edges correspond to its nodes and labeled points. Conversely, any marked tree with $N(\tau) \geq 2$ can be realized by a stable curve, so the map

$$\tau \mapsto S(\tau) = \{X \in \bar{\mathcal{M}}_{0,N(\tau)+1} : t(X) \cong \tau\}$$

gives a bijection between marked trees with $N(\tau) \geq 2$ and the strata of moduli spaces. The desired inversion formula now follows from the preceding corollary. \square

Proof of Corollary 2. Let $a_n = \chi(\mathcal{M}_{0,n+1})$. It is known that $\chi(X-Y) + \chi(Y) = \chi(X)$ whenever Y is a closed subvariety of a complex variety X , see [Ful], p. 141, note 13, and that $\chi(A \times B) = \chi(A) \times \chi(B)$. The first property implies that $\chi(\overline{\mathcal{M}}_{0,n+1})$ is the sum of the Euler characteristics of its strata S , and the second implies that

$$\chi(S) = a_{n_1} \cdots a_{n_s}$$

whenever $S \cong \mathcal{M}_{0,n_1+1} \times \cdots \times \mathcal{M}_{0,n_s+1}$. Thus the stated relationship between generating functions follows from Theorem 1. \square

Moduli space over \mathbb{R} . The real points of the moduli space form a submanifold $\mathcal{M}_{0,n}(\mathbb{R})$ with $(n-1)!/2$ connected components, each homeomorphic to \mathbb{R}^{n-3} . Let M_n be the component of $\mathcal{M}_{0,n}(\mathbb{R})$ where the marked points can be chosen to lie in \mathbb{R} , with $x_1 < x_2 < \cdots < x_n$. Let \overline{M}_n be the closure of M_n in $\overline{\mathcal{M}}_{0,n}$. The strata of \overline{M}_n are encoded by ribbon trees, since the cyclic ordering of the points (x_i) is preserved under stable limits (cf. [De]). Thus in this setting, Theorem 3 yields:

Corollary 5. *The formal inverse of $F(x) = x - \sum_2^\infty A_n x^n$ is given by $G(x) = x + \sum_2^\infty B_n x^n$, where*

$$B_n = \sum A_{n_1} \cdots A_{n_s} \times \left(\begin{array}{c} \text{the number of strata } S \subset \overline{M}_{0,n+1} \\ \text{isomorphic to } M_{n_1+1} \times \cdots \times M_{n_s+1} \end{array} \right).$$

Notes and references. A compendium of results on trees, generating functions and inversion can be found in [St], Chapter 5; see also [Ca]. For background on the many connections between graphs and moduli space, see e.g. [ACG], Chapter XVIII, [LZ], and the references therein.

Acknowledgements. I would like to thank S. Koch for useful conversations related to this note.

References

- [ACG] E. ARBARELLO, M. CORNALBA, and P.A. GRIFFITHS, *Geometry of Algebraic Curves*. Volume 2. With a contribution by J.D. Harris. Grundlehren der mathematischen Wissenschaften 268. Springer Verlag, Berlin etc., 2011.
MR 2807457 Zbl 1235.14002

- [Bi] F. BITTNER, The universal Euler characteristic for varieties of characteristic zero. *Compos. Math.* 140 (2004), 1011–1032. [MR 1086.14016](#) [Zbl 2059227](#)
- [Ca] D. CALLAN, Lagrange inversion and Schröder trees. Preprint 1999. www.stat.wisc.edu/~callan/notes/lagrange_schroder/lagrange_schroder.tex
- [De] S. L. DEVADOSS, Tessellations of moduli spaces and the mosaic operad. In J. P. MEYER, J. MORAVA and W. S. WILSON (eds.), *Homotopy Invariant Algebraic Structures*. Contemporary Mathematics 239. American Mathematical Society, Providence, RI, 1999, 91–114. [MR 1718078](#) [Zbl 0968.32009](#)
- [Ful] W. FULTON, *Introduction to Toric Varieties*. Annals of Mathematics Studies 131. Princeton University Press, Princeton, NJ, 1993. [MR 1234037](#) [Zbl 0813.14039](#)
- [Ge1] E. GETZLER, Operads and moduli spaces of genus 0 Riemann surfaces. In R. H. DIJKGRAAF, C. F. FABER, and G. VAN DER GEER (eds.), *The Moduli Space of Curves*. Progress in Mathematics 129. Birkhäuser Boston, Boston, MA, 1995, 199–230. [MR 1363058](#) [Zbl 0851.18005](#)
- [Ge2] —, The semi-classical approximation for modular operads. *Comm. Math. Phys.* **194** (1998), 481–492. [MR 1627677](#) [Zbl 0912.18007](#)
- [LZ] S. K. LANDO and A. K. ZVONKIN, *Graphs on Surfaces and Their Applications*. With an appendix by Don B. ZAGIER. Encyclopaedia of Mathematical Sciences 141. Low-Dimensional Topology II. Springer Verlag, Berlin, 2004. [MR 2036721](#) [Zbl 1040.05001](#)
- [St] R. P. STANLEY, *Enumerative Combinatorics*. Volume II. With a foreword by G.-C. ROTA and Appendix 1 by S. FOMIN. Cambridge Studies in Advanced Mathematics 62. Cambridge University Press, Cambridge, 1999. [MR 1676282](#) [Zbl 0928.05001](#)

(Reçu le 2 octobre 2012)

Curtis T. McMULLEN, Mathematics Department, Harvard University, 1 Oxford Street, Cambridge, MA 02138-2901, U.S.A.