

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 60 (2014)

Artikel: Schauder estimate for solutions of Poisson's equation with Neumann boundary condition
Autor: Nardi, Giacomo
DOI: <https://doi.org/10.5169/seals-630586>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 16.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Schauder estimate for solutions of Poisson's equation with Neumann boundary condition

Giacomo NARDI*

Abstract. We consider here the Neumann problem for the Laplace operator and prove an existence result in the Hölder spaces and obtain Schauder estimates. According to our knowledge this result is not explicitly proved in the several works devoted to Schauder theory, where similar theorems are proved for the Dirichlet and oblique derivative problems. Our contribution is to make explicit the existence and the estimate for the Neumann problem.

Mathematics Subject Classification (2010). Primary 35J25, 35J05

Keywords. Schauder estimate, Poisson's equation, Neumann boundary condition

1. Introduction

Let Ω be a $C^{2,\alpha}$ -domain of \mathbb{R}^N (we refer to Section 2 for notation and definitions). We consider the following problem:

$$\begin{cases} \Delta u = f & \text{in } \Omega, \\ \frac{\partial u}{\partial n} = g & \text{on } \partial\Omega, \end{cases}$$

with $f \in C^{0,\alpha}(\overline{\Omega})$ and $g \in C^{1,\alpha}(\overline{\Omega})$. The aim of this paper is to prove an existence result in $C^{2,\alpha}(\overline{\Omega})$ for this problem and an estimate of the form

$$\|u\|_{C^{2,\alpha}} \leq C(\|f\|_{C^{0,\alpha}} + \|g\|_{C^{1,\alpha}}).$$

In the 1930s, this kind of estimate was used by Schauder [Sch] and Caccioppoli [Ca] to prove an existence result in $C^{2,\alpha}(\overline{\Omega})$ for the Dirichlet problem for an elliptic equation ([GT, Theorem 6.8, p. 100]; [LU, Theorem 1.3, p. 107]). Using the same technique, in the 1950s, Fiorenza [Fi] proved a similar estimate and an existence result in $C^{2,\alpha}(\overline{\Omega})$ for the oblique derivative problem

* We thank A. Adimurthi, N. Fusco, R. Gianni, L. Orsina, and N. Trudinger, for their advice. We thank B. Dacorogna who supervised the researches on this subject.

$$l(x)u + m(x)\frac{\partial u}{\partial n} = g \quad \text{on } \partial\Omega \quad (lm > 0 \text{ on } \partial\Omega)$$

for elliptic equations ([GT, Theorem 6.31, p. 128]; [LU, Theorem 3.1, p. 126]).

Unfortunately the hypothesis $l \neq 0$ may not be removed in the proof of the existence result and the Schauder estimate for the oblique derivative problem ([GT, Theorem 6.31, p. 128]). Moreover, reading this proof, we can verify that it is not even possible to get the result for the Neumann problem taking the limit $l \rightarrow 0$.

So, the case of the Neumann problem needs to be considered independently and, according to our knowledge, it is not explicitly present in the classical literature on the subject (see for instance [GT], [LU]) where the Dirichlet and oblique derivative problems are studied in detail.

The main goal of this work is to formalize the existence result and a Schauder estimate for the Neumann problem for Poisson's equation.

Our main contribution is the following result (Theorems 3.1 and 4.1):

Theorem. *Let $\alpha \in (0, 1)$ and let Ω be a $C^{2,\alpha}$ -domain. Let $f \in C^{0,\alpha}(\overline{\Omega})$ and $g \in C^{1,\alpha}(\overline{\Omega})$ be such that*

$$\int_{\Omega} f = \int_{\partial\Omega} g.$$

Then there exists a solution $u \in C^{2,\alpha}(\overline{\Omega})$ (unique up to an additive constant) to the problem

$$\begin{cases} \Delta u = f & \text{in } \Omega, \\ \frac{\partial u}{\partial n} = g & \text{on } \partial\Omega. \end{cases}$$

Moreover, every solution to this problem verifies the estimate

$$\left\| u - \frac{1}{|\Omega|} \int_{\Omega} u \right\|_{C^{2,\alpha}} \leq C(\|f\|_{C^{0,\alpha}} + \|g\|_{C^{1,\alpha}}),$$

with $C = C(\Omega, N, \alpha)$.

The starting point of the proof is the alternative theorem for the oblique derivative problem for an uniformly elliptic operator L with coefficients in $C^{0,\alpha}(\overline{\Omega})$ and $c < 0$ (Theorem 5.1). This allows us to prove an existence result (uniqueness is given up to a constant) for the Neumann problem for elliptic operators with $c < 0$ (Theorem 5.2). Afterwards, using Theorem 5.2 and the Fredholm alternative, we can prove the existence and uniqueness of the solution to our initial problem in the class of functions belonging to $C^{2,\alpha}(\overline{\Omega})$ and having null average (Theorem 3.1).

Concerning the estimate we are led to estimate the quantity $u - \frac{1}{|\Omega|} \int u$ instead of u , because the solution to the Neumann problem, if it exists, is unique up to a constant. We give three proofs of this result (Theorem 4.1).

We finally obtain, for the Neumann problem, results similar to those for the Dirichlet and oblique derivative boundary conditions.

As already mentioned, the theorem is well known to all specialists in elliptic partial differential equations. However, after discussion with several of them, we were not able to find a precise reference for such a result, specially the one concerning the estimate.

This paper is organized as follows. In Section 2 we recall the main definitions used in this work. In Section 3 we prove an existence theorem for the Neumann problem for the Poisson equation (Theorem 3.1). In Section 4 we prove the estimate (Theorem 4.1). In Section 5 we recall some useful results to prove the existence theorem. We adopt the same notation as in [GT]. We refer to [Na] for a more detailed analysis of the problem.

2. Definitions and notation

In the sequel we denote by Ω an open bounded non-empty subset of \mathbb{R}^N ($N \geq 2$) and let u be a function defined on Ω . For every multi-index $\beta = (\beta_1, \dots, \beta_N)$ ($\beta_i \geq 0$ for $i = 1, \dots, N$) of length $|\beta| = \sum_{i=1}^N \beta_i$, we set

$$D^\beta u = \frac{\partial^{|\beta|} u}{\partial x_1^{\beta_1} \dots \partial x_N^{\beta_N}}.$$

We recall the definition of the usual functional spaces ($k \geq 0$):

$$C^k(\Omega) = \{u : \Omega \rightarrow \mathbb{R} \mid \forall \beta \text{ multi-index, } |\beta| \leq k, D^\beta u \text{ is continuous in } \Omega\},$$

$$C^k(\overline{\Omega}) = \{u \in C^k(\Omega) \mid D^\beta u, |\beta| \leq k, \text{ can be extended by continuity to } \partial\Omega\}.$$

Moreover $C^k(\overline{\Omega})$ is a Banach space equipped with the norm

$$\|u\|_{C^k} = \sum_{i=0}^k \sup_{|\beta|=i} \sup_{\Omega} |D^\beta u|.$$

We now recall the definition of Hölder spaces.

Definition 2.1 (Hölder-continuity). Let $\alpha \in]0, 1]$ and let $\Omega \subset \mathbb{R}^N$ be an open set. We define the α -Hölder coefficient of $u : \Omega \rightarrow \mathbb{R}$ as

$$[u]_{0,\alpha;\Omega} = \sup_{\substack{x,y \in \Omega \\ x \neq y}} \frac{|u(x) - u(y)|}{|x - y|^\alpha}.$$

If $[u]_{0,\alpha;\Omega} < \infty$ then we say that u is Hölder continuous with exponent α in Ω . If there is no ambiguity about the domain Ω we denote $[u]_{0,\alpha;\Omega}$ by $[u]_{0,\alpha}$.

We define the *Hölder space* $C^{k,\alpha}(\overline{\Omega})$ as the set of functions belonging to $C^k(\overline{\Omega})$ whose k th-order partial derivatives are Hölder continuous with exponent α in Ω .

$C^{k,\alpha}(\overline{\Omega})$ is a Banach space equipped with the following norm:

$$\|u\|_{C^{k,\alpha}} = \|u\|_{C^k} + [u]_{k,\alpha},$$

where

$$[u]_{k,\alpha} = \sup_{|\beta|=k} [D^\beta u]_{0,\alpha}.$$

We set $C^{k,0}(\overline{\Omega}) = C^k(\overline{\Omega})$, and one can easily verify that $C^{k,\alpha}(\overline{\Omega}) \subset C^{h,\alpha}(\overline{\Omega})$ for all integers h, k with $h < k$.

Moreover, for every vector-valued function $u : \Omega \rightarrow \mathbb{R}^N$, we say that it belongs to $C^{k,\alpha}(\overline{\Omega})$ if its components belong to $C^{k,\alpha}(\overline{\Omega})$.

In order to give the definition of $C^{k,\alpha}$ -domains, we need the following notation:

$$\begin{aligned} B &= \{x = (x_1, \dots, x_N) \in \mathbb{R}^N \mid |x| < 1\}, \\ B_+ &= \{x \in B \mid x_N > 0\}, \quad B_0 = \{x \in B \mid x_N = 0\}. \end{aligned}$$

Definition 2.2. We call *domain* every open, bounded, connected, and non-empty subset Ω of \mathbb{R}^N . Moreover, Ω is said to be a $C^{k,\alpha}$ -domain ($k \geq 1$, $\alpha \in [0, 1]$) if for every $p \in \partial\Omega$ there exists a neighborhood U_p of p in \mathbb{R}^N and a diffeomorphism $\varphi_p : B \rightarrow U_p$ such that

- (i) $\varphi_p \in C^{k,\alpha}(\overline{B})$ and $\varphi_p^{-1} \in C^{k,\alpha}(\overline{U_p})$;
- (ii) $\varphi_p(B_+) = U_p \cap \Omega$;
- (iii) $\varphi_p(B_0) = U_p \cap \partial\Omega$.

Remark 2.3 (Hölder-continuity on the boundary). Let Ω be a $C^{k,\alpha}$ -domain. We say that $u \in C^{k,\alpha}(\partial\Omega)$ if, for every $p \in \partial\Omega$, we have $u \circ \varphi_p \in C^{k,\alpha}(B_0)$, where φ_p is given by the previous definition.

Of course if $u \in C^{k,\alpha}(\overline{\Omega})$ its restriction to $\partial\Omega$ belongs to $C^{k,\alpha}(\partial\Omega)$. Moreover, for every $u \in C^{k,\alpha}(\partial\Omega)$ there exists a function belonging to $C^{k,\alpha}(\overline{\Omega})$ whose restriction to the boundary coincides with u (see [GT]: Lemma 6.38 p. 137).

Then, in order to study the Neumann problem for Poisson's equation, we can consider boundary values belonging to $C^{k,\alpha}(\overline{\Omega})$ instead of $C^{k,\alpha}(\partial\Omega)$.

We denote by L^p and $W^{m,p}$ the usual Lebesgue and Sobolev spaces and refer to [AF, Br] for their properties.

3. Existence of solutions to the Neumann problem for Poisson's equation in $C^{2,\alpha}(\bar{\Omega})$

In this section we consider Poisson's equation with Neumann boundary condition and prove the following result:

Theorem 3.1. *Let Ω be a $C^{2,\alpha}$ -domain and let $f \in C^{0,\alpha}(\bar{\Omega})$, $g \in C^{1,\alpha}(\bar{\Omega})$ be such that*

$$(3.1) \quad \int_{\Omega} f = \int_{\partial\Omega} g.$$

Then the problem

$$(3.2) \quad \begin{cases} \Delta u = f & \text{in } \Omega, \\ \frac{\partial u}{\partial n} = g & \text{on } \partial\Omega, \end{cases}$$

admits a unique solution in the class

$$\mathcal{C} = \left\{ u \in C^{2,\alpha}(\bar{\Omega}) : \frac{1}{|\Omega|} \int_{\Omega} u = 0 \right\}.$$

We start by recalling the following estimate which is very useful in the sequel:

Theorem 3.2 ([GT, Theorem 6.30, p. 127]; [LU, Theorem 3.1, p. 126]). *Let Ω be a $C^{2,\alpha}$ -domain and let $u \in C^{2,\alpha}(\bar{\Omega})$ be a solution of (3.2). Then*

$$(3.3) \quad \|u\|_{C^{2,\alpha}} \leq C(\|f\|_{C^{0,\alpha}} + \|g\|_{C^{1,\alpha}} + \|u\|_{C^0}),$$

with $C = C(\Omega, \alpha, N)$.

We prove two preliminary lemmas.

Lemma 3.3. *Let $\Omega \subset \mathbb{R}^N$ be a C^2 -domain and let $u \in C^2(\bar{\Omega})$. We suppose that there exists $p \in \partial\Omega$ such that:*

$$u(p) = \max_{\bar{\Omega}} u \quad (\min_{\bar{\Omega}} u) \quad \text{and} \quad Du(p) = 0.$$

Then $\Delta u(p) \leq 0$ (≥ 0).

Proof. Up to a translation, we may assume that p is the origin of \mathbb{R}^N . Moreover, as $\Omega \in C^2$, up to a rotation, we may also assume that, for some R , we have $-re_i \in \Omega$ for every $r \in [0, R]$ and $i = 1, \dots, N$ where (e_1, \dots, e_N) is an orthonormal basis of \mathbb{R}^N . Then, partial derivatives verify

$$D_{ii}u(p) = \lim_{h \rightarrow 0^-} \frac{D_i u(p + he_i) - D_i u(p)}{h} = \lim_{h \rightarrow 0^-} \frac{D_i u(p + he_i)}{h} \leq 0,$$

because $Du(p) = 0$ and, for $h < 0$ small enough we have $D_i u(p + he_i) \geq 0$.

If p is a minimum point the proof is similar. \square

Lemma 3.4. *Let $\Omega \subset \mathbb{R}^N$ be a C^2 -domain and $f \in C^{0,\alpha}(\overline{\Omega})$. Let $u \in C^{2,\alpha}(\overline{\Omega})$ be a solution to the problem*

$$\begin{cases} \Delta u - u = f & \text{in } \Omega, \\ \frac{\partial u}{\partial n} = 0 & \text{on } \partial\Omega; \end{cases}$$

then

$$\|u\|_{C^0} \leq \|f\|_{C^0}.$$

Proof. Let p be such that $|u(p)| = \max_{\overline{\Omega}} |u|$. We consider separately the following cases: $p \in \Omega$ and $p \in \partial\Omega$.

$p \in \Omega$. Let p be a maximum point for u then $u(p) \geq 0$ and $\Delta u(p) \leq 0$. As $\Delta u - u = f$ we get $f(p) = \Delta u(p) - u(p) \leq \Delta u(p) \leq 0$. Thus

$$\|u\|_{C^0} = u(p) = \Delta u(p) - f(p) \leq -f(p) \leq \|f\|_{C^0}.$$

If p is a minimum point for u ($u(p) \leq 0$) the proof is similar.

$p \in \partial\Omega$. If p is a maximum point for u then p is a maximum point for $u|_{\partial\Omega}$. So, for every vector τ tangent to $\partial\Omega$ at p we have:

$$\frac{\partial u}{\partial \tau}(p) = 0$$

and, using the hypothesis

$$\frac{\partial u}{\partial n}(p) = 0,$$

we get $Du(p) = 0$. Thanks to the previous lemma we have $\Delta u(p) \leq 0$ and, as $\Delta u - u = f$ on $\partial\Omega$, the lemma ensues. We can use the same arguments if p is a minimum point for u . \square

Now, we can prove the existence theorem for the Neumann problem for the Laplace operator:

Proof of Theorem 3.1. By Theorem 5.2 there exists a unique solution in $C^{2,\alpha}(\overline{\Omega})$, denoted by $\mathfrak{X}[f, g]$, to the problem:

$$\begin{cases} \Delta u - u = f & \text{in } \Omega, \\ \frac{\partial u}{\partial n} = g & \text{on } \partial\Omega, \end{cases}$$

for all f, g verifying the compatibility condition (3.1). Moreover

$$(3.4) \quad \int_{\Omega} u = \int_{\Omega} \Delta u - \int_{\Omega} f = \int_{\partial\Omega} \frac{\partial u}{\partial n} - \int_{\partial\Omega} g = 0,$$

so $\mathfrak{X}[f, g] \in \mathcal{C}$. Defining

$$\mathcal{A} = \left\{ (f, g) \in C^{0,\alpha}(\bar{\Omega}) \times C^{1,\alpha}(\bar{\Omega}) : \int_{\Omega} f = \int_{\partial\Omega} g \right\}$$

we have that $\mathfrak{L} : \mathcal{A} \rightarrow \mathcal{C}$ is a well defined bijective operator.

Now, we consider the following equation:

$$(3.5) \quad u - \mathfrak{L}[-u, 0] = \mathfrak{L}[f, g].$$

Then $u \in \mathcal{C}$ is a solution to the problem (3.2) if and only if it is a solution of (3.5), because

$$(f, g) = \mathfrak{L}^{-1}(u - \mathfrak{L}[-u, 0]) = \mathfrak{L}^{-1}u + (u, 0) = \left(\Delta u, \frac{\partial u}{\partial n} \right).$$

Then, we need to show that for every $(f, g) \in \mathcal{A}$ there exists a unique solution $u \in \mathcal{C}$ of (3.5). As \mathfrak{L} is bijective, we are led to prove that, for every $v \in \mathcal{C}$, the equation

$$(3.6) \quad u - \mathfrak{L}[-u, 0] = v$$

admits a unique solution on \mathcal{C} . For this we use the Fredholm alternative theorem.

We consider the space

$$\mathcal{F} = \left\{ f \in C^{0,\alpha}(\bar{\Omega}) : \frac{1}{|\Omega|} \int_{\Omega} f = 0 \right\},$$

equipped with the norm of $C^{0,\alpha}(\bar{\Omega})$. Let T be the operator:

$$\begin{aligned} T : \mathcal{F} &\rightarrow \mathcal{F}, \\ Tf &= \mathfrak{L}[-f, 0]. \end{aligned}$$

Using the properties of \mathfrak{L} , we get:

$$(3.7) \quad T(\mathcal{F}) \subset \mathcal{C}.$$

We first show that T is a compact operator. Let $\{f_k\} \subset \mathcal{F}$ then, because of Theorem 5.2 and (3.4), there exists $\{u_k\} \subset \mathcal{C}$ such that $Tf_k = u_k$ and, because of (3.3), we have

$$\|u_k\|_{C^{2,\alpha}} \leq C(\Omega, \alpha, N)(\|f_k\|_{C^{0,\alpha}} + \|u_k\|_{C^0}).$$

So, using the previous lemma, we get:

$$\|u_k\|_{C^{2,\alpha}} \leq 2C(\Omega, \alpha, N)\|f_k\|_{C^{0,\alpha}}.$$

If $\{f_k\}$ is a bounded sequence of \mathcal{F} then $\{u_k\}$ is bounded in $C^{2,\alpha}(\bar{\Omega})$ and in $W^{2,\infty}(\Omega)$. Thus there exists a subsequence $\{u_{k_h}\}$ and a function $u \in W^{2,\infty}(\Omega)$ such that

$$u_{k_h} \xrightarrow{*} u \text{ in } W^{2,\infty},$$

so, for every $p > 1$

$$u_{k_h} \xrightarrow{*} u \text{ in } W^{2,p}.$$

Now, choosing $\frac{N}{2-\alpha} < p \leq N$, by the Rellich–Kondrachov theorem, we obtain

$$u_{k_h} \rightarrow u \text{ in } C^{0,\alpha}(\bar{\Omega}),$$

which proves that T is compact.

Now, equation (3.6) can be rewritten as

$$(3.8) \quad u - Tu = v.$$

The equation

$$u - Tu = 0$$

is equivalent to the problem

$$\begin{cases} \Delta u = 0 & \text{in } \Omega, \\ \frac{\partial u}{\partial n} = 0 & \text{on } \partial\Omega, \end{cases}$$

which, in \mathcal{C} , admits only the trivial solution $u = 0$.

Then, as T is compact, applying the Fredholm alternative to (3.8) we have that for every $v \in \mathcal{C}$ there exists a unique solution $u \in \mathcal{C}$ of (3.8) and the theorem ensues. \square

Remark 3.5. If $u \in \mathcal{C}$ is a solution to the problem (3.2) then, for every $k \in \mathbb{R}$, the function $u + k$ is also a solution to (3.2) in $C^{2,\alpha}(\bar{\Omega})$ (but not in \mathcal{C}).

On the other hand, if u is a solution to (3.2) in $C^{2,\alpha}(\bar{\Omega})$, we can obtain a solution in \mathcal{C} setting

$$v = u - \frac{1}{|\Omega|} \int_{\Omega} u.$$

Then, using Theorem 3.1 we have an existence and uniqueness (up to an additive constant) result for the problem (3.2) in $C^{2,\alpha}(\bar{\Omega})$.

4. Schauder estimate

We prove an estimate of $\|u\|_{C^{2,\alpha}}$ in terms of $\|f\|_{C^{0,\alpha}}, \|g\|_{C^{1,\alpha}}$ for any solution to problem (3.2). In particular, as the uniqueness of the solution is proved up to an additive constant, we prove the estimate for a solution with null average.

We state the following theorem and give three proofs:

Theorem 4.1. Let Ω be a $C^{2,\alpha}$ -domain and $f \in C^{0,\alpha}(\overline{\Omega})$, $g \in C^{1,\alpha}(\overline{\Omega})$ be such that

$$\int_{\Omega} f = \int_{\partial\Omega} g.$$

Let $u \in C^{2,\alpha}(\overline{\Omega})$ be a solution to (3.2). Then

$$\left\| u - \frac{1}{|\Omega|} \int_{\Omega} u \right\|_{C^{2,\alpha}} \leq C(\|f\|_{C^{0,\alpha}} + \|g\|_{C^{1,\alpha}}),$$

with $C = C(\Omega, \alpha, N)$.

The first proof of Theorem 4.1 was suggested to us by A. Adimurthy who attributed it to Jacques-Louis Lions. However we did not find any reference for such a proof, which is detailed in the sequel.

First proof of Theorem 4.1. Let $u \in C^{2,\alpha}(\overline{\Omega})$ be a solution to (3.2) such that $\frac{1}{|\Omega|} \int_{\Omega} u = 0$. By (3.3), we have the following estimate for u :

$$(4.1) \quad \|u\|_{C^{2,\alpha}} \leq C_1(\Omega, \alpha, N)(\|u\|_{C^0} + \|f\|_{C^{0,\alpha}} + \|g\|_{C^{1,\alpha}}),$$

and we wish to prove that

$$(4.2) \quad \|u\|_{C^{2,\alpha}} \leq C_2(\Omega, \alpha, N)(\|f\|_{C^{0,\alpha}} + \|g\|_{C^{1,\alpha}}).$$

Let us suppose that (4.2) is false. Then for every $k \in \mathbb{N}$ there exist $\{u_k\} \in C^{2,\alpha}(\overline{\Omega})$ and $\{f_k\} \in C^{0,\alpha}(\overline{\Omega})$, $\{g_k\} \in C^{1,\alpha}(\overline{\Omega})$ such that

$$(4.3) \quad \begin{cases} \Delta u_k = f_k & \text{in } \Omega, \\ \frac{\partial u_k}{\partial n} = g_k & \text{on } \partial\Omega, \end{cases}$$

$$(4.4) \quad \frac{1}{|\Omega|} \int_{\Omega} u_k = 0,$$

$$(4.5) \quad \|u_k\|_{C^{2,\alpha}} = 1,$$

$$(4.6) \quad \|u_k\|_{C^{2,\alpha}} > k(\|f_k\|_{C^{0,\alpha}} + \|g_k\|_{C^{1,\alpha}}).$$

Thus, we have $f_k \rightarrow 0$ in $C^{0,\alpha}(\overline{\Omega})$ and $g_k \rightarrow 0$ in $C^{1,\alpha}(\overline{\Omega})$.

Using (4.5) we have, that for every multi-index β , $|\beta| = 0, 1, 2$, $\{D^{\beta} u_k\}$ is uniformly bounded in $C^0(\overline{\Omega})$ and equicontinuous because

$$|D^{\beta} u_k(x) - D^{\beta} u_k(y)| \leq |x - y|^{\alpha} \quad \forall x, y \in \Omega, \forall |\beta| = 2,$$

which implies that

$$|D^{\beta} u_k(x) - D^{\beta} u_k(y)| \leq C(\Omega) |x - y|^{\alpha} \quad \forall x, y \in \Omega, \forall |\beta| = 0, 1.$$

Iterating the Ascoli–Arzelà theorem we get a subsequence $\{u_{k_h}\}$ such that

$$(4.7) \quad \begin{aligned} u_{k_h} &\rightarrow u_0 \text{ in } C^0(\bar{\Omega}), \\ D^\beta u_{k_h} &\rightarrow u_\beta \text{ in } C^0(\bar{\Omega}) \quad \forall \beta, |\beta| = 1, 2, \end{aligned}$$

which implies that

$$u_{k_h} \rightarrow u_0 \text{ in } C^2(\bar{\Omega}).$$

Then

$$\begin{aligned} \Delta u_0 &= \lim_h \Delta u_{k_h} = \lim_h f_{k_h} = 0, \\ \frac{\partial u_0}{\partial n} &= \lim_h \frac{\partial u_{k_h}}{\partial n} = \lim_h g_{k_h} = 0, \end{aligned}$$

and

$$\begin{cases} \Delta u_0 = 0 & \text{in } \Omega, \\ \frac{\partial u_0}{\partial n} = 0 & \text{on } \partial\Omega, \\ \frac{1}{|\Omega|} \int_{\Omega} u_0 = 0, \end{cases}$$

which implies that $u_0 = 0$. Comparing with (4.1), we get a contradiction because

$$1 = \|u_{k_h}\|_{C^{2,\alpha}} \leq C_1(\Omega, \alpha, N) (\|u_{k_h}\|_{C^0} + \|f_{k_h}\|_{C^{0,\alpha}} + \|g_{k_h}\|_{C^{1,\alpha}}) \rightarrow 0.$$

□

For the second proof of Theorem 4.1 we need two more lemmas. The first one states an estimate in $L^2(\Omega)$ for $u - \frac{1}{|\Omega|} \int_{\Omega} u$:

Lemma 4.2. *Let Ω be a $C^{2,\alpha}$ -domain and let $f \in C^{0,\alpha}(\bar{\Omega})$, $g \in C^{1,\alpha}(\bar{\Omega})$ be such that*

$$\int_{\Omega} f = \int_{\partial\Omega} g.$$

Let $u \in C^{2,\alpha}(\bar{\Omega})$ be a solution to (3.2). Then

$$\left\| u - \frac{1}{|\Omega|} \int_{\Omega} u \right\|_{L^2} \leq C(\|f\|_{C^{0,\alpha}} + \|g\|_{C^{1,\alpha}}),$$

with $C = C(\Omega, N)$.

Proof. We may assume that $\frac{1}{|\Omega|} \int_{\Omega} u = 0$. Integrating by parts we have

$$\int_{\Omega} |Du|^2 = \int_{\partial\Omega} ug - \int_{\Omega} uf$$

and using Young's inequality with $\varepsilon > 0$ we get

$$\int_{\Omega} |Du|^2 \leq \int_{\partial\Omega} |ug| + \int_{\Omega} |uf| \leq \varepsilon \int_{\partial\Omega} |u|^2 + \frac{1}{4\varepsilon} \int_{\partial\Omega} |g|^2 + \varepsilon \int_{\Omega} |u|^2 + \frac{1}{4\varepsilon} \int_{\Omega} |f|^2.$$

Now, we have

$$\int_{\partial\Omega} |u|^2 \leq C_1(\Omega) \|u\|_{W^{1,2}}^2 \quad \forall u \in W^{1,2}(\Omega) \cap C(\bar{\Omega});$$

so

$$\int_{\Omega} |Du|^2 \leq \varepsilon C_2(\Omega) \|u\|_{W^{1,2}}^2 + C_3(\Omega, \varepsilon) [\|g\|_{W^{1,2}}^2 + \|f\|_{L^2}^2]$$

and, as $u \in \mathcal{C}$, using the Poincaré inequality, we get

$$\int_{\Omega} |Du|^2 \leq \varepsilon C_4(\Omega, N) \|Du\|_{L^2}^2 + C_5(\Omega, \varepsilon) [\|g\|_{C^{1,\alpha}}^2 + \|f\|_{C^{0,\alpha}}^2].$$

Choosing $\varepsilon < 1/C_4(\Omega, N)$ and using the Poincaré inequality we have

$$\|u\|_{L^2} \leq C_6(\Omega, N) \|Du\|_{L^2} \leq C (\|g\|_{C^{1,\alpha}} + \|f\|_{C^{0,\alpha}}),$$

with $C = C(\Omega, N)$. □

The second lemma proves a local estimate for a solution of Poisson's equation:

Lemma 4.3 ([Se, Theorem 1, p. 255, and Theorem 2, p. 259]). *Let Ω be a domain of \mathbb{R}^N . Let $f \in L^p(\Omega)$ with $p > N/2$ and $u \in C^2(\Omega)$ be a solution of $\Delta u = f$ in Ω . Then, for every ball $B(y, 2R) \subset \Omega$, we have*

$$\sup_{B(y, R)} |u| \leq C \left(R^{-\frac{N}{2}} \|u\|_{L^2(B(y, 2R))} + R^{2-\frac{N}{p}} \|f\|_{L^p(\Omega)} \right),$$

with $C = C(N, p)$.

Following an idea of N. Fusco, we can now give the second proof of Theorem 4.1.

Second proof of Theorem 4.1. We may assume that $\frac{1}{|\Omega|} \int_{\Omega} u = 0$. We give an estimate for $\sup_{\partial\Omega} |u|$. As Ω is a $C^{2,\alpha}$ -domain, for every small enough ε we have

$$(x - \varepsilon n(x)) \in \Omega \quad \forall x \in \partial\Omega$$

where $n(x)$ denotes the unit outer-pointing normal to $\partial\Omega$ at x . Then, by the Lagrange theorem, for every $x \in \partial\Omega$ there exists $\tau = \tau(x, \varepsilon) \in (0, 1)$ such that

$$u(x - \varepsilon n(x)) = u(x) + \left(Du(x - \tau \varepsilon n(x)), -\varepsilon n(x) \right).$$

Defining $\Omega_{\varepsilon} = \{x \in \Omega \mid \text{dist}(x, \partial\Omega) \geq \varepsilon\}$ and taking the supremum for $x \in \partial\Omega$ in the previous relationship we get:

$$(4.8) \quad \sup_{\partial\Omega} |u| \leq \sup_{\Omega_\varepsilon} |u| + \varepsilon \sup_{\Omega} |Du|.$$

Let $R < \varepsilon/2$. We consider a finite cover of Ω_ε , denoted by $\{B(y_i, R)\}_{i=1}^M$ with $M = M(\varepsilon, \Omega)$ ($B(y_i, R)$ denotes an open ball of radius R centered at y_i). As $f \in C^0(\overline{\Omega})$ then $f \in L^{N+1}(\Omega)$ and, by Lemma 4.3, we have

$$\sup_{\Omega_\varepsilon} |u| \leq \sum_{i=1}^M \sup_{B(y_i, R)} |u| \leq MC_1(N) \left(R^{-\frac{N}{2}} \|u\|_{L^2(B(y_i, 2R))} + R^{2-\frac{N}{N+1}} \|f\|_{L^{N+1}(\Omega)} \right)$$

which implies that

$$(4.9) \quad \sup_{\Omega_\varepsilon} |u| \leq C_2(\varepsilon, \Omega, N) (\|u\|_{L^2} + \|f\|_{L^{N+1}}).$$

By Theorem 3.7 in [GT, p. 36] and (4.8), we have

$$\sup_{\Omega} |u| \leq \sup_{\Omega_\varepsilon} |u| + \varepsilon \sup_{\Omega} |Du| + C_3(\Omega) \sup_{\Omega} |f|$$

and, by (4.9), we get

$$\sup_{\Omega} |u| \leq C_2(\varepsilon, \Omega, N) (\|u\|_{L^2} + \|f\|_{L^{N+1}}) + \varepsilon \sup_{\Omega} |Du| + C_3(\Omega) \sup_{\Omega} |f|.$$

Using Lemma 4.2, we get

$$\sup_{\Omega} |u| \leq C_4(\varepsilon, \Omega, N) (\|f\|_{C^{0,\alpha}} + \|g\|_{C^{1,\alpha}}) + \varepsilon \|u\|_{C^{2,\alpha}}$$

and, by (3.3), we have

$$\|u\|_{C^{2,\alpha}} \leq C_5(\Omega, \alpha, N, \varepsilon) (\|f\|_{C^{0,\alpha}} + \|g\|_{C^{1,\alpha}}) + \varepsilon C_6(\Omega, \alpha, N) \|u\|_{C^{2,\alpha}}.$$

Choosing $\varepsilon < 1/C_6(N, \Omega, \alpha)$ the theorem ensues. \square

We now give the third proof of Theorem 4.1. We thank R. Gianni who gave us some ideas for this proof.

Third proof of Theorem 4.1. We assume that $\int_{\Omega} u = 0$ and, because of (3.3), we just need to prove an estimate for $\|u\|_{C^0}$ in terms of $\|f\|_{C^{0,\alpha}}$ and $\|g\|_{C^{1,\alpha}}$.

Let $M = \max_{\overline{\Omega}} |u|$ and $p \in \overline{\Omega}$ be such that $|u(p)| = M$. Moreover, we suppose $M > (\|f\|_{C^{0,\alpha}} + \|g\|_{C^{1,\alpha}})$, otherwise there is nothing to prove. By (3.3), we have

$$(4.10) \quad \|Du\|_{C^0} \leq C_1 (\|f\|_{C^{0,\alpha}} + \|g\|_{C^{1,\alpha}} + \|u\|_{C^0}) = K,$$

with $C_1 = C_1(\Omega, \alpha, N)$. We distinguish two cases: $p \in \Omega$ and $p \in \partial\Omega$.

We suppose that $p \in \Omega$ and we denote by $B(p, r)$ the open ball of radius r centered at p . We consider in particular $r < r_0 = \text{dist}(p, \partial\Omega)$ so that such a ball is contained in Ω .

Now, we can prove that for every $r < r_0$, we have

$$|u| > M - rK, \quad \text{in } B(p, r).$$

In fact, for every $x \in B(p, r)$ we have

$$u(x) - u(p) = \int_0^1 \frac{d}{dt} u(p + t(x - p)) dt = \int_0^1 \langle Du(p + t(x - p)), x - p \rangle dt$$

so

$$|u(x)| \geq |u(p)| - Kr = M - Kr.$$

Choosing $r < \min\{1/4C_1, r_0\}$, for every $x \in B(p, r)$, we have:

$$\begin{aligned} |u(x)| \geq M - rK &= (1 - rC_1)M - rC_1(\|f\|_{C^{0,\alpha}} + \|g\|_{C^{1,\alpha}}) \\ &> \frac{3}{4}M - rC_1(\|f\|_{C^{0,\alpha}} + \|g\|_{C^{1,\alpha}}). \end{aligned}$$

Then, as $M > (\|f\|_{C^{0,\alpha}} + \|g\|_{C^{1,\alpha}})$ and $r < 1/4C_1$, we get

$$|u(x)| > \frac{3}{4}M - C_1rM > \frac{M}{2} \quad \forall x \in B(p, r)$$

and, denoting by ω_N the measure of the unit ball of \mathbb{R}^N , we get

$$\frac{M}{2} \frac{\omega_N^{1/2}}{2^{N/2}} r^{N/2} \leq \left(\int_{B(p,r)} |u|^2 \right)^{1/2} \leq \|u\|_{L^2}.$$

Then, by Lemma 4.2, we can state that

$$\|u\|_{C^0} \leq \frac{2^{\frac{N+2}{2}} r^{-\frac{N}{2}}}{\omega_N^{1/2}} C_2(\Omega, N) (\|f\|_{C^{0,\alpha}} + \|g\|_{C^{1,\alpha}}),$$

which implies that

$$\|u\|_{C^{2,\alpha}} \leq C(\|f\|_{C^{0,\alpha}} + \|g\|_{C^{1,\alpha}}),$$

with $C = C(\Omega, \alpha, N)$.

We suppose now that $p \in \partial\Omega$. As Ω is a C^2 -domain, it satisfies the interior sphere condition (see [GT, p. 33]). Then, there exists $r_0 > 0$ such that the open ball $B(p - rn(p), r)$ with $r < r_0$ is contained in Ω ($n(p)$ denotes the unit outer-pointing normal to the boundary at p).

Thus, as p is not the center of the ball, similarly to the previous case we obtain

$$|u| > M - 2rK \quad \text{in } B(p - rn(p), r) \quad r < r_0.$$

Then, by choosing $r < \min\{1/8C_1, r_0\}$, we get the result by the same argument used above applied to the ball $B(p - rn(p), r)$. \square

5. Appendix: A preliminary result

In this section we recall some results used in Section 3. We consider a $C^{2,\alpha}$ -domain Ω and the operator

$$L : C^{2,\alpha}(\bar{\Omega}) \rightarrow \mathbb{R},$$

$$Lu = \sum_{i,j=1}^N a_{ij}(x) \frac{\partial^2 u}{\partial x_i \partial x_j} + \sum_{i=1}^N b_i(x) \frac{\partial u}{\partial x_i} + c(x)u,$$

where $a_{ij}, b_i, c \in C^{0,\alpha}(\bar{\Omega})$ for every $i, j = 1, \dots, N$ and $c(x) \leq 0$ for every $x \in \bar{\Omega}$.

We say that L is elliptic if the matrix $A(x) = [a_{i,j}(x)]$ is positive definite for every $x \in \bar{\Omega}$. Moreover, denoting by $\lambda(x)$ the smallest eigenvalue of $A(x)$, we say that L is uniformly elliptic if there exists $\lambda_0 > 0$ such that $\lambda(x) \geq \lambda_0$ for every $x \in \bar{\Omega}$.

We refer to [GT, p. 130] for the following alternative result for the oblique derivative problem:

Theorem 5.1. *Let Ω be a $C^{2,\alpha}$ -domain. Let L be an elliptic operator, which is uniformly elliptic in Ω and with $C^{0,\alpha}(\bar{\Omega})$ -coefficients. Let $l, m \in C^{1,\alpha}(\bar{\Omega})$ be such that $m \neq 0$ for every $x \in \partial\Omega$. Then exactly one of the following holds:*

(1) *The homogeneous problem*

$$\begin{cases} Lu = 0 & \text{in } \Omega, \\ l(x)u + m(x)\frac{\partial u}{\partial n} = 0 & \text{on } \partial\Omega, \end{cases}$$

admits nontrivial solutions;

(2) *the homogeneous problem has only the trivial solution, in which case for every $f \in C^{0,\alpha}(\bar{\Omega})$, $g \in C^{1,\alpha}(\bar{\Omega})$ there exists a solution $u \in C^{2,\alpha}(\bar{\Omega})$ to the inhomogeneous problem*

$$\begin{cases} Lu = f & \text{in } \Omega, \\ l(x)u + m(x)\frac{\partial u}{\partial n} = g & \text{on } \partial\Omega. \end{cases}$$

We can now prove the following theorem:

Theorem 5.2. *Let Ω be $C^{2,\alpha}$ -domain. Then for every $f \in C^{0,\alpha}(\bar{\Omega})$ and $g \in C^{1,\alpha}(\bar{\Omega})$ there exists a unique solution $u \in C^{2,\alpha}(\bar{\Omega})$ to the problem*

$$\begin{cases} \Delta u - u = f & \text{in } \Omega, \\ \frac{\partial u}{\partial n} = g & \text{on } \partial\Omega. \end{cases}$$

Proof. The problem

$$\begin{cases} \Delta u - u = 0 & \text{in } \Omega, \\ \frac{\partial u}{\partial n} = 0 & \text{on } \partial\Omega, \end{cases}$$

admits only the trivial solution. Using Point 2 of Theorem 5.1 the result ensues. \square

References

- [AF] R. A. ADAMS and J. J. F. FOURNIER, *Sobolev Spaces*. Academic Press, Paris-New York-London, 2003 (Second Edition). [Zbl 1098.46001](#) [MR 2424078](#)
- [Br] H. BREZIS, *Analyse Fonctionnelle*. Masson, Paris-New York-Milan, 1987 (2^e tirage).
- [Ca] R. CACCIOPPOLI, Sulle equazioni ellittiche a derivate parziali con n variabili indipendenti. *Atti della Accademia Nazionale dei Lincei. Rendiconti Lincei: Matematica e applicazioni. Serie VI* **19** (1934), 83–89. [Zbl 0009.06802](#)
- [Fi] R. FIORENZA, Sui problemi di derivata obliqua per le equazioni ellittiche. *Ricerche di Matematica* **8** (1959), 83–110. [Zbl 0090.31404](#) [MR 0114995](#)
- [GT] D. GILBARG and N. S. TRUDINGER, Elliptic partial differential equations of second order. Springer-Verlag, Berlin-Heidelberg, 2001 (Reprint of the 1998 edition). [Zbl 1042.35002](#) [MR 1814364](#)
- [LU] O. A. LADYŽENSKAJA and N. N. URAL'CEVA, *Équations aux Dérivées Partielles de Type Elliptique*. Dunod, Paris, 1968. [MR 0239273](#)
- [Na] G. NARDI, Risolubilità nel senso classico e nel senso delle distribuzioni per alcuni problemi ellittici. Master thesis, 2007.
- [Sch] J. P. SCHAUDER, Über lineare elliptische Differentialgleichungen zweiter Ordnung. *Mathematische Zeitschrift* B **38** (1934), 257–282. [Zbl 0008.25502](#) [MR 1545448](#)
- [Se] J. SERRIN, Local behavior of solutions of quasi-linear equations. *Acta Mathematica* **111** (1964), 247–302. [Zbl 0128.09101](#) [MR 0170096](#)

(Reçu le 13 février 2014)

Giacomo NARDI, Centre de Recherches en Mathématiques de la Decision (CEREMADE), CNRS : UMR7534, Université Paris IX – Paris Dauphine
e-mail: nardi@ceremade.dauphine.fr