

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 61 (2015)
Heft: 1-2

Artikel: A functorial extension of the abelian Reidemeister torsions of three-manifolds
Autor: Florens, Vincent / Massuyeau, Gwénaél
DOI: <https://doi.org/10.5169/seals-630593>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A functorial extension of the abelian Reidemeister torsions of three-manifolds

Vincent FLORENS and Gwénaél MASSUYEAU

Abstract. Let \mathbb{F} be a field and let $G \subset \mathbb{F} \setminus \{0\}$ be a multiplicative subgroup. We consider the category Cob_G of 3-dimensional cobordisms equipped with a representation of their fundamental group in G , and the category $\text{Vect}_{\mathbb{F}, \pm G}$ of \mathbb{F} -linear maps defined up to multiplication by an element of $\pm G$. Using the elementary theory of Reidemeister torsions, we construct a “Reidemeister functor” from Cob_G to $\text{Vect}_{\mathbb{F}, \pm G}$. In particular, when the group G is free abelian and \mathbb{F} is the field of fractions of the group ring $\mathbb{Z}[G]$, we obtain a functorial formulation of an Alexander-type invariant introduced by Lescop for 3-manifolds with boundary; when G is trivial, the Reidemeister functor specializes to the TQFT developed by Frohman and Nicas to enclose the Alexander polynomial of knots. The study of the Reidemeister functor is carried out for any multiplicative subgroup $G \subset \mathbb{F} \setminus \{0\}$. We obtain a duality result and we show that the resulting projective representation of the monoid of homology cobordisms is equivalent to the Magnus representation combined with the relative Reidemeister torsion.

Mathematics Subject Classification (2010). Primary: 57M27, 57Q10.

Keywords. 3-manifold, cobordism, Reidemeister torsion, Alexander polynomial, TQFT

Contents

1	Introduction	162
2	The Alexander functor A	167
3	Alexander functor and knots	173
4	The Reidemeister functor R	176
5	Back to the Alexander functor	184
6	Reidemeister functor and knots	187
7	The monoid of homology cobordisms	191
8	Computations with Heegaard splittings	195

9 Duality	198
A A short review of combinatorial torsions	205
References	207

1. Introduction

Let Cob be the category of 3-dimensional cobordisms introduced by Crane and Yetter [CY], and whose definition we briefly recall. The objects of Cob are integers $g \geq 0$, and correspond to compact connected oriented surfaces F_g of genus g with one boundary component. Indeed, we fix for every $g \geq 0$ a model surface F_g whose boundary is identified with S^1 , and we also fix a base point \star on $\partial F_g = S^1$. The morphisms $g_- \rightarrow g_+$ in the category Cob are the equivalence classes of cobordisms between the surfaces F_{g_-} and F_{g_+} . To be more specific, a *cobordism* from F_{g_-} to F_{g_+} is a pair (M, m) consisting of a compact connected oriented 3-manifold M and an orientation-preserving homeomorphism $m : F(g_-, g_+) \rightarrow \partial M$ where

$$F(g_-, g_+) := -F_{g_-} \cup_{S^1 \times \{-1\}} (S^1 \times [-1, 1]) \cup_{S^1 \times \{1\}} F_{g_+};$$

two such pairs (M, m) and (M', m') are *equivalent* if there exists a homeomorphism $f : M \rightarrow M'$ such that $m' = f|_{\partial M} \circ m$. We shall denote a pair (M, m) simply by the upper-case letter M , with the convention that the boundary-parametrization is always denoted by the lower-case letter m ; besides, we denote by $m_{\pm} : F_{g_{\pm}} \rightarrow M$ the restriction of m composed with the inclusion of ∂M into M . Thus the cobordism M “runs” from the *bottom surface* $\partial_- M := m_-(F_{g_-})$ to the *top surface* $\partial_+ M := m_+(F_{g_+})$. The *degree* of the cobordism M is the integer $g_+ - g_-$.

The composition $N \circ M$ of two cobordisms M, N in Cob is defined by identifying $\partial_+ M$ to $\partial_- N$ and, for any integer $g \geq 0$, the identity of the object g is the cylinder $F_g \times [-1, 1]$ with the obvious boundary-parametrization. Our model surfaces F_0, F_1, F_2, \dots also come with an identification of the boundary-connected sum $F_g \#_{\partial} F_h$ with the surface F_{g+h} for any $g, h \geq 0$. Thus the category Cob is enriched with a monoidal structure \otimes : the tensor product $g \otimes h$ of two integers g, h is the sum $g + h$, and the tensor product $M \otimes N$ of two cobordisms M, N is their boundary-connected sum $M \#_{\partial} N$.

Let now G be an abelian group. The category Cob can be refined to the category Cob_G of cobordisms equipped with a representation of the first integral homology group in G . To be more specific, an object of Cob_G is a pair (g, φ) consisting of an integer $g \geq 0$ and a group homomorphism $\varphi : H_1(F_g; \mathbb{Z}) \rightarrow G$. A morphism $(g_-, \varphi_-) \rightarrow (g_+, \varphi_+)$ in the category Cob_G

is a pair (M, φ) where $M \in \text{Cob}(g_-, g_+)$ and $\varphi : H_1(M; \mathbb{Z}) \rightarrow G$ is a group homomorphism such that $\varphi \circ m_{\pm, * } = \varphi_{\pm}$. The composition of two morphisms $(M, \varphi) \in \text{Cob}_G((g_-, \varphi_-), (g_+, \varphi_+))$ and $(N, \psi) \in \text{Cob}_G((h_-, \psi_-), (h_+, \psi_+))$, such that $(g_+, \varphi_+) = (h_-, \psi_-)$, is defined by

$$(N, \psi) \circ (M, \varphi) := (N \circ M, \psi + \varphi)$$

where $N \circ M$ is the composition in Cob and $\psi + \varphi : H_1(N \circ M; \mathbb{Z}) \rightarrow G$ is defined from φ and ψ by using the Mayer–Vietoris theorem. The monoidal structure of Cob also extends to the category Cob_G : the tensor product of objects is

$$(g, \varphi) \otimes (h, \psi) := (g + h, \varphi \oplus \psi)$$

where $H_1(F_{g+h}; \mathbb{Z}) = H_1(F_g \#_{\partial} F_h; \mathbb{Z})$ is identified with $H_1(F_g; \mathbb{Z}) \oplus H_1(F_h; \mathbb{Z})$, and the tensor product of morphisms is

$$(M, \varphi) \otimes (N, \psi) := (M \#_{\partial} N, \varphi \oplus \psi)$$

where $H_1(M \#_{\partial} N; \mathbb{Z})$ is identified with $H_1(M; \mathbb{Z}) \oplus H_1(N; \mathbb{Z})$.

Consider now a commutative ring R and fix a subgroup $G \subset R^\times$ of its group of units. Let $\text{grMod}_{R, \pm G}$ be the category whose objects are \mathbb{Z} -graded R -modules and whose morphisms are graded R -linear maps of arbitrary degree, up to multiplication by an element of $\pm G$. The usual tensor product of graded R -modules defines a monoidal structure on the category $\text{grMod}_{R, \pm G}$: here the tensor product $a \otimes b$ of two graded R -linear maps $a : U \rightarrow U'$ and $b : V \rightarrow V'$ is defined with Koszul’s rule, i.e. we set $(a \otimes b)(u \otimes v) := (-1)^{|b||u|} a(u) \otimes b(v)$ for any homogeneous elements $u \in U, v \in V$. In this paper, we construct and study two functors from Cob_G to $\text{grMod}_{R, \pm G}$ for some specific rings R and specific subgroups $G \subset R^\times$.

Our first functor is based on the “Alexander function” introduced by Lescop [Les]. For any compact orientable 3-manifold M with boundary, this function is defined on an exterior power of the Alexander module of M relative to a boundary point, and it takes values in a ring of Laurent polynomials. Lescop’s definition proceeds in a rather elementary way using a presentation of the Alexander module.

Theorem I. *Let G be a finitely generated free abelian group, and let $\mathbb{Z}[G]$ be its group ring. Then there is a degree-preserving monoidal functor*

$$A := A_G : \text{Cob}_G \longrightarrow \text{grMod}_{\mathbb{Z}[G], \pm G}$$

which, at the level of objects, assigns to any (g, φ) the exterior algebra of the φ -twisted relative homology group of the pair (F_g, \star) .

The $\mathbb{Z}[G]$ -linear map $A(M, \varphi)$ associated to a morphism (M, φ) of Cob_G is defined in a very simple way from the Alexander function of M using the

decomposition of ∂M into two parts, $\partial_- M$ and $\partial_+ M$. The fact that the Alexander function gives rise to a functor on the category of cobordisms is somehow implicit in [Les], where Lescop studies the behaviour of her invariant under some specific gluing operations. As it contains the Alexander polynomial of knots in a natural way, we call A the *Alexander functor*.

Since the works of Milnor [Mil1] and Turaev [Tur1], it is known that the Alexander polynomial of knots and 3-manifolds can be interpreted as a special kind of abelian Reidemeister torsion. We follow this direction to define our second functor, which we call the *Reidemeister functor*. In the sequel, the category $\text{grMod}_{R, \pm G}$ associated to a field $R := \mathbb{F}$ and a subgroup G of $\mathbb{F}^\times = \mathbb{F} \setminus \{0\}$ is denoted by $\text{grVect}_{\mathbb{F}, \pm G}$.

Theorem II. *Let \mathbb{F} be a field and let G be a subgroup of \mathbb{F}^\times . Then there is a degree-preserving monoidal functor*

$$R := R_{\mathbb{F}, G} : \text{Cob}_G \longrightarrow \text{grVect}_{\mathbb{F}, \pm G}$$

which, at the level of objects, assigns to any (g, φ) the exterior algebra of the φ -twisted relative homology group of the pair (F_g, \star) .

The construction of the functor R uses the elementary theory of Reidemeister torsions, but note that we need to consider cell chain complexes which are *not* necessarily acyclic. When G is a finitely generated free abelian group and $\mathbb{F} := Q(G)$ is the field of fractions of $\mathbb{Z}[G]$, we recover the functor A by extension of scalars. Thus it suffices to study the functor R and this is done using basic properties of combinatorial torsions. For instance, we compute its restriction to the monoid of homology cobordisms (which includes the mapping class group of a surface): we find that the representation induced by R is equivalent to the Magnus representation combined with the Reidemeister torsion of cobordisms relative to the top surface. We also give a formula for R in terms of Heegaard splittings and we show that R satisfies some duality properties, which generalize the symmetry properties of the Alexander polynomial of knots and 3-manifolds.

It is expected that Turaev's refinements of the Reidemeister torsion [Tur2, Tur3] can be adapted to refine R to a kind of "monoidal" degree-preserving functor from Cob_G to the category $\text{grVect}_{\mathbb{F}}$ of graded \mathbb{F} -vector spaces: the sign ambiguity would presumably be fixed using homological orientations on the manifolds, while the ambiguity in G would be dispelled by adding Euler structures. (Observe however that, since we use Koszul's rule and we allow morphisms in $\text{grVect}_{\mathbb{F}}$ to have non-zero degree, this category is not monoidal in the usual sense of the word.)

We now explain how our constructions are related to prior work. Soon after the emergence of quantum invariants of 3-manifolds in the late eighties, there were several works which showed how to interpret the classical Alexander polynomial

in this new framework. A more general problem was then to extend the Alexander polynomial to a functor from a category of cobordisms to a category of vector spaces following, as closely as possible, the axioms of a TQFT [Aty]. This problem was solved by Frohman and Nicas who used elementary intersection theory in $U(1)$ -representation varieties of surfaces [FN1]. (See also [FN2] for a much more general construction using $PU(N)$ -representations.) Later, Kerler showed that the Frohman–Nicas functor is in fact equivalent to a TQFT based on a certain quasitriangular Hopf algebra [Ker1]. The Alexander polynomial of a knot K in an integral homology 3-sphere N is recovered from this functor by taking the “graded” trace of the endomorphism associated to the cobordism that one obtains by “cutting” $N \setminus K$ along a Seifert surface of K . It turns out that, in the case $G = \{1\}$, the Alexander functor A is equivalent to the Frohman–Nicas functor. Note that the way how their functor determines the Alexander polynomial is somehow *extrinsic*, in that it goes through the choice of a Seifert surface. On the contrary, the functor A for $G = \mathbb{Z}$ *intrinsically* contains the Alexander polynomial of oriented knots in oriented integral homology 3-spheres by considering any knot of this type as a “bottom knot” in the manner of [Hab], i.e. by regarding its exterior as a morphism $1 \rightarrow 0$ in Cob_G . Since this functorial extension of the Alexander polynomial applies to cobordisms M equipped with an element of $H^1(M; \mathbb{Z})$, it should be regarded as a kind of HQFT with target $K(\mathbb{Z}, 1)$ – see [Tur6] – rather than a TQFT.

Our constructions are also related to the work of Bigelow, Cattabriga and the first author [BCF], which provides a functorial extension of the Alexander polynomial to the category of tangles instead of the category of cobordisms. To describe this relation, let TangCob be the monoidal category whose objects are pairs of non-negative integers (g, n) – corresponding to surfaces F_g with n punctures – and whose morphisms are cobordisms with tangles inside. Clearly the category TangCob contains the category Cob of [CY] as well as the usual category Tang of (unoriented) tangles in the standard ball; for any abelian group G , there is an obvious refinement TangCob_G of the category TangCob . When G is the infinite cyclic group generated by t , the usual category Tang_+ of oriented tangles in the standard ball can be regarded as a subcategory of TangCob_G by only considering those representations of tangle exteriors that send any oriented meridian to the generator t . The functors A and R constructed in this paper could be extended to the category TangCob_G using similar methods, but with more technicality. When G is infinite cyclic, the restriction of the resulting functor $A : \text{TangCob}_G \rightarrow \text{grMod}_{\mathbb{Z}[G], \pm G}$ to Tang_+ would coincide with the “Alexander representation of tangles” constructed in [BCF]. We also mention Archibald’s extension of the Alexander polynomial [Arc], which is based on diagrammatic presentations of tangles: her invariant seems to be very close to the invariant

constructed in [BCF] and is stronger since it is defined *without* ambiguity in $\pm G$.

Finally, our approach is related to the work of Cimasoni and Turaev on “Lagrangian representations of tangles” [CT1, CT2]. These representations are functors from the category Tang_+ to the category of “Lagrangian relations” (which generalizes the category of $\mathbb{Z}[t^{\pm 1}]$ -modules equipped with non-degenerate skew-hermitian forms) and, for string links, they are equivalent to the (reduced) Burau representation [LDi, KLW]. The constructions of [CT1, CT2] could be adapted to the case of cobordisms in order to obtain a functor from Cob_G to the category of “Lagrangian relations” over the ring $\mathbb{Z}[G]$. In the case of homology cobordisms, the resulting functor would be equivalent to the (reduced) Magnus representation but it would miss the relative Reidemeister torsion: so it would be weaker than the functor A .

The paper is organized as follows. A first part deals exclusively with the Alexander functor: §2 gives the construction of the functor A (Theorem I) and §3 explains how the classical Alexander polynomial of knots is contained in A . Next, the Reidemeister functor is constructed in §4 (Theorem II) and is proved to be a generalization of A in §5. (Thus, we provide two different proofs of the functoriality of A .) Starting from there, we focus on the study of R and indicate the resulting properties for A . The abelian Reidemeister torsions of knot exteriors and closed 3-manifolds are shown to be determined by R in §6. The functor R restricts to a projective representation of the monoid of homology cobordisms, which we fully compute in §7. We also explain in §8 how to calculate R using Heegaard splittings of cobordisms, and we prove in §9 a duality result for R which involves the twisted intersection form of surfaces. Finally, the paper ends with a short appendix recalling the definition and basic properties of the torsion of chain complexes.

Notation and conventions. Let R be a commutative ring. The exterior algebra of an R -module N is denoted by

$$\Lambda N = \bigoplus_{i \geq 0} \Lambda^i N \quad \text{where } \Lambda^0 N = R;$$

the multivector $v_1 \wedge \cdots \wedge v_i \in \Lambda^i N$ defined by a finite family $v = (v_1, \dots, v_i)$ of elements of N is still denoted by v . If N is free of rank n , a *volume form* on N is an isomorphism of R -modules $\Lambda^n N \rightarrow R$.

Let X be a topological space with base point \star . The maximal abelian cover of X based at \star is denoted by $p_X : \widehat{X} \rightarrow X$, and the preferred lift of \star is denoted by $\widehat{\star}$. (Here we assume the appropriate assumptions on X to have a universal cover.) For any oriented loop α in X based at \star , the unique lift of α to \widehat{X} starting at $\widehat{\star}$ is denoted by $\widehat{\alpha}$.

Unless otherwise specified, (co)homology groups are taken with coefficients in the ring of integers \mathbb{Z} ; (co)homology classes are denoted with square brackets $[-]$. For any subspace $Y \subset X$ such that $\star \in Y$ and any ring homomorphism $\varphi : \mathbb{Z}[H_1(X)] \rightarrow R$, we denote by $H^\varphi(X, Y)$ the φ -twisted homology of the pair (X, Y) , namely

$$H^\varphi(X, Y) = H(C^\varphi(X, Y)) \quad \text{where } C^\varphi(X, Y) := R \otimes_{\mathbb{Z}[H_1(X)]} C(\widehat{X}, p_X^{-1}(Y)).$$

If (X', Y') is another pair of spaces and $f : (X', Y') \rightarrow (X, Y)$ is a continuous map, the corresponding homomorphism $H(X') \rightarrow H(X)$ is still denoted by f . If a base point $\star' \in Y'$ is given and $f(\star') = \star$, the R -linear map $H^{\varphi f}(X', Y') \rightarrow H^\varphi(X, Y)$ induced by f is also denoted by f .

Acknowledgements. This work was partially supported by the French ANR research project ‘‘Interlow’’ (ANR-09-JCJC-0097-01). The authors would like to thank the referee for some useful comments.

2. The Alexander functor \mathbf{A}

We firstly review the Alexander function of a 3-manifold with boundary following [Les]. (Note that the terminology ‘‘Alexander function’’ has a very different meaning in [Tur2].) Next, we construct the Alexander functor \mathbf{A} . In this section, we fix a finitely generated free abelian group G ; the extension of a group homomorphism $\varphi : A \rightarrow G$ to a ring homomorphism $\mathbb{Z}[A] \rightarrow \mathbb{Z}[G]$ is still denoted by φ .

2.1. The Alexander function. Let M be a compact connected orientable 3-manifold with connected boundary. We fix a base point $\star \in \partial M$ and a group homomorphism $\varphi : H_1(M) \rightarrow G$. The *genus* of M is the integer $g(M) := 1 - \chi(M)$, i.e. the genus of the surface ∂M .

Lemma 2.1. *There exists a presentation of the $\mathbb{Z}[G]$ -module $H_1^\varphi(M, \star)$ whose deficiency is $g(M)$.*

Proof. We consider a decomposition of M with a single 0-handle, s 1-handles and r 2-handles. Since the boundary of M has genus $g(M)$, we have $s - r = g(M)$. This handle decomposition defines a 2-dimensional complex $X \subset M$ onto which M deformation retracts. The complex X has a single 0-cell (which we assume to be \star), s 1-cells and r 2-cells. Thus we obtain a presentation of the $\mathbb{Z}[G]$ -module $H_1^\varphi(M, \star) \simeq H_1^\varphi(X, \star)$ with s generators and r relations. □

We now simplify our notation by setting $g := g(M)$ and $H := H_1^\varphi(M, \star)$.

Definition 2.2 (Lescop [Les]). Consider a presentation of the $\mathbb{Z}[G]$ -module H with deficiency g :

$$(2.1) \quad H = \langle \gamma_1, \dots, \gamma_{g+r} \mid \rho_1, \dots, \rho_r \rangle.$$

Let Γ be the $\mathbb{Z}[G]$ -module freely generated by the symbols $\gamma_1, \dots, \gamma_{g+r}$, and regard ρ_1, \dots, ρ_r as elements of Γ . Then the *Alexander function* of M with coefficients φ is the $\mathbb{Z}[G]$ -linear map $\mathcal{A}_M^\varphi : \Lambda^g H \rightarrow \mathbb{Z}[G]$ defined by

$$\mathcal{A}_M^\varphi(u_1 \wedge \dots \wedge u_g) \cdot \gamma_1 \wedge \dots \wedge \gamma_{g+r} = \rho_1 \wedge \dots \wedge \rho_r \wedge \widetilde{u}_1 \wedge \dots \wedge \widetilde{u}_g \in \Lambda^{g+r} \Gamma$$

for any $u_1, \dots, u_g \in H$, which we lift to some $\widetilde{u}_1, \dots, \widetilde{u}_g \in \Gamma$ in an arbitrary way.

The map \mathcal{A}_M^φ can be concretely computed as follows: if one considers the $r \times (g+r)$ matrix defined by the presentation (2.1) of H , and if one adjoins to this matrix some row vectors giving u_1, \dots, u_g in the generators $\gamma_1, \dots, \gamma_{g+r}$, then $\mathcal{A}_M^\varphi(u_1 \wedge \dots \wedge u_g)$ is the determinant of the resulting $(g+r) \times (g+r)$ matrix. It is shown in [Les, §3.1] that, up to multiplication by a unit of $\mathbb{Z}[G]$ (i.e., an element of $\pm G$), the map \mathcal{A}_M^φ does not depend on the choice of the presentation (2.1).

Let $Q(G)$ be the field of fractions of $\mathbb{Z}[G]$. The following lemma, which is implicit in [Les], shows that either the Alexander function is trivial or it induces by extension of scalars a volume form on $H_Q := Q(G) \otimes_{\mathbb{Z}[G]} H$.

Lemma 2.3. *We have $\dim H_Q \geq g$, and $\mathcal{A}_M^\varphi \neq 0$ if and only if $\dim H_Q = g$.*

Proof. Let A be the $r \times (g+r)$ matrix with entries in $\mathbb{Z}[G]$ corresponding to the presentation (2.1) of the $\mathbb{Z}[G]$ -module H . The multiplication $v \mapsto vA$ defines a linear map $Q(G)^r \rightarrow Q(G)^{g+r}$ whose cokernel is H_Q . Therefore

$$\dim H_Q = (g+r) - \text{rank } A.$$

Clearly, we have $\text{rank } A \leq r$ so that $\dim H_Q \geq g$.

Assume that $\dim H_Q > g$ and let A' be a matrix obtained by adding g arbitrary rows to A . Then $\text{rank } A' < r$ so that all the minors of A' of order r vanish. By expanding the determinant of A' successively along the last g rows, we see that $\det A' = 0$ and deduce that $\mathcal{A}_M^\varphi = 0$.

Assume that $\dim H_Q = g$. Then $\text{rank } A = r$ so that A has a non-zero minor D of order r . Let $1 \leq i_1 < \dots < i_g \leq g+r$ be the indices of the columns of A not pertaining to D . Then $\mathcal{A}_M^\varphi(\gamma_{i_1} \wedge \dots \wedge \gamma_{i_g}) = D \neq 0$. \square

2.2. Definition of A. In order to define a functor A, we associate to any object (g, φ) of Cob_G the exterior algebra

$$A(g, \varphi) := \Lambda H_1^\varphi(F_g, \star)$$

of the $\mathbb{Z}[G]$ -module $H^\varphi(F_g, \star) = H_1^\varphi(F_g, \star)$, which is free of rank $2g$. Next, we associate to any morphism $(M, \varphi) \in \text{Cob}_G((g_-, \varphi_-), (g_+, \varphi_+))$ a $\mathbb{Z}[G]$ -linear map

$$A(M, \varphi) : \Lambda H_1^{\varphi_-}(F_{g_-}, \star) \longrightarrow \Lambda H_1^{\varphi_+}(F_{g_+}, \star)$$

of degree $\delta g := g_+ - g_-$ as follows. We denote by I the interval $m(\star \times [-1, 1])$, which connects the base point of the bottom surface $\partial_- M$ to that of the top surface $\partial_+ M$. We set $H := H_1^\varphi(M, I)$, $H_\pm := H_1^{\varphi_\pm}(F_{g_\pm}, \star)$ and $g := g_+ + g_-$. Then, for any integer $j \geq 0$, the image $A(M, \varphi)(x) \in \Lambda^{j+\delta g} H_+$ of any $x \in \Lambda^j H_-$ is defined by the following property:

$$\forall y \in \Lambda^{g-j} H_+, \mathcal{A}_M^\varphi(\Lambda^j m_-(x) \wedge \Lambda^{g-j} m_+(y)) = \omega(A(M, \varphi)(x) \wedge y).$$

Here $\omega : \Lambda^{2g} H_+ \rightarrow \mathbb{Z}[G]$ is an arbitrary volume form on H_+ . Due to the choices of ω and of the presentation of H , the map $A(M, \varphi)$ is only defined up to multiplication by an element of $\pm G$. Besides, observe that $A(M, \varphi)$ is trivial on $\Lambda^j H_-$ for any $j < \max(0, -\delta g)$ and any $j > \min(g, 2g_-)$.

The next two lemmas show that the above paragraph defines a monoidal functor A from Cob_G to $\text{grMod}_{\mathbb{Z}[G], \pm G}$, which proves Theorem I of the Introduction. The first lemma is related to Property 6 of the Alexander function in [Les], while the second lemma seems to be new.

Lemma 2.4. *For any morphisms $(M, \varphi) \in \text{Cob}_G((g_-, \varphi_-), (g_+, \varphi_+))$ and $(N, \psi) \in \text{Cob}_G((h_-, \psi_-), (h_+, \psi_+))$, we have*

$$(2.2) \quad A((M, \varphi) \otimes (N, \psi)) = A(M, \varphi) \otimes A(N, \psi).$$

Proof. We set $g := g_+ + g_-$, $h := h_+ + h_-$, $\delta g := g_+ - g_-$, $\delta h := h_+ - h_-$ and

$$H_\pm^M := H_1^{\varphi_\pm}(F_{g_\pm}, \star), \quad H_\pm^N := H_1^{\psi_\pm}(F_{h_\pm}, \star), \quad H_\pm := H_1^{\varphi_\pm \oplus \psi_\pm}(F_{g_\pm + h_\pm}, \star),$$

$$H^M := H_1^\varphi(M, I), \quad H^N := H_1^\psi(N, I), \quad H := H_1^{\varphi \oplus \psi}(M \sharp_\partial N, I).$$

In the statement of the lemma and in the proof below, we identify

$$A((g_\pm, \varphi_\pm) \otimes (h_\pm, \psi_\pm)) = A(g_\pm + h_\pm, \varphi_\pm \oplus \psi_\pm) = \Lambda H_\pm = \Lambda(H_\pm^M \oplus H_\pm^N)$$

in the obvious way with

$$\Lambda H_\pm^M \otimes \Lambda H_\pm^N = A(g_\pm, \varphi_\pm) \otimes A(h_\pm, \psi_\pm).$$

Since the intersection of M and N in $M \#_{\partial} N$ is a 2-disk which retracts onto I , the Mayer–Vietoris theorem gives an isomorphism $H^M \oplus H^N \xrightarrow{\cong} H$. If $\text{rank } H^M > g$, then $\mathcal{A}_M^\varphi = 0$ by Lemma 2.3 so that $A(M, \varphi) = 0$; the same lemma applied to N shows that

$$\text{rank } H = \text{rank } H^M + \text{rank } H^N > g + h$$

so that $A((M, \varphi) \otimes (N, \psi)) = 0$ and (2.2) trivially holds true. Therefore, we can assume in the sequel that $\text{rank}(H^M) = g$ and $\text{rank}(H^N) = h$.

Let $x := x^M \otimes x^N \in \Lambda^i H_-^M \otimes \Lambda^j H_-^N \subset \Lambda^{i+j} H_-$: we aim at showing that $a := A((M, \varphi) \otimes (N, \psi))(x)$ is equal to

$$a' := (A(M, \varphi) \otimes A(N, \psi))(x) = (-1)^{i\delta h} A(M, \varphi)(x^M) \otimes A(N, \psi)(x^N).$$

(Recall that we are using Koszul’s rule in the definition of the tensor product of morphisms in the category $\text{grMod}_{\mathbb{Z}[G], \pm G}$.) It is enough to prove that, for any integers $p, q \geq 0$ such that $p + q = (g + h) - (i + j)$ and any $y := y^M \otimes y^N \in \Lambda^p H_+^M \otimes \Lambda^q H_+^N \subset \Lambda^{p+q} H_+$, the identity

$$(2.3) \quad \omega(a \wedge y) = \omega(a' \wedge y)$$

holds true up to multiplication by an element of $\pm G$ independent of x, y (and, in particular, independent of i, j, p, q). In the sequel, we fix some volume forms ω^M and ω^N on H_+^M and H_+^N respectively, and we assume that the volume form ω on $H_+ = H_+^M \oplus H_+^N$ is defined by

$$(2.4) \quad \omega(u \wedge v) = \omega^M(u) \cdot \omega^N(v)$$

for any $u \in \Lambda^{2g} H_+^M$ and $v \in \Lambda^{2h} H_+^N$. By definition of A , we have

$$(2.5) \quad \omega(a \wedge y) = \mathcal{A}_{M \#_{\partial} N}^{\varphi \oplus \psi} (\Lambda^i m_-(x^M) \wedge \Lambda^j n_-(x^N) \wedge \Lambda^p m_+(y^M) \wedge \Lambda^q n_+(y^N)).$$

If $p > g - i$, then $i + p > \text{rank}(H^M)$ by our assumptions, so that $\Lambda^i m_-(x^M) \wedge \Lambda^p m_+(y^M) \in \Lambda^{i+p} H^M$ is torsion; we deduce that $\omega(a \wedge y) = 0$; on the other hand, the degree of $A(M, \varphi)(x^M) \wedge y^M \in \Lambda H_+^M$ is $i + \delta g + p > 2g$ so that $\omega(a' \wedge y) = 0$ as well; thus (2.3) trivially holds true if $p > g - i$. If $p < g - i$, then $q > h - j$ and the same conclusion applies. Therefore, we can assume in the sequel that $p = g - i$ and $q = h - j$.

To proceed, we consider a presentation $H^M = \langle \gamma_1, \dots, \gamma_{g+r} \mid \rho_1, \dots, \rho_r \rangle$ and a presentation $H^N = \langle \mu_1, \dots, \mu_{h+s} \mid \zeta_1, \dots, \zeta_s \rangle$. By the above-mentioned isomorphism between $H^M \oplus H^N$ and H , we obtain a presentation

$$H = \langle \gamma_1, \dots, \gamma_{g+r}, \mu_1, \dots, \mu_{h+s} \mid \rho_1, \dots, \rho_r, \zeta_1, \dots, \zeta_s \rangle.$$

Note that, with these choices of presentations, the matrix corresponding to H is the direct sum of the matrices corresponding to H^M and H^N . Therefore, we get

$$\begin{aligned}
\omega(a \wedge y) &\stackrel{(2.5)}{=} (-1)^{is+p(s+j)} \mathcal{A}_M^\varphi (\Lambda^i m_-(x^M) \wedge \Lambda^{g-i} m_+(y^M)) \\
&\quad \cdot \mathcal{A}_N^\psi (\Lambda^j n_-(x^N) \wedge \Lambda^{h-j} n_+(y^N)) \\
&= (-1)^{is+p(s+j)} \omega^M (A(M, \varphi)(x^M) \wedge y^M) \cdot \omega^N (A(N, \psi)(x^N) \wedge y^N) \\
&\stackrel{(2.4)}{=} (-1)^{is+p(s+j)} \omega (A(M, \varphi)(x^M) \wedge y^M \wedge A(N, \psi)(x^N) \wedge y^N) \\
&= (-1)^{is+p(s+j)+p(j+\delta h)} \omega (A(M, \varphi)(x^M) \wedge A(N, \psi)(x^N) \wedge y^M \wedge y^N) \\
&= (-1)^{g(s+h)} \omega (a' \wedge y).
\end{aligned}$$

□

Lemma 2.5. *For any morphisms $(M, \varphi) \in \text{Cob}_G((g_-, \varphi_-), (g_+, \varphi_+))$ and $(N, \psi) \in \text{Cob}_G((h_-, \psi_-), (h_+, \psi_+))$ such that $(g_+, \varphi_+) = (h_-, \psi_-)$, we have*

$$A((N, \psi) \circ (M, \varphi)) = A(N, \psi) \circ A(M, \varphi).$$

The next subsection is devoted to the proof of Lemma 2.5.

2.3. Proof of the functoriality of A . We use the notations of Lemma 2.5 and we set

$$\begin{aligned}
g &:= g_- + g_+, & h &:= h_- + h_+, & f &:= g_- + h_+, \\
\delta g &:= g_+ - g_-, & \delta h &:= h_+ - h_-, & \delta f &:= h_+ - g_-, \\
H^M &:= H_1^\varphi(M, I), & H^N &:= H_1^\psi(N, I), & H &:= H_1^{\psi+\varphi}(N \circ M, I).
\end{aligned}$$

Let $v = (v_1, \dots, v_{2g_+})$ be a basis of $H_1^{\varphi+}(F_{g_+}, \star)$: we set $mv_i := m_+(v_i)$ and $nv_i := n_-(v_i)$ for all $i = 1, \dots, 2g_+$. We consider presentations of the following form:

$$\begin{aligned}
H^M &= \langle mv_1, \dots, mv_{2g_+}, u_1, \dots, u_r \mid \zeta_1, \dots, \zeta_{r+\delta g} \rangle, \\
H^N &= \langle nv_1, \dots, nv_{2h_-}, w_1, \dots, w_s \mid \rho_1, \dots, \rho_{s-\delta h} \rangle.
\end{aligned}$$

Applying the Mayer–Vietoris theorem to $N \circ M$, we obtain that the $\mathbb{Z}[G]$ -module H is generated by

$$(2.6) \quad mv_1, \dots, mv_{2g_+}, nv_1, \dots, nv_{2h_-}, u_1, \dots, u_r, w_1, \dots, w_s$$

subject to the relations $\zeta_1, \dots, \zeta_{r+\delta g}, \rho_1, \dots, \rho_{s-\delta h}, mv_1 - nv_1, \dots, mv_{2g_+} - nv_{2g_+}$.

In the sequel, we set $H_- := H_1^{\varphi-}(F_{g_-}, \star)$ and $H_+ := H_1^{\psi+}(F_{h_+}, \star)$. Let $x \in \Lambda^j H_-$ and $y \in \Lambda^{f-j} H_+$: we wish to compute

$$\mathcal{A}_{N \circ M}^{\psi+\varphi} (\Lambda^j m_-(x) \wedge \Lambda^{f-j} n_+(y))$$

using the previous presentation of H . For this, we perform some computations in $\Lambda^k \Gamma$ where $k := 4g_+ + r + s$ and Γ denotes the free $\mathbb{Z}[G]$ -module generated by the k symbols listed at (2.6). Set $\zeta := \zeta_1 \wedge \cdots \wedge \zeta_{r+\delta g}$, $\rho := \rho_1 \wedge \cdots \wedge \rho_{s-\delta h}$. Then, we have

$$\begin{aligned} & \zeta \wedge \rho \wedge (mv_1 - nv_1) \wedge \cdots \wedge (mv_{2g_+} - nv_{2g_+}) \wedge \widetilde{\Lambda^j m_-(x)} \wedge \widetilde{\Lambda^{f-j} n_+(y)} \\ &= \sum_P (-1)^{|P|} \varepsilon_P \cdot \zeta \wedge \rho \wedge mv_P \wedge nv_{\overline{P}} \wedge \widetilde{\Lambda^j m_-(x)} \wedge \widetilde{\Lambda^{f-j} n_+(y)} \\ &= \sum_P (-1)^{|P|(j+1)} \varepsilon_P \cdot \left(\zeta \wedge mv_P \wedge \widetilde{\Lambda^j m_-(x)} \right) \\ & \quad \wedge \left(\rho \wedge nv_{\overline{P}} \wedge \widetilde{\Lambda^{f-j} n_+(y)} \right) \in \Lambda^k \Gamma. \end{aligned}$$

Here the sums are taken over all parts $P \subset \{1, \dots, 2g_+\}$, \overline{P} denotes the complement of P , mv_P is the wedge of the mv_i for $i \in P$, $nv_{\overline{P}}$ is the wedge of the nv_i for $i \in \overline{P}$ and ε_P is the signature of the permutation $P\overline{P}$ (where the elements of P in increasing order are followed by the elements of \overline{P} in increasing order). A sign $(-1)^{(s-\delta h)(j+|P|)}$ is missing in the second sum but, since the presentation of H^N is arbitrary of deficiency h , we can assume that its number of relations $(s - \delta h)$ is even.

In the sequel, we omit the ‘‘tilde’’ notation to distinguish elements of ΛH from their lifts to $\Lambda \Gamma$. Note that, in the above sums, the multivector $\zeta \wedge mv_P \wedge \Lambda^j m_-(x)$ has degree $(r + \delta g) + |P| + j$ which is greater than $2g_+ + r$ as soon as $|P| > g - j$; similarly, the multivector $\rho \wedge nv_{\overline{P}} \wedge \Lambda^{f-j} n_+(y)$ has degree $(s - \delta h) + (2g_+ - |P|) + (f - j)$ which is greater than $2h_- + s$ as soon as $|P| < g - j$; since $2g_+ + r$ and $2h_- + s$ are respectively the numbers of generators of H^M and H^N in the above presentations, the summand corresponding to P vanishes for $|P| > g - j$ and for $|P| < g - j$. Therefore the above sums are actually indexed by the subsets $P \subset \{1, \dots, 2g_+\}$ having cardinality $g - j$, and we get

$$\begin{aligned} & \zeta \wedge \rho \wedge (mv_1 - nv_1) \wedge \cdots \wedge (mv_{2g_+} - nv_{2g_+}) \wedge \Lambda^j m_-(x) \wedge \Lambda^{f-j} n_+(y) \\ &= \sum_{|P|=g-j} \varepsilon'_P \cdot \left(\zeta \wedge mv_P \wedge \Lambda^j m_-(x) \right) \wedge \left(\rho \wedge nv_{\overline{P}} \wedge \Lambda^{f-j} n_+(y) \right) \end{aligned}$$

where we have set $\varepsilon'_P := (-1)^{|P|(j+1)} \varepsilon_P$. The summand is here equal to

$$\begin{aligned} & \varepsilon'_P \cdot \left(\zeta \wedge mv_P \wedge \Lambda^j m_-(x) \right) \wedge \left(\rho \wedge nv_{\overline{P}} \wedge \Lambda^{f-j} n_+(y) \right) \\ &= \varepsilon'_P \cdot \left(\mathcal{A}_M^\varphi(mv_P \wedge \Lambda^j m_-(x)) \cdot (mv \wedge u) \right) \\ & \quad \wedge \left(\mathcal{A}_N^\psi(nv_{\overline{P}} \wedge \Lambda^{f-j} n_+(y)) \cdot (nv \wedge w) \right) \\ &= \varepsilon'_P \cdot \mathcal{A}_M^\varphi(mv_P \wedge \Lambda^j m_-(x)) \mathcal{A}_N^\psi(nv_{\overline{P}} \wedge \Lambda^{f-j} n_+(y)) \cdot (mv \wedge nv \wedge u \wedge w). \end{aligned}$$

We deduce that

$$\begin{aligned}
 & \mathcal{A}_{N \circ M}^{\psi + \varphi}(\Lambda^j m_-(x) \wedge \Lambda^{f-j} n_+(y)) \\
 &= \sum_{|P|=g-j} \varepsilon'_P \cdot \mathcal{A}_M^\varphi(mv_P \wedge \Lambda^j m_-(x)) \cdot \mathcal{A}_N^\psi(nv_{\overline{P}} \wedge \Lambda^{f-j} n_+(y)) \\
 &= \mathcal{A}_N^\psi \left(\sum_{|P|=g-j} \varepsilon'_P \cdot \mathcal{A}_M^\varphi(mv_P \wedge \Lambda^j m_-(x)) \cdot nv_{\overline{P}} \wedge \Lambda^{f-j} n_+(y) \right) \\
 &= \mathcal{A}_N^\psi \left(\sum_{|P|=g-j} (-1)^{|P|} \varepsilon_P \cdot \omega(A(M, \varphi)(x) \wedge v_P) \cdot nv_{\overline{P}} \wedge \Lambda^{f-j} n_+(y) \right).
 \end{aligned}$$

We can assume that the basis v of $H_1^{\varphi+}(F_{g+}, \star)$ is compatible with the chosen volume form ω , in the sense that $\omega(v_1 \wedge \cdots \wedge v_{2g+}) = 1$. Observe that, for all $z \in \Lambda^{j+\delta g} H_1^{\varphi+}(F_{g+}, \star)$, we have the identities

$$z = \sum_{|P|=g-j} \varepsilon_{\overline{P}} \cdot \omega(z \wedge v_P) \cdot v_{\overline{P}} = \sum_{|P|=g-j} (-1)^{|P|} \cdot \varepsilon_P \cdot \omega(z \wedge v_P) \cdot v_{\overline{P}}$$

where the sums range over all subsets $P \subset \{1, \dots, 2g+\}$ of cardinality $g-j$. Hence

$$\begin{aligned}
 \mathcal{A}_{N \circ M}^{\psi + \varphi}(\Lambda^j m_-(x) \wedge \Lambda^{f-j} n_+(y)) &= \mathcal{A}_N^\psi(\Lambda^{j+\delta g} n_{-A(M, \varphi)(x)} \wedge \Lambda^{f-j} n_+(y)) \\
 &= \omega(A(N, \psi)(A(M, \varphi)(x)) \wedge y).
 \end{aligned}$$

It follows that $\omega(A((N, \psi) \circ (M, \varphi))(x) \wedge y) = \omega(A(N, \psi)(A(M, \varphi)(x)) \wedge y)$, which concludes the proof of Lemma 2.5.

3. Alexander functor and knots

In this section, we relate the functor A to the classical Alexander polynomial of knots. We fix a finitely generated free abelian group G ; the extension of a group homomorphism $\varphi : A \rightarrow G$ to a ring homomorphism $\mathbb{Z}[A] \rightarrow \mathbb{Z}[G]$ is still denoted by φ .

3.1. The Alexander polynomial of a topological pair. Given a finitely generated $\mathbb{Z}[G]$ -module N and an integer $i \geq 0$, the i -th Alexander polynomial of N is the greatest common divisor of all minors of order $n-i$ in an $m \times n$ presentation matrix of N . This algebraic invariant is denoted by $\Delta_i N \in \mathbb{Z}[G]/\pm G$.

Let (X, Y) be a pair of topological spaces, and assume that they have the homotopy type of finite CW-complexes. Consider a group homomorphism $\varphi : H_1(X) \rightarrow G$. The Alexander polynomial of (X, Y) with coefficients φ is

$$\Delta^\varphi(X, Y) := \Delta_0 H_1^\varphi(X, Y) \in \mathbb{Z}[G]/\pm G.$$

If Y is empty, we set $\Delta^\varphi(X) := \Delta_0 H_1^\varphi(X)$.

3.2. The Alexander function in genus one. Let M be a compact connected orientable 3-manifold with connected boundary, and fix a base point $\star \in \partial M$. Let also $\varphi : H_1(M) \rightarrow G$ be a group homomorphism. The next lemma generalizes Property 1 of the Alexander function given in [Les].

Lemma 3.1. *Assume that $g(M) = 1$ and that φ is not trivial. Then, for any $h \in H := H_1^\varphi(M, \star)$, we have*

$$\mathcal{A}_M^\varphi(h) = \begin{cases} \Delta^\varphi(M) \cdot \partial_*(h) & \text{if } \text{rank } \varphi(H_1(M)) \geq 2, \\ \Delta^\varphi(M) \cdot \frac{\partial_*(h)}{t-1} & \text{if } \text{rank } \varphi(H_1(M)) = 1 \text{ and } t \text{ is a generator.} \end{cases}$$

Here $\partial_* : H \rightarrow \mathbb{Z}[G]$ is the connecting homomorphism $H_1^\varphi(M, \star) \rightarrow H_0^\varphi(\star)$ in the long exact sequence of the pair (M, \star) , followed by the canonical isomorphism $H_0^\varphi(\star) \simeq \mathbb{Z}[G]$.

We shall deduce Lemma 3.1 from the following.

Lemma 3.2. *If φ is not trivial, then $\Delta^\varphi(M) = \Delta_1 H_1^\varphi(M, \star)$.*

Proof. The long exact sequence in φ -twisted homology for the pair (M, \star) gives

$$0 \longrightarrow H_1^\varphi(M) \longrightarrow H_1^\varphi(M, \star) \longrightarrow H_0^\varphi(\star) \longrightarrow H_0^\varphi(M) \longrightarrow 0.$$

Since the $\mathbb{Z}[G]$ -module $H_0^\varphi(\star) \simeq \mathbb{Z}[G]$ is torsion-free, we deduce that

$$(3.1) \quad \text{Tors } H_1^\varphi(M) \simeq \text{Tors } H_1^\varphi(M, \star).$$

Besides, the above exact sequence implies that

$$\text{rank } H_1^\varphi(M) - \text{rank } H_1^\varphi(M, \star) + 1 - \text{rank } H_0^\varphi(M) = 0.$$

We now show that $\text{rank } H_0^\varphi(M) = 0$. By considering a cell decomposition of M with \star as a single 0-cell and some 1-cells e_1, \dots, e_r , we see that

$$H_0^\varphi(M) \simeq \mathbb{Z}[G] / \langle (g_1 - 1), \dots, (g_r - 1) \rangle_{\text{ideal}}$$

where $g_i := \varphi([e_i]) \in G$. Thus we have the short exact sequence of modules

$$0 \longrightarrow I_\varphi \longrightarrow \mathbb{Z}[G] \longrightarrow H_0^\varphi(M) \longrightarrow 0,$$

where I_φ is the ideal generated by the $\varphi(h) - 1$ for all $h \in H_1(M)$. By tensoring with the field of fractions $Q(G)$, we obtain

$$0 \longrightarrow Q(G) \otimes_{\mathbb{Z}[G]} I_\varphi \longrightarrow Q(G) \longrightarrow Q(G) \otimes_{\mathbb{Z}[G]} H_0^\varphi(M) \longrightarrow 0.$$

Since φ is not trivial, $Q(G) \otimes_{\mathbb{Z}[G]} I_\varphi \neq 0$ so that $Q(G) \otimes_{\mathbb{Z}[G]} H_0^\varphi(M) = 0$. Hence

$$(3.2) \quad \text{rank } H_1^\varphi(M, \star) = \text{rank } H_1^\varphi(M) + 1.$$

We conclude thanks to (3.1) and (3.2) using the following:

Fact. [Bla, Lemma 4.10]. *Let N be a finitely generated $\mathbb{Z}[G]$ -module. Then*

$$\Delta_i(N) = \begin{cases} 0 & \text{if } i < \text{rank}(N), \\ \Delta_{i-\text{rank } N}(\text{Tors } N) & \text{if } i \geq \text{rank}(N). \end{cases} \quad \square$$

Proof of Lemma 3.1. Observe that, for any oriented loop ρ in M based at \star , we have $\partial_*([\widehat{\rho}]) = \varphi([\rho]) - 1$. Thus, the greatest common divisor of $\partial_*(H)$ is

$$\text{gcd } \partial_*(H) = \text{gcd } \{\varphi(x) - 1 \mid x \in H_1(M)\} \in \mathbb{Z}[G]/\pm G.$$

Since φ is assumed to be non-trivial, we deduce that

$$\text{gcd } \partial_*(H) = \begin{cases} 1 & \text{if } \text{rank } \varphi(H_1(M)) \geq 2, \\ t - 1 & \text{if } \text{rank } \varphi(H_1(M)) = 1 \text{ and } t \text{ is a generator.} \end{cases}$$

Therefore, we have to prove that

$$(3.3) \quad \mathcal{A}_M^\varphi(h) = \Delta^\varphi(M) \cdot \frac{\partial_*(h)}{\text{gcd } \partial_*(H)}.$$

For this, we consider a presentation $H = \langle \gamma_1, \dots, \gamma_{r+1} \mid \rho_1, \dots, \rho_r \rangle$ and let A be the associated $r \times (r + 1)$ matrix. We have

$$\forall z_1, \dots, z_{r+1} \in \mathbb{Z}[G], \mathcal{A}_M^\varphi(z_1\gamma_1 + \dots + z_{r+1}\gamma_{r+1}) = \sum_{i=1}^{r+1} (-1)^{i+r+1} \det(A_i) z_i$$

where A_i is the matrix A with the i -th column removed. Then Lemma 3.2 gives

$$(3.4) \quad \Delta^\varphi(M) = \Delta_1 H = \text{gcd } \mathcal{A}_M^\varphi(H).$$

It follows that $\Delta^\varphi(M) = 0$ if and only if $\mathcal{A}_M^\varphi = 0$. In that case (3.3) trivially holds true: thus we assume in the sequel that $\mathcal{A}_M^\varphi \neq 0$. Lemma 2.3 implies that $\text{rank } H = 1$: it follows that any two $Q(G)$ -linear maps $Q(G) \otimes_{\mathbb{Z}[G]} H \rightarrow Q(G)$ are linearly dependent. Since $\mathcal{A}_M^\varphi \neq 0$ and $\partial_* \neq 0$, we deduce that there exist non-zero elements $D, E \in \mathbb{Z}[G]$ such that

$$(3.5) \quad \forall h \in H, \mathcal{A}_M^\varphi(h) = \frac{D}{E} \cdot \partial_*(h)$$

or, equivalently, $D\partial_*(h) = E\mathcal{A}_M^\varphi(h)$ for all $h \in H$. Hence $D \text{gcd } \partial_*(H) = E \text{gcd } \mathcal{A}_M^\varphi(H)$ and we deduce from (3.4) that

$$(3.6) \quad \frac{D}{E} = \frac{\Delta^\varphi(M)}{\text{gcd } \partial_*(H)}.$$

The identity (3.3) is then deduced from (3.5) and (3.6). □

3.3. The functor \mathbf{A} on knot exteriors. Let K be an oriented knot in an oriented homology 3-sphere N . The *Alexander polynomial* of K is classically defined as

$$\Delta(K) := \Delta^{\varphi_K}(M_K) = \Delta_0 H_1^{\varphi_K}(M_K) \in \mathbb{Z}[G]/\pm G$$

where M_K is the complement of an open tubular neighborhood of K in N , G is the infinite cyclic group spanned by t , and $\varphi_K : H_1(M_K) \rightarrow G$ is the isomorphism mapping an oriented meridian $\mu \subset \partial M_K$ of K to t . Note that $\Delta(K)$ is a Laurent polynomial in the variable t , which is defined up to multiplication by a monomial $\pm t^k$ for $k \in \mathbb{Z}$.

We make M_K a morphism $1 \rightarrow 0$ in the category Cob by choosing a boundary-parametrization $m : F(1, 0) \rightarrow \partial M_K$ such that $\mu_- := m^{-1}(\mu)$ is contained in the bottom surface F_1 and goes through the base point \star . Set $H_- := H_1^{\varphi_K m_-}(F_1, \star)$. The following proposition shows that the knot invariants $\Delta(K)$ and $\mathbf{A}(M_K, \varphi_K)$ carry the same topological information. This is deduced from Lemma 3.1 applied to $M := M_K$.

Proposition 3.3. *With the above notation and for any $h \in \Lambda^i H_-$, we have*

$$\mathbf{A}(M_K, \varphi_K)(h) = \begin{cases} \Delta(K) \cdot \partial_*(h)/(t-1) & \text{if } i = 1, \\ 0 & \text{otherwise,} \end{cases}$$

where $\partial_* : H_- \rightarrow \mathbb{Z}[G]$ is the connecting homomorphism for the pair (F_1, \star) . In particular, we have $\Delta(K) = \mathbf{A}(M_K, \varphi_K)([\widehat{\mu}_-])$.

4. The Reidemeister functor \mathbf{R}

In this section, we construct the Reidemeister functor \mathbf{R} . We fix a field \mathbb{F} and a subgroup G of \mathbb{F}^\times . In this section, the extension of a group homomorphism $\varphi : A \rightarrow G$ to a ring homomorphism $\mathbb{Z}[A] \rightarrow \mathbb{F}$ is still denoted by φ .

4.1. The Reidemeister function. We use the elementary theory of abelian Reidemeister torsions to construct an analogue of the Alexander function considered in §2.1. Let M be a compact connected orientable 3-manifold with connected boundary, and let $\varphi : H_1(M) \rightarrow G$ be a group homomorphism. We fix a base point $\star \in \partial M$ and we set $g := g(M) = 1 - \chi(M)$.

Lemma 4.1. *We have $H_i^\varphi(M, \star) = 0$ if $i = 0$ or $i > 2$. Moreover, we have*

$$\dim H_1^\varphi(M, \star) = g + \dim H_2^\varphi(M, \star).$$

Proof. Since ∂M is non-empty, M deformation retracts to a connected 2-dimensional complex whose only 0-cell is \star : the first assertion follows. Moreover, we have

$$-g = \chi(M) - 1 = \chi(M, \star) = -\dim H_1^\varphi(M, \star) + \dim H_2^\varphi(M, \star). \quad \square$$

Denote $H := H_1^\varphi(M, \star)$ and assume in this paragraph that $\dim H = g$. We choose a cell decomposition of M where \star is a 0-cell: by Lemma 4.1, the homology of the φ -twisted cell chain complex $C^\varphi(M, \star)$ is concentrated in degree 1. For every dimension $i \in \{0, \dots, 3\}$, let $n_i \geq 0$ be the number of relative i -cells of (M, \star) and order them $\sigma_1^{(i)}, \dots, \sigma_{n_i}^{(i)}$ in an arbitrary way. For every cell σ of (M, \star) , we also choose an orientation of σ and a lift $\hat{\sigma}$ of σ to the maximal abelian cover \widehat{M} of M . Thus, we get a basis $c := (c_3, c_2, c_1, c_0)$ of the \mathbb{F} -chain complex $C^\varphi(M, \star)$ where, for every $i \in \{0, \dots, 3\}$, the basis of the \mathbb{F} -vector space $C_i^\varphi(M, \star)$ is given by $c_i := (1 \otimes \hat{\sigma}_1^{(i)}, \dots, 1 \otimes \hat{\sigma}_{n_i}^{(i)})$. Then we consider the function $H^g \rightarrow \mathbb{F}$ defined by

$$(4.1) \quad (h_1, \dots, h_g) \mapsto \begin{cases} \tau(C^\varphi(M, \star); c, (h_1, \dots, h_g)) & \text{if } h_1 \wedge \dots \wedge h_g \neq 0, \\ 0 & \text{otherwise.} \end{cases}$$

Here $\tau(C; c, h)$ denotes the torsion of the finite \mathbb{F} -chain complex C with basis c and homological basis h : see §A.1. It follows from the definition of the torsion that the map (4.1) is multilinear and alternate: see Lemma A.2.

Definition 4.2. The *Reidemeister function* of M with coefficients φ is the \mathbb{F} -linear map $\mathcal{R}_M^\varphi : \Lambda^g H \rightarrow \mathbb{F}$ defined by (4.1) if $\dim H = g$ and by $\mathcal{R}_M^\varphi := 0$ if $\dim H \neq g$.

Because of the choice of the orders, orientations, and lifts of the cells of (M, \star) , the map \mathcal{R}_M^φ is only defined up to multiplication by an element of $\pm G \subset \mathbb{F}$. It remains to justify that $\mathcal{R}_M^\varphi \in \text{Hom}(\Lambda^g H, \mathbb{F})/\pm G$ defines a topological invariant of M (i.e., it does not depend on the choice of the cell decomposition). Note that we do not need Chapman's result on the topological invariance of the torsion of CW-complexes [Cha, Coh] since we are considering here manifolds of dimension 3. Specifically, using Whitehead's theory of smooth triangulations and the fact that the Reidemeister torsion of CW-complexes is invariant under cellular subdivisions, we obtain that the above definition of \mathcal{R}_M^φ applied to a smooth triangulation of (M, \star) produces an invariant of smooth 3-manifolds. (See [Mil2, §9] or [Tur3, §3] for similar arguments which are valid in any dimension.) Next, we appeal to the 3-dimensional Hauptvermutung to conclude that \mathcal{R}_M^φ is an invariant of topological 3-manifolds. Thus, we can consider in Definition 4.2 an arbitrary cell decomposition of (M, \star) provided it can be subdivided to a smooth triangulation of M .

4.2. Definition of \mathbf{R} . The definition of the functor \mathbf{R} from the Reidemeister function \mathcal{R} goes parallel to the definition of \mathbf{A} from \mathcal{A} (see §2.2). Thus we associate to any object (g, φ) of Cob_G the exterior algebra

$$\mathbf{R}(g, \varphi) := \Lambda H_1^\varphi(F_g, \star)$$

of the \mathbb{F} -vector space $H^\varphi(F_g, \star) = H_1^\varphi(F_g, \star)$, which has dimension $2g$. Next, we associate to any morphism (M, φ) from (g_-, φ_-) to (g_+, φ_+) an \mathbb{F} -linear map

$$\mathbf{R}(M, \varphi) : \Lambda H_1^{\varphi_-}(F_{g_-}, \star) \longrightarrow \Lambda H_1^{\varphi_+}(F_{g_+}, \star)$$

of degree $\delta g := g_+ - g_-$ in the following way. We set $H := H_1^\varphi(M, I)$ where $I := m(\star \times [-1, 1])$, $H_\pm := H_1^{\varphi_\pm}(F_{g_\pm}, \star)$ and $g := g_+ + g_-$. Then, for any integer $j \geq 0$, the image $\mathbf{R}(M, \varphi)(x) \in \Lambda^{j+\delta g} H_+$ of any $x \in \Lambda^j H_-$ is defined by the following property:

$$\forall y \in \Lambda^{g-j} H_+, \quad \mathcal{R}_M^\varphi(\Lambda^j m_-(x) \wedge \Lambda^{g-j} m_+(y)) = \omega(\mathbf{R}(M, \varphi)(x) \wedge y).$$

Here $\omega : \Lambda^{2g+} H_+ \rightarrow \mathbb{F}$ is an arbitrary volume form which is *integral* in the following sense: regarding H_+ as $\mathbb{F} \otimes_{\mathbb{Z}[H_1(F_{g_+})]} H_1(F_{g_+}, \star; \mathbb{Z}[H_1(F_{g_+})])$, we assume that ω arises from an arbitrary volume form on the free $\mathbb{Z}[H_1(F_{g_+})]$ -module $H_1(F_{g_+}, \star; \mathbb{Z}[H_1(F_{g_+})])$. Due to the choices of this volume form and of the ordered/oriented lifts of the cells to \widehat{M} , the map $\mathbf{R}(M, \varphi)$ is only defined up to multiplication by an element of $\pm G \subset \mathbb{F}$. Besides, $\mathbf{R}(M, \varphi)$ is trivial on $\Lambda^j H_-$ for any $j < \max(0, -\delta g)$ and any $j > \min(g, 2g_-)$.

The next two lemmas show that the above paragraph defines a monoidal functor $\mathbf{R} : \text{Cob}_G \rightarrow \text{grVect}_{\mathbb{F}, \pm G}$, which proves Theorem II of the Introduction.

Lemma 4.3. *For any morphisms $(M, \varphi) \in \text{Cob}_G((g_-, \varphi_-), (g_+, \varphi_+))$ and $(N, \psi) \in \text{Cob}_G((h_-, \psi_-), (h_+, \psi_+))$, we have*

$$(4.2) \quad \mathbf{R}((M, \varphi) \otimes (N, \psi)) = \mathbf{R}(M, \varphi) \otimes \mathbf{R}(N, \psi).$$

Proof. We set $g := g_+ + g_-$, $h := h_+ + h_-$, $\delta g := g_+ - g_-$, $\delta h := h_+ - h_-$ and

$$H_\pm^M := H_1^{\varphi_\pm}(F_{g_\pm}, \star), \quad H_\pm^N := H_1^{\psi_\pm}(F_{h_\pm}, \star), \quad H_\pm := H_1^{\varphi_\pm \oplus \psi_\pm}(F_{g_\pm+h_\pm}, \star),$$

$$H^M := H_1^\varphi(M, I), \quad H^N := H_1^\psi(N, I), \quad H := H_1^{\varphi \oplus \psi}(M \sharp_\partial N, I).$$

Since M and N intersect in $M \sharp_\partial N$ along a 2-disk which retracts onto I , the Mayer–Vietoris theorem gives an isomorphism $H^M \oplus H^N \xrightarrow{\simeq} H$. If $\dim(H^M) > g$, then $\mathcal{R}_M^\varphi = 0$ by definition, so that $\mathbf{R}(M, \varphi) = 0$; moreover,

$$\dim(H) = \dim(H^M) + \dim(H^N) > g + h$$

so that $R((M, \varphi) \otimes (N, \psi)) = 0$ as well, and (4.2) trivially holds true in that case. Therefore, we can assume that $\dim(H^M) = g$ and $\dim(H^N) = h$.

Let $x^M = (x_1^M, \dots, x_i^M)$ be a family of vectors in H_-^M and let $x^N = (x_1^N, \dots, x_j^N)$ be a family of vectors in H_-^N . We consider the element

$$x := x^M \otimes x^N \in \Lambda^i H_-^M \otimes \Lambda^j H_-^N \subset \Lambda^{i+j}(H_-^M \oplus H_-^N) = \Lambda^{i+j} H_-.$$

We aim at showing that $r := R((M, \varphi) \otimes (N, \psi))(x)$ is equal to

$$r' := (R(M, \varphi) \otimes R(N, \psi))(x) = (-1)^{i\delta h} \cdot R(M, \varphi)(x^M) \otimes R(N, \psi)(x^N).$$

It is enough to prove that, for any integers $p, q \geq 0$ such that $p + q = (g + h) - (i + j)$ and for any families $y^M = (y_1^M, \dots, y_p^M) \subset H_+^M$ and $y^N = (y_1^N, \dots, y_q^N) \subset H_+^N$, we have

$$(4.3) \quad \omega(r \wedge y) = \omega(r' \wedge y)$$

where $y := y^M \otimes y^N \in \Lambda^p H_+^M \otimes \Lambda^q H_+^N \subset \Lambda^{p+q} H_+$. In fact, we only need to prove (4.3) up to multiplication by an element of $\pm G$, provided this factor is independent of i, j, p, q, x and y .

In the sequel, we fix integral volume forms ω^M and ω^N on H_+^M and H_+^N respectively, and we assume that the volume form ω on $H_+ = H_+^M \oplus H_+^N$ is defined by

$$(4.4) \quad \omega(u \wedge v) = \omega^M(u) \cdot \omega^N(v)$$

for any $u \in \Lambda^{2g+} H_+^M, v \in \Lambda^{2h+} H_+^N$. (So ω is integral too.) By definition of R , we have

$$(4.5) \quad \omega(r \wedge y) = \mathcal{R}_{M\#_0 N}^{\varphi \oplus \psi} (\Lambda^i m_-(x^M) \wedge \Lambda^j n_-(x^N) \wedge \Lambda^p m_+(y^M) \wedge \Lambda^q n_+(y^N)).$$

If $p > g - i$, then we have $i + p > \dim(H^M)$ by our assumptions and we obtain $\Lambda^i m_-(x^M) \wedge \Lambda^p m_+(y^M) = 0 \in \Lambda^{i+p} H^M$; we deduce that $\omega(r \wedge y) = 0$; on the other hand, the degree of the multivector $R(M, \varphi)(x^M) \wedge y^M \in \Lambda H_+^M$ is $i + \delta g + p > 2g_+$ so that $\omega(r' \wedge y) = 0$ as well; thus (4.3) trivially holds true if $p > g - i$. If $p < g - i$, then $q > h - j$ and the same conclusion applies. Therefore, we can assume that $p = g - i$ and $q = h - j$ in the sequel.

Since $H^M \oplus H^N \simeq H$, $k := (m_-(x^M), m_+(y^M), n_-(x^N), n_+(y^N))$ is a basis of H if, and only if, the families $k^M := (m_-(x^M), m_+(y^M))$ and $k^N := (n_-(x^N), n_+(y^N))$ are bases of H^M and H^N respectively. If the former condition is not satisfied, then $\omega(r \wedge y)$ is zero by (4.5) and, if the latter condition is not satisfied, then $\omega(r' \wedge y)$ is trivial as well since we have

$$\begin{aligned}
\omega(r' \wedge y) &= (-1)^{i\delta h} \omega \left(\mathbb{R}(M, \varphi)(x^M) \wedge \mathbb{R}(N, \psi)(x^N) \wedge y^M \wedge y^N \right) \\
&= (-1)^{i\delta h + p(j + \delta h)} \omega \left(\mathbb{R}(M, \varphi)(x^M) \wedge y^M \wedge \mathbb{R}(N, \psi)(x^N) \wedge y^N \right) \\
&\stackrel{(4.4)}{=} (-1)^{gh + pj} \omega^M \left(\mathbb{R}(M, \varphi)(x^M) \wedge y^M \right) \cdot \omega^N \left(\mathbb{R}(N, \psi)(x^N) \wedge y^N \right)
\end{aligned}$$

or, equivalently,

$$\begin{aligned}
(4.6) \quad \omega(r' \wedge y) &= (-1)^{gh + pj} \mathcal{R}_M^\varphi \left(\Lambda^i m_-(x^M) \wedge \Lambda^{g-i} m_+(y^M) \right) \\
&\quad \cdot \mathcal{R}_N^\psi \left(\Lambda^j n_-(x^N) \wedge \Lambda^{h-j} n_+(y^N) \right).
\end{aligned}$$

Therefore, we can assume in the sequel that k is a basis of H .

Consider next the twisted cell chain complexes $C := C^{\varphi \oplus \psi}(M \#_{\partial} N, I)$, $C^M := C^\varphi(M, I)$ and $C^N := C^\psi(N, I)$. There is a short exact sequence of \mathbb{F} -chain complexes

$$(4.7) \quad 0 \longrightarrow D \longrightarrow C^M \oplus C^N \longrightarrow C \longrightarrow 0$$

where D is the (un-)twisted cell chain complex of the disk $M \cap N \subset M \#_{\partial} N$ relatively to I . Clearly, D is acyclic. By the multiplicativity property of torsions (see Theorem A.3 and Example A.4), we obtain

$$\varepsilon \cdot \tau(C; c, k) \cdot \tau(D; d) \cdot \tau(\mathcal{H}; ((k^M, k^N), k)) = \tau(C^M; c^M, k^M) \cdot \tau(C^N; c^N, k^N)$$

for some appropriate choices of ordered/oriented lifts of the relative cells, which result in bases c, d, c^M, c^N of the chain complexes. Here ε is a sign not depending on i, j, p, q, x, y , and \mathcal{H} is the long exact sequence in homology

$$0 \longrightarrow \dots \longrightarrow 0 \longrightarrow H^M \oplus H^N \longrightarrow H \longrightarrow 0 \longrightarrow 0 \longrightarrow 0$$

induced by (4.7), which we view as a finite acyclic \mathbb{F} -chain complex concentrated in degrees 3, 4 and with basis $((k^M, k^N), k)$. By definition of k, k^M and k^N , we have $\tau(\mathcal{H}; ((k^M, k^N), k)) = 1$ and, since the intersection disk $M \cap N$ can be reduced to I by elementary collapses, the scalar $T := \tau(D; d)$ belongs to $\pm G$. We conclude that

$$\begin{aligned}
\omega(r \wedge y) &\stackrel{(4.5)}{=} (-1)^{pj} \cdot \tau(C; c, k) \\
&= (-1)^{pj} \varepsilon T^{-1} \cdot \tau(C^M; c^M, k^M) \cdot \tau(C^N; c^N, k^N) \\
&\stackrel{(4.6)}{=} (-1)^{gh} \varepsilon T^{-1} \cdot \omega(r' \wedge y). \quad \square
\end{aligned}$$

Lemma 4.4. *For any morphisms $(M, \varphi) \in \text{Cob}_G((g_-, \varphi_-), (g_+, \varphi_+))$ and $(N, \psi) \in \text{Cob}_G((h_-, \psi_-), (h_+, \psi_+))$ such that $(g_+, \varphi_+) = (h_-, \psi_-)$, we have*

$$(4.8) \quad \mathbb{R}((N, \psi) \circ (M, \varphi)) = \mathbb{R}(N, \psi) \circ \mathbb{R}(M, \varphi).$$

The next subsection is devoted to the proof of Lemma 4.4.

4.3. Proof of the functoriality of \mathbf{R} . We use the notations of Lemma 4.4 and we set

$$\begin{aligned} g &:= g_- + g_+, & h &:= h_- + h_+, & f &:= g_- + h_+, \\ \delta g &:= g_+ - g_-, & \delta h &:= h_+ - h_-, & \delta f &:= h_+ - g_-, \\ H^M &:= H_1^\varphi(M, I), & H^N &:= H_1^\psi(N, I), & H &:= H_1^{\psi+\varphi}(N \circ M, I), \\ K^M &:= H_2^\varphi(M, I), & K^N &:= H_2^\psi(N, I), & K &:= H_2^{\psi+\varphi}(N \circ M, I), \\ H_- &:= H_1^{\varphi-}(F_{g_-}, \star), & V &:= H_1^{\varphi+}(F_{g_+}, \star), & H_+ &:= H_1^{\psi+}(F_{h_+}, \star). \end{aligned}$$

Since $N \circ M$ is obtained from M and N by identifying $\partial_+ M$ to $\partial_- N$, there is a short exact sequence of chain complexes

$$(4.9) \quad 0 \longrightarrow \underbrace{C^{\varphi+}(F_{g_+}, \star)}_{D:=} \longrightarrow \underbrace{C^\psi(N, I)}_{C^N:=} \oplus \underbrace{C^\varphi(M, I)}_{C^M:=} \longrightarrow \underbrace{C^{\psi+\varphi}(N \circ M, I)}_{C:=} \longrightarrow 0.$$

Let \mathcal{H} be the corresponding long exact sequence in homology:

$$0 \rightarrow \dots \rightarrow 0 \rightarrow K^N \oplus K^M \rightarrow K \rightarrow V \xrightarrow{(-n_-, m_+)} H^N \oplus H^M \rightarrow H \rightarrow 0 \rightarrow 0 \rightarrow 0.$$

If $K^M \neq 0$, then $\dim(H^M) > g$ by Lemma 4.1 so that $\mathcal{R}_M^\varphi = 0$ and $\mathbf{R}(M, \varphi) = 0$; besides, the long exact sequence \mathcal{H} implies that $K \neq 0$ so that $\mathbf{R}((N, \psi) \circ (M, \varphi)) = 0$; therefore, (4.8) trivially holds true in that case. If $K^N \neq 0$, the same conclusion applies. So, we can assume that $K^M = 0$ and $K^N = 0$ or, equivalently, $\dim H^M = g$ and $\dim H^N = h$.

Let $j \in \{0, \dots, f\}$, and let $x = (x_1, \dots, x_j)$ and $y = (y_1, \dots, y_{f-j})$ be families of vectors in H_- and H_+ respectively. Let $v = (v_1, \dots, v_{2g_+})$ be an arbitrary basis of V and let $\omega^v : \Lambda^{2g_+} V \rightarrow \mathbb{F}$ be the volume form such that $\omega^v(v_1 \wedge \dots \wedge v_{2g_+}) = 1$; there exists an $\alpha_v \in \mathbb{F} \setminus \{0\}$ such that $\omega = \alpha_v \cdot \omega^v$ is the integral volume form chosen in the definition of the functor \mathbf{R} . We have $\mathbf{R}(M, \varphi)(x) \in \Lambda^{j+\delta g} V$, hence

$$\mathbf{R}(M, \varphi)(x) = \sum_{|P|=g-j} \varepsilon_{\overline{P}} \cdot \omega^v(\mathbf{R}(M, \varphi)(x) \wedge v_P) \cdot v_{\overline{P}}$$

where the sum is taken over all subsets $P \subset \{1, \dots, 2g_+\}$ of cardinality $g-j$, \overline{P} denotes the complement of P , v_P (respectively $v_{\overline{P}}$) is the wedge of the v_i 's for $i \in P$ (respectively $i \in \overline{P}$), and $\varepsilon_{\overline{P}}$ is the signature of the permutation $\overline{P}P$ (where the elements of \overline{P} in increasing order are followed by the elements of P in increasing order). We deduce that

(4.10)

$$\begin{aligned}
& \omega \left(\mathbf{R}(N, \psi) \left(\mathbf{R}(M, \varphi)(x) \right) \wedge y \right) \\
&= \mathcal{R}_N^\psi \left(\Lambda^{j+\delta g} n_- \mathbf{R}(M, \varphi)(x) \wedge \Lambda^{f-j} n_+(y) \right) \\
&= \mathcal{R}_N^\psi \left(\sum_{|P|=g-j} \varepsilon_{\overline{P}} \cdot \omega^v \left(\mathbf{R}(M, \varphi)(x) \wedge v_P \right) \cdot \Lambda^{j+\delta g} n_-(v_{\overline{P}}) \wedge \Lambda^{f-j} n_+(y) \right) \\
&= \alpha_v^{-1} \mathcal{R}_N^\psi \left(\sum_{|P|=g-j} \varepsilon'_P \cdot \mathcal{R}_M^\varphi \left(\Lambda^{g-j} m_+(v_P) \wedge \Lambda^j m_-(x) \right) \right. \\
&\quad \left. \cdot \Lambda^{j+\delta g} n_-(v_{\overline{P}}) \wedge \Lambda^{f-j} n_+(y) \right) \\
&= \alpha_v^{-1} \sum_{|P|=g-j} \varepsilon'_P \cdot \mathcal{R}_M^\varphi \left(\Lambda^{g-j} m_+(v_P) \wedge \Lambda^j m_-(x) \right) \\
&\quad \cdot \mathcal{R}_N^\psi \left(\Lambda^{j+\delta g} n_-(v_{\overline{P}}) \wedge \Lambda^{f-j} n_+(y) \right)
\end{aligned}$$

where $\varepsilon'_P := \varepsilon_{\overline{P}} \cdot (-1)^{j(g-j)}$. If $K \neq 0$, then $\mathbf{R}((N, \psi) \circ (M, \varphi)) = 0$; besides, the long exact sequence in homology \mathcal{H} shows that there exists a $w \in V \setminus \{0\}$ such that $n_-(w) = 0 \in H^N$ and $m_+(w) = 0 \in H^M$; since the basis v of V is arbitrary in (4.10), we can assume that $v_1 = w$. In the last sum indexed by P , the vector w appears either in v_P or in $v_{\overline{P}}$, so that the corresponding summand is always zero; it follows that $\mathbf{R}(N, \psi) \left(\mathbf{R}(M, \varphi)(x) \right) \wedge y = 0$ for any $x \in \Lambda^j H_-$ and $y \in \Lambda^{f-j} H_+$; therefore, (4.8) trivially holds true in that case. Thus, we can assume in the sequel that $K = 0$ or, equivalently, $\dim H = f$.

It now remains to prove using the above assumptions that, for any families of vectors $x = (x_1, \dots, x_j)$ in H_- and $y = (y_1, \dots, y_{f-j})$ in H_+ ,

$$\begin{aligned}
(4.11) \quad & \omega \left(\mathbf{R}((N, \psi) \circ (M, \varphi))(x) \wedge y \right) \\
&= \alpha_v^{-1} \sum_{|P|=g-j} \varepsilon'_P \cdot \mathcal{R}_M^\varphi \left(\Lambda^{g-j} m_+(v_P) \wedge \Lambda^j m_-(x) \right) \\
&\quad \cdot \mathcal{R}_N^\psi \left(\Lambda^{j+\delta g} n_-(v_{\overline{P}}) \wedge \Lambda^{f-j} n_+(y) \right)
\end{aligned}$$

where, as in the previous paragraph, v is an arbitrary basis of V . Assume firstly that $k := (m_-(x), n_+(y))$ is not a basis of H . Then

$$\mathbf{R}((N, \psi) \circ (M, \varphi))(x) \wedge y = \mathcal{R}_{N \circ M}^{\psi+\varphi} \left(\Lambda^j m_-(x) \wedge \Lambda^{f-j} n_+(y) \right)$$

is zero. Besides, the long exact sequence \mathcal{H} implies that there exists $w \in V$ such that

$$\begin{aligned}
m_+(w) &= a_1 m_-(x_1) + \dots + a_j m_-(x_j) \in H^M, \\
-n_-(w) &= b_1 n_+(y_1) + \dots + b_{f-j} n_+(y_{f-j}) \in H^N
\end{aligned}$$

where $a_1, \dots, a_j, b_1, \dots, b_{f-j} \in \mathbb{F}$ are not all zeros. If $w = 0$, then we have $\Lambda^j m_-(x) = 0 \in \Lambda^j H^M$ or $\Lambda^{f-j} n_+(y) = 0 \in \Lambda^{f-j} H^N$ (depending on whether

we can find a non-zero scalar among the a_i 's or among the b_i 's); in both cases, the second term of (4.11) is trivial. If $w \neq 0$, then we take a basis v of V such that $v_1 = w$ and we easily see that the second term of (4.11) is trivial in that case too. Therefore, we can assume in the sequel that $k = (m_-(x), n_+(y))$ is a basis of H .

We now fix a basis $v = (v_1, \dots, v_{2g_+})$ of V such that $\omega(v) = 1$ and we prove (4.11) with $\alpha_v = 1$. Let also k^M and k^N be arbitrary bases of H^M and H^N , respectively. By the multiplicativity property of torsions (see Theorem A.3 and Example A.4), we deduce from (4.9) that

$$(4.12) \quad \begin{aligned} \tau(D; d, v) \cdot \tau(C; c, k) \cdot \tau(\mathcal{H}; (v, (k^N, k^M), k)) \\ = \pm \tau(C^N; c^N, k^N) \cdot \tau(C^M; c^M, k^M) \in \mathbb{F} \end{aligned}$$

for some appropriate choices of ordered/oriented lifts of the relative cells, which result in bases d, c, c^M, c^N of the chain complexes. The sign appearing in (4.12) depends only on the dimensions of the complexes C, D, C^M, C^N and the dimensions of their homology groups. The sequence \mathcal{H} is viewed here as a finite acyclic \mathbb{F} -chain complex concentrated in degrees 3, 4, 5; its torsion is

$$\begin{aligned} \tau(\mathcal{H}; (v, (k^N, k^M), k)) &= \left[\frac{((-n_-, m_+)(v), \text{lift of } k \text{ to } H^N \oplus H^M)}{(k^N, k^M)} \right]^{-1} \\ &= \left[\frac{(k^N, k^M)}{((-n_-, m_+)(v), \text{lift of } k \text{ to } H^N \oplus H^M)} \right], \end{aligned}$$

where the symbol $\left[\frac{a}{b} \right]$ stands for the determinant of the square matrix expressing a family of vectors a in the basis b of $H^N \oplus H^M$. We have $\tau(D; d, v) \in \pm G$ since (F_{g_+}, \star) has the simple homotopy type of a wedge of circles relative to its vertex. We deduce from (4.12) that

$$\begin{aligned} \mathcal{R}_{N \circ M}^{\psi+\varphi}(\Lambda^j m_-(x) \wedge \Lambda^{f-j} n_+(y)) \cdot \left[\frac{(k^N, k^M)}{((-n_-, m_+)(v), \text{lift of } k \text{ to } H^N \oplus H^M)} \right] \\ = \beta_v \cdot \mathcal{R}_M^\varphi(k^M) \cdot \mathcal{R}_N^\psi(k^N) \end{aligned}$$

where $\beta_v \in \pm G$ does not depend on j, x, y, k^M, k^N (but depends on v). The previous identity makes sense, and holds true, when k^M is an arbitrary family of g vectors in H^M and k^N is an arbitrary family of h vectors in H^N . (Indeed, if k^M is not a basis of H^M or k^N is not a basis of H^N , then both sides of this identity are zero.) In particular, we obtain for any subset $P \subset \{1, \dots, 2g_+\}$ of cardinality $g - j$

$$\begin{aligned} & \mathcal{R}_{N \circ M}^{\psi+\varphi}(\Lambda^j m_-(x) \wedge \Lambda^{f-j} n_+(y)) \cdot \left[\frac{(n_-(v_{\bar{P}}), n_+(y), m_+(v_P), m_-(x))}{((-n_-, m_+)(v), \text{lift of } k \text{ to } H^N \oplus H^M)} \right] \\ &= \beta_v \cdot \mathcal{R}_M^\varphi(\Lambda^{g-j} m_+(v_P) \wedge \Lambda^j m_-(x)) \cdot \mathcal{R}_N^\psi(\Lambda^{\delta g+j} n_-(v_{\bar{P}}) \wedge \Lambda^{f-j} n_+(y)). \end{aligned}$$

By multilinearity of the determinant and using the facts that $\dim H^M = g$ and $\dim H^N = h$, we have

$$\begin{aligned} 1 &= \left[\frac{(-n_-(v_1) + m_+(v_1), \dots, -n_-(v_{2g_+}) + m_+(v_{2g_+}), m_-(x), n_+(y))}{((-n_-, m_+)(v), \text{lift of } k \text{ to } H^N \oplus H^M)} \right] \\ &= \sum_{|P|=g-j} \varepsilon_P (-1)^{|\bar{P}|} \left[\frac{(m_+(v_P), n_-(v_{\bar{P}}), m_-(x), n_+(y))}{((-n_-, m_+)(v), \text{lift of } k \text{ to } H^N \oplus H^M)} \right] \\ &= (-1)^{g(f+1)} \sum_{|P|=g-j} \varepsilon'_P \left[\frac{(n_-(v_{\bar{P}}), n_+(y), m_+(v_P), m_-(x))}{((-n_-, m_+)(v), \text{lift of } k \text{ to } H^N \oplus H^M)} \right]. \end{aligned}$$

Thus we obtain identity (4.11), up to multiplication by an element of $\pm G$ not depending on j, x, y . This concludes the proof of Lemma 4.4.

5. Back to the Alexander functor

We show in this section that the functor A is an instance of the functor R .

5.1. A formula for the Reidemeister function. Let M be a compact connected orientable 3-manifold with connected boundary, and fix a base point $\star \in \partial M$. Let also $\varphi : H_1(M) \rightarrow G$ be a group homomorphism with values in a multiplicative subgroup G of a field \mathbb{F} . We use the same notation as in §4.1, where we have introduced \mathcal{R}_M^φ .

When it does not vanish, the Reidemeister function \mathcal{R}_M^φ is defined as an alternated product of 4 determinants since the \mathbb{F} -chain complex $C^\varphi(M, \star)$ has length 3. We now give a recipe to compute it by means of a single determinant using Fox's free derivatives. We consider for this purpose a *spine* X^+ of M , i.e. a 2-dimensional subcomplex X^+ of a smooth triangulation of M such that M retracts to X^+ by elementary collapses; we also assume that \star is a vertex of X^+ . (It is well known that any 3-manifold with boundary has a spine: see for instance [Mat, Remark 1.1.5].) Next, we choose a maximal tree in the 1-skeleton of X^+ which contains \star , and let X be the 2-dimensional CW-complex obtained from X^+ by collapsing that tree to the vertex \star . Hence X has a single 0-cell \star . We denote by $\gamma_1, \dots, \gamma_{g+r}$ the 1-cells of X and we denote by R_1, \dots, R_r the 2-cells of X ; besides, each of these cells is given an arbitrary orientation. The

fundamental group $\pi_1(\Gamma) = \pi_1(\Gamma, \star)$ of the 1-skeleton $\Gamma := \gamma_1 \cup \dots \cup \gamma_{g+r}$ of X is freely generated by the oriented loops $\gamma_1, \dots, \gamma_{g+r}$, hence the free derivatives $\frac{\partial}{\partial \gamma_1}, \dots, \frac{\partial}{\partial \gamma_{g+r}} : \mathbb{Z}[\pi_1(\Gamma)] \rightarrow \mathbb{Z}[\pi_1(\Gamma)]$ are defined. Note that the attaching maps of the oriented 2-cells R_1, \dots, R_r define some elements $\rho_1, \dots, \rho_r \in \pi_1(\Gamma)$.

Lemma 5.1. *Let $\kappa_1, \dots, \kappa_g$ be oriented loops in Γ based at \star and, for all $i \in \{1, \dots, g\}$, let $k_i \in H \simeq H_1^\varphi(X, \star)$ be the homology class of $1 \otimes \widehat{\kappa}_i \in C_1^\varphi(X, \star)$. Then*

$$(5.1) \quad \mathcal{R}_M^\varphi(k_1 \wedge \dots \wedge k_g) = \det \varphi i_* \begin{pmatrix} \frac{\partial \rho_1}{\partial \gamma_1} & \dots & \dots & \dots & \frac{\partial \rho_1}{\partial \gamma_{g+r}} \\ \vdots & & & & \vdots \\ \frac{\partial \rho_r}{\partial \gamma_1} & \dots & \dots & \dots & \frac{\partial \rho_r}{\partial \gamma_{g+r}} \\ \frac{\partial \kappa_1}{\partial \gamma_1} & \dots & \dots & \dots & \frac{\partial \kappa_1}{\partial \gamma_{g+r}} \\ \vdots & & & & \vdots \\ \frac{\partial \kappa_g}{\partial \gamma_1} & \dots & \dots & \dots & \frac{\partial \kappa_g}{\partial \gamma_{g+r}} \end{pmatrix}.$$

Here the composition of φ with the isomorphism $H_1(M) \simeq H_1(X)$ induced by the homotopy equivalence $M \simeq X$ is still denoted by φ , and the ring homomorphism $i_* : \mathbb{Z}[\pi_1(\Gamma)] \rightarrow \mathbb{Z}[\pi_1(M)]$ is induced by the map $i : \Gamma \rightarrow M$ which is the inclusion $\Gamma \subset X$ composed with the homotopy equivalence $X \simeq M$.

Proof. The lemma is proved in a way similar to Milnor’s result relating the Reidemeister torsion of a knot exterior to the Alexander polynomial of the knot [Mil, Theorem 4]. (See also [Tur5, Theorem II.1.2].) By assumption, the pair (M, \star) has the simple homotopy type of (X^+, \star) and, using the multiplicativity property of torsions (Theorem A.3), it can be checked that the Reidemeister torsions of (X, \star) and (X^+, \star) are equal for any choice of homological bases. Therefore we can safely replace M by X in our computation of \mathcal{R}_M^φ . Thus we now consider the φ -twisted cell chain complex

$$C := C^\varphi(X, \star) = \mathbb{F} \otimes_{\mathbb{Z}[H_1(X)]} C(\widehat{X}, p_X^{-1}(\star)).$$

The lifts $\widehat{\gamma}_1, \dots, \widehat{\gamma}_{g+r}$ of $\gamma_1, \dots, \gamma_{g+r}$ define a basis $c_1 := (1 \otimes \widehat{\gamma}_1, \dots, 1 \otimes \widehat{\gamma}_{g+r})$ of C in degree 1. Similarly, the lifts $\widehat{R}_1, \dots, \widehat{R}_r$ of R_1, \dots, R_r that contain $\widehat{\star}$ define a basis $c_2 := (1 \otimes \widehat{R}_1, \dots, 1 \otimes \widehat{R}_r)$ of C in degree 2.

Let A' be the square matrix with entries in \mathbb{F} defined by the right-hand side of (5.1), and let A be the $r \times (g+r)$ matrix defined by the first r rows of A' . Observe that A is the matrix of $\partial_2 : C_2 \rightarrow C_1$ in the bases c_2 and c_1 . Since (X, \star) has no relative cells in degree 0, $H \simeq H_1(C)$ is the cokernel of the linear map $\mathbb{F}^r \rightarrow \mathbb{F}^{g+r}$ defined by the multiplication $v \mapsto vA$. Assume that $\dim H > g$: then the rank of A is less than r , so that all the minors of A of

order r vanish; by expanding the determinant of A' successively along its last g rows, we obtain that $\det A' = 0$ and the lemma trivially holds true in that case. Therefore we can assume that $\dim H = g$.

Observe, next, that the last g rows of A' give the vectors $k_1, \dots, k_g \in H \simeq H_1^\varphi(X, \star)$ as linear combinations of the generators $[1 \otimes \widehat{\gamma}_1], \dots, [1 \otimes \widehat{\gamma}_{g+r}]$ of $H_1^\varphi(X, \star) \simeq H$. If $k := (k_1, \dots, k_g)$ is not a basis of H , then k_1, \dots, k_g are linearly dependent: since the first r rows of A' give a system of relations for the previous set of generators, we deduce that $\det A' = 0$ and the lemma is trivially true in that case too. Thus we can assume that k is a basis of H . Let c be the basis of C given by c_1 in degree 1 and c_2 in degree 2. By Lemma 4.1, the homology of C is concentrated in degree 1 and, for all $i \in \{1, \dots, g\}$, $1 \otimes \widehat{\kappa}_i$ is a 1-cycle of C representing $k_i \in H \simeq H_1(C)$. So, by definition of the function \mathcal{R}_M^φ , we get

$$(5.2) \quad \begin{aligned} \mathcal{R}_M^\varphi(k_1 \wedge \dots \wedge k_g) &= \tau(C; c, k) \\ &= \det(\text{matrix of } (\partial_2(c_2), 1 \otimes \widehat{\kappa}) \text{ in the basis } c_1). \end{aligned}$$

The conclusion follows from the previous two observations. □

Remark 5.2. It follows from Lemma 5.1 that the Reidemeister function has the following *integrality* property: for all $h_1, \dots, h_g \in H_1(M, \star; \mathbb{Z}[H_1(M)])$, we have

$$\mathcal{R}_M^\varphi(\varphi_*(h_1) \wedge \dots \wedge \varphi_*(h_g)) \in \varphi(\mathbb{Z}[H_1(M)])$$

where $\varphi_* : H_1(M, \star; \mathbb{Z}[H_1(M)]) \rightarrow H_1^\varphi(M, \star)$ is the canonical map.

5.2. Specialization of \mathbf{R} to \mathbf{A} . We now assume that G is a finitely generated free abelian group, and we denote by $Q(G)$ the field of fractions of $\mathbb{Z}[G]$. Let M be a compact connected orientable 3-manifold with connected boundary, and fix a base point $\star \in \partial M$. Let $\varphi : H_1(M) \rightarrow G$ be a group homomorphism: we denote by $\varphi_{\mathbb{Z}} : \mathbb{Z}[H_1(M)] \rightarrow \mathbb{Z}[G]$ and by $\varphi : \mathbb{Z}[H_1(M)] \rightarrow Q(G)$ the extensions of φ to ring homomorphisms. Set

$$g := g(M), \quad H_{\mathbb{Z}} := H_1^{\varphi_{\mathbb{Z}}}(M, \star), \quad H := H_1^\varphi(M, \star).$$

Lemma 5.3. *We have the following commutative diagram:*

$$\begin{array}{ccc} \Lambda^g H_{\mathbb{Z}} & \xrightarrow{A_M^\varphi} & \mathbb{Z}[G] \\ \Lambda^g \iota \downarrow & & \downarrow \\ \Lambda^g H & \xrightarrow{\mathcal{R}_M^\varphi} & Q(G), \end{array}$$

where $\iota : H_{\mathbb{Z}} \rightarrow H \simeq Q(G) \otimes_{\mathbb{Z}[G]} H_{\mathbb{Z}}$ denotes the canonical map.

Proof. We proceed as in §5.1: we consider a spine X^+ of M , and we obtain a 2-dimensional CW-complex X with a single vertex \star by collapsing a maximal tree in the 1-skeleton of X^+ . The cells of X are $\gamma_1, \dots, \gamma_{g+r}$ in dimension 1, and R_1, \dots, R_r in dimension 2. Orient $\gamma_1, \dots, \gamma_{g+r}$ and R_1, \dots, R_r arbitrarily, and set

$$C_{\mathbb{Z}} := C^{\varphi_{\mathbb{Z}}}(X, \star), \quad C := C^{\varphi}(X, \star) = Q(G) \otimes_{\mathbb{Z}[G]} C_{\mathbb{Z}}.$$

Since M deformation retracts to X , $H_{\mathbb{Z}}$ is isomorphic to $H_1^{\varphi_{\mathbb{Z}}}(X, \star)$ so that $H_{\mathbb{Z}}$ is the cokernel of $\partial_2 : C_{\mathbb{Z},2} \rightarrow C_{\mathbb{Z},1}$. Let $\widehat{\gamma}_1, \dots, \widehat{\gamma}_{g+r}$ be the preferred lifts of $\gamma_1, \dots, \gamma_{g+r}$ to \widehat{X} , and let $\widehat{R}_1, \dots, \widehat{R}_r$ be the lifts of R_1, \dots, R_r that contain $\widehat{\star}$: we denote by A the matrix of ∂_2 in the bases $(1 \otimes \widehat{R}_1, \dots, 1 \otimes \widehat{R}_r)$ and $(1 \otimes \widehat{\gamma}_1, \dots, 1 \otimes \widehat{\gamma}_{g+r})$. This presentation matrix of the $\mathbb{Z}[G]$ -module $H_{\mathbb{Z}}$ can be used to compute \mathcal{A}_M^{φ} . Specifically, let $k_1, \dots, k_g \in H_{\mathbb{Z}}$ and assume that each k_i has the form $[1 \otimes \widehat{\kappa}_i]$ where κ_i is an oriented loop in the 1-skeleton of X based at \star : then $\mathcal{A}_M^{\varphi}(k_1 \wedge \dots \wedge k_g)$ is the determinant of the matrix obtained from A by adding g rows that express the vectors $1 \otimes \widehat{\kappa}_1, \dots, 1 \otimes \widehat{\kappa}_g$ in the basis $(1 \otimes \widehat{\gamma}_1, \dots, 1 \otimes \widehat{\gamma}_{g+r})$ of $C_{\mathbb{Z},1}$. We deduce from formula (5.2) that $\mathcal{A}_M^{\varphi}(k_1 \wedge \dots \wedge k_g) = \mathcal{R}_M^{\varphi}(\iota(k_1) \wedge \dots \wedge \iota(k_g))$. \square

The next theorem, which compares the Alexander functor to the Reidemeister functor, is a direct application of Lemma 5.3.

Theorem 5.4. *The following diagram is commutative:*

$$\begin{array}{ccc}
 & \text{grMod}_{\mathbb{Z}[G], \pm G} & \\
 \text{Cob}_G & \xrightarrow{A} & \\
 & \text{grVect}_{Q(G), \pm G} & \\
 & \xleftarrow{R} & \\
 & & Q(G) \otimes_{\mathbb{Z}[G]} (-)
 \end{array}$$

6. Reidemeister functor and knots

We now compute the functor R on knot exteriors and we consider, next, the situation of closed 3-manifolds. In this section, we fix a field \mathbb{F} and a multiplicative subgroup G of \mathbb{F} . The extension of a group homomorphism $\varphi : H \rightarrow G$ to a ring homomorphism $\mathbb{Z}[H] \rightarrow \mathbb{F}$ is still denoted by φ .

6.1. The abelian Reidemeister torsion of a CW-pair. Let (X, Y) be a pair of finite CW-complexes, and let $\varphi : \mathbb{Z}[H_1(X)] \rightarrow \mathbb{F}$ be a ring homomorphism. We consider the φ -twisted cell chain complex $C^{\varphi}(X, Y)$ of the pair (X, Y) , which

is a finite \mathbb{F} -chain complex of length $p := \dim X$. For every $i \in \{0, \dots, p\}$, let $n_i \geq 0$ be the number of relative i -cells of (X, Y) and order them $\sigma_1^{(i)}, \dots, \sigma_{n_i}^{(i)}$ in an arbitrary way. For every cell σ of (X, Y) , we also choose an orientation of σ and a lift $\hat{\sigma}$ of σ to the maximal abelian cover \widehat{X} of X . Thus, we obtain a basis $c := (c_p, \dots, c_0)$ of the \mathbb{F} -chain complex $C^\varphi(X, Y)$ where, for every $i \in \{0, \dots, p\}$, the basis of $C_i^\varphi(X, Y)$ is $c_i := (1 \otimes \hat{\sigma}_1^{(i)}, \dots, 1 \otimes \hat{\sigma}_{n_i}^{(i)})$. Recall that the *Reidemeister torsion* of the pair (X, Y) with coefficients φ is the scalar

$$\tau^\varphi(X, Y) := \begin{cases} 0 & \text{if } H^\varphi(X, Y) \neq 0, \\ \tau(C^\varphi(X, Y); c) & \text{if } H^\varphi(X, Y) = 0, \end{cases}$$

where $\tau(C; c)$ denotes the torsion of a finite acyclic \mathbb{F} -chain complex C based by c : see §A.1. The reader is referred to the monograph [Tur4] for an introduction to this combinatorial invariant. Without further structure on the CW-pair (X, Y) , the scalar $\tau^\varphi(X, Y)$ is only defined up to multiplication by an element of $\pm\varphi(H_1(X))$. If $Y = \emptyset$, we denote it by $\tau^\varphi(X)$.

6.2. The Reidemeister function in genus one. We now consider a compact connected orientable 3-manifold M with connected boundary and a group homomorphism $\varphi : H_1(M) \rightarrow G$. Let $\star \in \partial M$ and set $H := H_1^\varphi(M, \star)$. The next lemma relates the Reidemeister function \mathcal{R}_M^φ to the Reidemeister torsion $\tau^\varphi(M)$ in genus one.

Lemma 6.1. *Assume that $g(M) = 1$ and that φ is not trivial. Then, for any $k \in H$,*

$$(6.1) \quad \mathcal{R}_M^\varphi(k) = \tau^\varphi(M) \cdot \partial_*(k).$$

Here $\partial_* : H \rightarrow \mathbb{F}$ is the connecting homomorphism $H_1^\varphi(M, \star) \rightarrow H_0^\varphi(\star)$ in the long exact sequence of the pair (M, \star) , followed by the canonical isomorphism $H_0^\varphi(\star) \simeq \mathbb{F}$.

Proof. Consider a cell decomposition of M where \star is a 0-cell. The short exact sequence of \mathbb{F} -chain complexes

$$(6.2) \quad 0 \longrightarrow \underbrace{C^\varphi(\star)}_{C' :=} \longrightarrow \underbrace{C^\varphi(M)}_{C :=} \longrightarrow \underbrace{C^\varphi(M, \star)}_{C'' :=} \longrightarrow 0$$

induces the following long exact sequence in homology:

$$(6.3) \quad 0 \longrightarrow 0 \longrightarrow 0 \longrightarrow 0 \longrightarrow H_2^\varphi(M) \longrightarrow H_2^\varphi(M, \star) \longrightarrow 0 \longrightarrow H_1^\varphi(M) \longrightarrow H_1^\varphi(M, \star) \xrightarrow{\partial_*} H_0^\varphi(\star) \longrightarrow H_0^\varphi(M) \longrightarrow 0.$$

We regard (6.3) as an acyclic \mathbb{F} -chain complex \mathcal{H} of length 12: let (h', h, h'') be the basis of \mathcal{H} obtained by choosing bases h', h, h'' of $H(C'), H(C), H(C'')$ in each degree. We choose an orientation and a lift to \widehat{M} for every cell of M and, for all $i \in \{0, \dots, 3\}$, we order the i -cells in an arbitrary way. Thus, we obtain bases c', c, c'' of the complexes C', C, C'' , respectively, which are compatible in the sense of §A.1. By the multiplicativity property of torsions (see Theorem A.3), we obtain

$$(6.4) \quad \tau(C; c, h) = \varepsilon \cdot \tau(C'; c', h') \cdot \tau(C''; c'', h'') \cdot \tau(\mathcal{H}; (h', h, h''))$$

where ε is a sign independent of h, h', h'' . If $H_2^\varphi(M) \neq 0$, then $\tau^\varphi(M) = 0$ by definition, but (6.3) gives $H_2^\varphi(M, \star) \neq 0$ and Lemma 4.1 implies that $\dim H_1^\varphi(M, \star) > g(M)$: hence $\mathcal{R}_M^\varphi = 0$ by definition and (6.1) trivially holds true. Therefore we can assume that $H_2^\varphi(M) = 0$.

Besides $H_0^\varphi(M) = 0$ since φ is non-trivial: the fact that $\chi(M) = 1 - g(M)$ is zero implies that $H_1^\varphi(M) = 0$ as well. Thus the chain complex \mathcal{H} defined by (6.3) is concentrated in degrees 2 and 3. Let $k \in H \setminus \{0\}$ which defines a basis h'' of $H(C'')$, and let h' be the basis of $H(C')$ defined by the canonical generator of $H_0^\varphi(\star)$. Then we obtain

$$\tau(\mathcal{H}; (h', h, h'')) = [\partial_*(h''_1)/h'_0]^{(-1)^{2+1}} = \partial_*(k)^{-1}.$$

Besides we have $\tau(C'; c', h') = 1$ by our choices of c' and h' . We conclude using (6.4) that $\tau^\varphi(M) = \varepsilon \cdot \mathcal{R}_M^\varphi(k) \cdot \partial_*(k)^{-1}$. \square

Remark 6.2. If $g(M) = 0$ and φ is not trivial, then $\mathcal{R}_M^\varphi : \mathbb{F} = \Lambda^0 H \rightarrow \mathbb{F}$ is the zero map. Indeed, pick an oriented loop α in M based at \star such that $\varphi([\alpha]) \neq 1$; then $\partial_* : H \rightarrow \mathbb{F}$ does not vanish on $[\hat{\alpha}]$ and it follows that $\dim H > g(M)$.

6.3. The functor \mathbf{R} on knot exteriors. Let K be an oriented knot in a closed connected oriented 3-manifold N , and denote by M_K the complement of an open tubular neighborhood of K in N . We assume given a group homomorphism $\varphi_K : M_K \rightarrow G$ and an oriented closed curve $\lambda \subset \partial M_K$ such that $\varphi_K([\lambda]) \neq 1$. Thus the Reidemeister torsion $\tau^{\varphi_K}(M_K) \in \mathbb{F} / \pm G$ is defined.

We make M_K a morphism $1 \rightarrow 0$ in the category Cob by choosing a boundary-parametrization $m : F(1, 0) \rightarrow \partial M_K$, such that $\lambda_- := m^{-1}(\lambda)$ is contained in the bottom surface F_1 and goes through the base point \star . Set $H_- := H_1^{\varphi_K m^-}(F_1, \star)$. The following proposition, which is easily deduced from Lemma 6.1, shows that the topological invariants $\tau^{\varphi_K}(M_K)$ and $\mathbf{R}(M_K, \varphi_K)$ are equivalent.

Proposition 6.3. *With the above notation and for any $h \in \Lambda^i H_-$, we have*

$$\mathbf{R}(M_K, \varphi_K)(h) = \begin{cases} \tau^{\varphi_K}(M_K) \cdot \partial_*(h) & \text{if } i = 1, \\ 0 & \text{otherwise,} \end{cases}$$

where $\partial_* : H_- \rightarrow \mathbb{F}$ is the connecting homomorphism for the pair (F_1, \star) . In particular, we have $\tau^{\varphi_K}(M_K) = R(M_K, \varphi_K)([\widehat{\lambda}_-]) / (\varphi_K([\lambda]) - 1)$.

Example 6.4. If G is the infinite cyclic group generated by t , N is a homology 3-sphere and φ_K maps the oriented meridian μ of K to t , then we know from [Mill] that $\tau^{\varphi_K}(M_K) = \Delta(K)/(t - 1)$. Thus we recover Proposition 3.3 by taking $\lambda := \mu$.

6.4. The situation of closed 3-manifolds . Let N be a closed connected orientable 3-manifold, and let $\varphi : H_1(N) \rightarrow G$ be a non-trivial group homomorphism. We wish to compute the Reidemeister torsion $\tau^\varphi(N)$ with coefficients $\varphi : \mathbb{Z}[H_1(N)] \rightarrow \mathbb{F}$ from the Reidemeister functor R . For this, we have to transform N into a cobordism. Note that removing an open 3-ball B from N and regarding $N \setminus B$ as an element of $\text{Cob}(0, 0)$ is not fruitful, since the functor R maps this morphism to zero (see Remark 6.2).

We proceed in the following (rather indirect) way. Choose a knot $K \subset N$ such that $\varphi([K]) \neq 1$. Consider the complement M_K of an open tubular neighborhood of K in N , and fix a parallel $\rho \subset \partial M_K$ of K . Let $\varphi_K : H_1(M_K) \rightarrow G$ be the homomorphism obtained from φ by restriction to $M_K \subset N$. Make M_K a morphism $1 \rightarrow 0$ in Cob by choosing a boundary-parametrization $m : F(1, 0) \rightarrow \partial M_K$ such that $\rho_- := m^{-1}(\rho)$ is contained in the bottom surface F_1 and $\star \in \rho_-$.

Proposition 6.5. *With the above notation, we have*

$$\tau^\varphi(N) = \frac{R(M_K, \varphi_K)([\widehat{\rho}_-])}{(\varphi([K]) - 1)^2} \in \mathbb{F} / \pm G.$$

Proof. There is a formula describing (under certain circumstances) how the abelian Reidemeister torsion changes when a solid torus is glued along a 3-manifold with toroidal boundary: see [Tur5, §VII.1]. This formula applies to our situation and gives

$$\tau^{\varphi_K}(M_K) = (\varphi([K]) - 1) \cdot \tau^\varphi(N).$$

We conclude by applying Proposition 6.3 to $\lambda := \rho$. □

As an application, we relate the functor A to the Alexander polynomial of closed 3-manifolds. Thus, we now assume that G is a finitely generated free abelian group and we take $\mathbb{F} := Q(G)$. We consider the *Alexander polynomial* of N with coefficients φ , namely

$$\Delta^\varphi(N) = \Delta_0 H_1^{\varphi_{\mathbb{Z}}}(N) \in \mathbb{Z}[G] / \pm G$$

where $\varphi_{\mathbb{Z}} : \mathbb{Z}[H_1(N)] \rightarrow \mathbb{Z}[G]$ is the extension of $\varphi : H_1(N) \rightarrow G$.

Proposition 6.6. *With the above notation, we have*

$$\Delta^\varphi(N) = \begin{cases} \frac{A(M_K, \varphi_K)([\widehat{\rho}_-])}{(\varphi([K]) - 1)^2} & \text{if } \text{rank } \varphi(H_1(N)) \geq 2, \\ \frac{A(M_K, \varphi_K)([\widehat{\rho}_-])}{(t^{n-1} + \dots + t + 1)^2} & \text{if } \text{rank } \varphi(H_1(N)) = 1. \end{cases}$$

In the second case, $t \in \varphi(H_1(N))$ is a generator and $n \in \mathbb{N}$ is such that $\varphi([K]) = t^n$.

Proof. Proposition 6.5 and Theorem 5.4 give

$$(6.5) \quad \tau^\varphi(N) = \frac{R(M_K, \varphi_K)([\widehat{\rho}_-])}{(\varphi([K]) - 1)^2} = \frac{A(M_K, \varphi_K)([\widehat{\rho}_-])}{(\varphi([K]) - 1)^2} \in Q(G)/\pm G.$$

Besides, according to [Tur1], we have

$$(6.6) \quad \tau^\varphi(N) = \begin{cases} \Delta^\varphi(N) & \text{if } \text{rank } \varphi(H_1(N)) \geq 2, \\ \Delta^\varphi(N)/(t - 1)^2 & \text{if } \text{rank } \varphi(H_1(N)) = 1. \end{cases}$$

We conclude by combining (6.5) with (6.6). □

7. The monoid of homology cobordisms

In this section, we fix an integer $k \geq 1$, an abelian group G and a group homomorphism $\psi : H_1(F_k) \rightarrow G$. We shall compute the functors A and R on the monoid of homology cobordisms over the surface F_k .

7.1. Homology cobordisms. A *homology cobordism* over F_k is a morphism $M : k \rightarrow k$ in the category Cob such that $m_\pm : H_1(F_k) \rightarrow H_1(M)$ is an isomorphism. The set of equivalence classes of homology cobordisms defines a submonoid

$$\mathcal{C}(F_k) \subset \text{Cob}(k, k).$$

We restrict ourselves to homology cobordisms M such that the composition

$$H_1(F_k) \xrightarrow[\simeq]{m_-} H_1(M) \xrightarrow[\simeq]{m_+^{-1}} H_1(F_k) \xrightarrow{\psi} G$$

coincides with ψ . Thus we obtain a submonoid

$$\mathcal{C}^\psi(F_k) \subset \mathcal{C}(F_k),$$

which we also view as a submonoid of $\text{Cob}_G((k, \psi), (k, \psi))$ by equipping every cobordism M of the above form with the homomorphism $\psi := \psi \circ m_-^{-1} = \psi \circ m_+^{-1} : H_1(M) \rightarrow G$.

Example 7.1. A *homology cylinder* is a homology cobordism M over F_k such that $m_- = m_+ : H_1(F_k) \rightarrow H_1(M)$. Homology cylinders constitute a submonoid $\mathcal{IC}(F_k)$ of $\mathcal{C}(F_k)$ such that $\mathcal{IC}(F_k) \subset \mathcal{C}^\psi(F_k)$, whatever ψ is.

7.2. The Magnus representation. Assume now that G is a multiplicative subgroup of a field \mathbb{F} . The extension of $\psi : H_1(F_k) \rightarrow G$ to a ring homomorphism $\mathbb{Z}[H_1(F_k)] \rightarrow \mathbb{F}$ is still denoted by ψ . We set

$$H^\psi := H_1^\psi(F_k, \star)$$

and, when we are given an $M \in \mathcal{C}^\psi(F_k)$, we denote $H := H_1^\psi(M, I)$. The fact that the map $m_\pm : H_1(F_k) \rightarrow H_1(M)$ is an isomorphism of abelian groups implies that $m_\pm : H^\psi \rightarrow H$ is an isomorphism of \mathbb{F} -vector spaces. (See [KLW, Proposition 2.1] for a similar statement.) Consequently, we are allowed to set $r^\psi(M) := m_+^{-1} \circ m_- : H^\psi \rightarrow H^\psi$. This results in a monoid homomorphism

$$r^\psi : \mathcal{C}^\psi(F_k) \longrightarrow \text{Aut}(H^\psi),$$

which is called the *Magnus representation*. See [Sak3] for a survey of this invariant.

7.3. The restriction of \mathbf{R} to homology cobordisms. The Reidemeister functor restricts to a monoid homomorphism

$$\mathbf{R} : \mathcal{C}^\psi(F_k) \longrightarrow \text{grVect}_{\mathbb{F}, \pm G}(\Lambda H^\psi, \Lambda H^\psi).$$

We now compute this projective representation of the monoid $\mathcal{C}^\psi(F_k)$.

Proposition 7.2. *For any $M \in \mathcal{C}^\psi(F_k)$ with top surface $\partial_+ M$, we have*

$$\mathbf{R}(M, \psi) = \tau^\psi(M, \partial_+ M) \cdot \Lambda(r^\psi(M)) : \Lambda H^\psi \longrightarrow \Lambda H^\psi$$

where $\tau^\psi(M, \partial_+ M)$ is the Reidemeister torsion of $(M, \partial_+ M)$ as defined in §6.1.

Proof. We shall prove a slightly more general statement: let $\psi_\pm : H_1(F_k) \rightarrow G$ be any group homomorphisms and assume that $M \in \mathcal{C}(F_k)$ is a cobordism such that $\psi_- \circ m_-^{-1} = \psi_+ \circ m_+^{-1} : H_1(M) \rightarrow G$. Then we claim that

$$(7.1) \quad \mathbf{R}(M, \psi) = \tau^\psi(M, \partial_+ M) \cdot \Lambda(m_+^{-1} \circ m_-) : \Lambda H_- \longrightarrow \Lambda H_+$$

where $H_\pm := H_1^{\psi_\pm}(F_k, \star)$ and $\psi := \psi_\pm \circ m_\pm^{-1}$. (The proposition is the particular case where $\psi_+ = \psi_- : H_1(F_k) \rightarrow G$.)

To prove this claim, we set $g := g(M) = 2k$, $H := H_1^\psi(M, I)$ and let $h = (h_1, \dots, h_g)$ be a basis of H . In order to compute $\mathcal{R}_M^\psi(h_1 \wedge \dots \wedge h_g)$, we consider the short exact sequence of \mathbb{F} -chain complexes:

$$(7.2) \quad 0 \longrightarrow \underbrace{C^{\psi_+}(F_k, \star)}_{C' :=} \xrightarrow{m_+} \underbrace{C^\psi(M, \star)}_{C :=} \longrightarrow \underbrace{C^\psi(M, \partial_+ M)}_{C'' :=} \longrightarrow 0.$$

The complex C'' is acyclic while C' and C have their homology concentrated in degree 1. Therefore, the long exact sequence in homology \mathcal{H} induced by (7.2) is concentrated in degrees 4 and 5 where it reduces to the map $m_+ : H_+ = H_1(C') \rightarrow H_1(C) = H$.

There exists a wedge of circles $S_1 \vee \dots \vee S_g$ based at \star onto which the surface F_k retracts by elementary collapses. Let $h' = (h'_1, \dots, h'_g)$ be the basis of H_+ obtained by lifting each of the loops S_1, \dots, S_g to the maximal abelian cover of F_k . Then we have

$$\tau(C'; c', h') \in \pm G \subset \mathbb{F}$$

for any choice of ordered/oriented lifts of the relative cells of (F_k, \star) inducing a basis c' of C' . Besides, by the multiplicativity property of torsions (see Theorem A.3), we have

$$\tau(C; c, h) = \varepsilon \cdot \tau(C'; c', h') \cdot \tau(C''; c'') \cdot \tau(\mathcal{H}; (h', h)) \in \mathbb{F} \setminus \{0\}$$

for some appropriate choices of ordered/oriented lifts of the relative cells, which result in bases c', c, c'' of the chain complexes. Here ε is a sign not depending on h and \mathcal{H} is regarded as an acyclic \mathbb{F} -chain complex based by (h', h) . We deduce that

$$\begin{aligned} \mathcal{R}_M^\psi(h_1 \wedge \dots \wedge h_g) &= \tau(C; c, h) = \tau^\psi(M, \partial_+ M) \cdot [m_+(h')/h]^{(-1)^{4+1}} \\ &= \tau^\psi(M, \partial_+ M) \cdot [h/m_+(h')]. \end{aligned}$$

(Here the identities are up to multiplication by an element of $\pm G$ not depending on h .)

To proceed, we consider any integer $j \geq 0$ and any $x \in \Lambda^j H_-$. Let $\omega : \Lambda^g H_+ \rightarrow \mathbb{F}$ be the volume form defined by $\omega(h'_1 \wedge \dots \wedge h'_g) = 1$. (Note that ω is integral.) Then, for any $y \in \Lambda^{g-j} H_+$, we have

$$\begin{aligned} \omega(\mathbf{R}(M, \psi)(x) \wedge y) &= \mathcal{R}_M^\psi(\Lambda^j m_-(x) \wedge \Lambda^{g-j} m_+(y)) \\ &= \tau^\psi(M, \partial_+ M) \cdot [(\Lambda^j m_-(x) \wedge \Lambda^{g-j} m_+(y))/m_+(h')] \\ &= \tau^\psi(M, \partial_+ M) \cdot [(\Lambda^j (m_+^{-1} m_-)(x) \wedge y)/h'] \\ &= \tau^\psi(M, \partial_+ M) \cdot \omega(\Lambda^j (m_+^{-1} m_-)(x) \wedge y). \end{aligned}$$

We conclude that $\mathbf{R}(M, \psi)(x) = \tau^\psi(M, \partial_+ M) \cdot \Lambda^j (m_+^{-1} m_-)(x)$ up to multiplication by an element of $\pm G$ not depending on x , which proves (7.1). \square

7.4. The restriction of A to homology cobordisms. Assume now that the abelian group G is finitely generated and free, and assume that $\mathbb{F} := Q(G)$. We denote by $\psi_{\mathbb{Z}} : \mathbb{Z}[H_1(F_k)] \rightarrow \mathbb{Z}[G]$ the extension of $\psi : H_1(F_k) \rightarrow G$ to a ring homomorphism and we set $H_{\mathbb{Z}}^{\psi} := H_1^{\psi_{\mathbb{Z}}}(F_k, \star)$. The Alexander functor restricts to a monoid homomorphism

$$A : \mathcal{C}^{\psi}(F_k) \longrightarrow \text{grMod}_{\mathbb{Z}[G], \pm G}(\Lambda H_{\mathbb{Z}}^{\psi}, \Lambda H_{\mathbb{Z}}^{\psi}).$$

This projective representation of the monoid $\mathcal{C}^{\psi}(F_k)$ is computed as follows.

Proposition 7.3. *For any $M \in \mathcal{C}^{\psi}(F_k)$, we have the commutative diagram*

$$\begin{array}{ccc} \Lambda H_{\mathbb{Z}}^{\psi} & \xrightarrow{A(M, \psi)} & \Lambda H_{\mathbb{Z}}^{\psi} \\ \downarrow & & \downarrow \\ \Lambda H^{\psi} & \xrightarrow{\Delta^{\psi}(M, \partial_+ M) \cdot \Lambda r^{\psi}(M)} & \Lambda H^{\psi} \end{array}$$

where $\Delta^{\psi}(M, \partial_+ M)$ is the Alexander polynomial of the pair $(M, \partial_+ M)$ as defined in §3.1.

Proof. The proposition can be proved directly from the definition of A , using an appropriate presentation of the $\mathbb{Z}[G]$ -module $H_1^{\psi_{\mathbb{Z}}}(M, I)$. It also follows from Theorem 5.4, Proposition 7.2 and the fact that

$$\tau^{\psi}(M, \partial_+ M) = \Delta^{\psi}(M, \partial_+ M) \in \mathbb{Z}[G]/\pm G.$$

The latter identity is shown using the fact that M collapses, relatively to $\partial_+ M$, onto a cell complex having only 1-cells and 2-cells in equal number. (For instance, consider the CW-complex resulting from a handle decomposition of M as discussed in §8.1.) Thus, the computation of both invariants $\tau^{\psi}(M, \partial_+ M)$ and $\Delta^{\psi}(M, \partial_+ M)$ reduces to the determinant of a same matrix. (See [FJR, Lemma 3.6] for instance.) \square

Example 7.4. Assume that $G := \{1\}$ is the trivial group. Then $\mathcal{C}^{\psi}(F_k) = \mathcal{C}(F_k)$. Moreover $\mathbb{Z}[G] = \mathbb{Z}$ and $Q(G) = \mathbb{Q}$, so that $H_{\mathbb{Z}}^{\psi} = H_1(F_k)$ and $H^{\psi} = H_1(F_k; \mathbb{Q})$. Note that $\Delta^{\psi}(M, \partial_+ M) = 1$ since $H_1^{\psi_{\mathbb{Z}}}(M, \partial_+ M) = H_1(M, \partial_+ M)$ is trivial in that case. It follows from Proposition 7.3 that $A(M, \psi) : \Lambda H_1(F_k) \rightarrow \Lambda H_1(F_k)$ is induced by the isomorphism $(m_+)^{-1}m_- : H_1(F_k) \rightarrow H_1(F_k)$.

Remark 7.5. If two cobordisms $M, M' \in \mathcal{C}^{\psi}(F_k)$ are homology cobordant, then we have $r^{\psi}(M) = r^{\psi}(M')$ (see [Sak2, Theorem 3.6]), but it may happen that $\Delta^{\psi}(M, \partial_+ M) \neq \Delta^{\psi}(M', \partial_+ M')$ (see [MM, Lemma 3.15] for an example). It follows from Proposition 7.3 that the restriction of A to $\mathcal{C}^{\psi}(F_k)$ is *stronger* than the representation r^{ψ} .

8. Computations with Heegaard splittings

Let G be a multiplicative subgroup of a field \mathbb{F} . We give a simple recipe to compute the functor $R = R_{\mathbb{F},G}$ using Heegaard splittings of cobordisms. In this section, the extension of a group homomorphism $\rho : H \rightarrow G$ to a ring homomorphism $\mathbb{Z}[H] \rightarrow \mathbb{F}$ is still denoted by ρ .

8.1. Heegaard splittings. In order to obtain concrete formulas for the functor R , it is convenient to fix compatible systems of “meridians and parallels” on the model surfaces. Specifically, we choose on the model surface F_1 a *meridian* α and a *parallel* β , which means the following: α and β are oriented simple closed curves in the interior of F_1 meeting transversely at a single point with homological intersection $[\alpha] \bullet [\beta] = +1$. Then the identification between $F_1 \#_{\partial} \cdots \#_{\partial} F_1$ and F_k induces, for any integer $k \geq 1$, a *system of meridians and parallels* $(\alpha_1, \dots, \alpha_k, \beta_1, \dots, \beta_k)$ on the surface F_k .

For any $k \geq 0$, we denote by $C_0^k \in \text{Cob}(0, k)$ the cobordism obtained from $F_k \times [-1, 1]$ by attaching k 2-handles along the curves $\alpha_1 \times \{-1\}, \dots, \alpha_k \times \{-1\}$. Similarly, let $C_k^0 \in \text{Cob}(k, 0)$ be the cobordism obtained from $F_k \times [-1, 1]$ by attaching k 2-handles along the curves $\beta_1 \times \{1\}, \dots, \beta_k \times \{1\}$. Observe that $C_k^0 \circ C_0^k = C_0^0 \in \text{Cob}(0, 0)$ is the 3-dimensional ball $F_0 \times [-1, 1]$. Thus we shall refer to C_k^0 and C_0^k as the *upper* and *lower* handlebodies, respectively. (Clearly, these notions depend on the above choice of meridians and parallels.)

Let also $\mathcal{M}(F_k)$ be the *mapping class group* of the surface F_k , which consists of isotopy classes of (orientation-preserving) homeomorphisms $f : F_k \rightarrow F_k$ fixing ∂F_k pointwisely. The *mapping cylinder* construction, which associates to any such homeomorphism f the cobordism

$$\mathbf{c}(f) := (F_k \times [-1, 1], (f \times \{-1\}) \cup (\partial F_k \times \text{Id}) \cup (\text{Id} \times \{1\})),$$

defines an embedding $\mathbf{c} : \mathcal{M}(F_k) \rightarrow \mathcal{C}(F_k)$ of the mapping class group into the monoid of homology cobordisms (see §7.1).

Let $M \in \text{Cob}(g_-, g_+)$ be an arbitrary cobordism. By elementary Morse theory, the 3-manifold underlying M can be obtained from the trivial cobordism $F_{g_+} \times [-1, 1]$ by attaching simultaneously some 1-handles (say, $r_+ \geq 0$) along the “bottom surface” $F_{g_+} \times \{-1\}$, and by attaching subsequently some 2-handles (say, $r_- \geq 0$) along the new “bottom surface.” We obtain in that way a *Heegaard splitting* of M , i.e. a decomposition in the monoidal category Cob of the form

$$(8.1) \quad M = \left(C_{r_+}^0 \otimes \text{Id}_{g_+} \right) \circ \mathbf{c}(f) \circ \left(C_0^{r_-} \otimes \text{Id}_{g_-} \right)$$

where $g_+ + r_+ = g_- + r_-$ and $f \in \mathcal{M}(F_{g_{\pm} + r_{\pm}})$. See [Ker2, Theorem 5].

8.2. Computation of R with Heegaard splittings. We now assume that the above cobordism M comes with a group homomorphism $\varphi : H_1(M) \rightarrow G$:

$$(M, \varphi) \in \text{Cob}_G((g_-, \varphi_-), (g_+, \varphi_+)).$$

The Heegaard splitting (8.1) of M induces a decomposition in the monoidal category Cob_G by endowing each submanifold S of that decomposition with the group homomorphism $\bar{\varphi} : H_1(S) \rightarrow G$ obtained by restricting φ to $S \subset M$. Hence we obtain

$$R(M, \varphi) = \left(R(C_{r_+}^0, \bar{\varphi}) \otimes \text{Id}_{\Lambda H_+} \right) \circ R(\mathbf{c}(f), \bar{\varphi}) \circ \left(R(C_0^{r_-}, \bar{\varphi}) \otimes \text{Id}_{\Lambda H_-} \right)$$

where $H_{\pm} := H_1^{\varphi_{\pm}}(F_{g_{\pm}}, \star)$ and the symbol $\bar{\varphi}$ denotes a representation in G induced by φ . Thus the computation of $R(M, \varphi)$ reduces to three cases: upper handlebodies, lower handlebodies and mapping cylinders.

To describe the values of R in those three cases, we need to fix further notation. Let $k \geq 0$ be an integer and let $\psi : H_1(F_k) \rightarrow G$ be a group homomorphism. We assume that, in our model surface F_1 , the intersection point $\alpha \cap \beta$ is connected by an arc to the base point $\star \in \partial F_1$: hence the curves $\alpha_1, \dots, \alpha_k, \beta_1, \dots, \beta_k$ are now viewed as oriented loops based at $\star \in \partial F_k$. We denote by $(a_1^{\psi}, \dots, a_k^{\psi}, b_1^{\psi}, \dots, b_k^{\psi})$ the basis of $H_1^{\psi}(F_k, \star)$ obtained by lifting these loops to the maximal abelian cover:

$$(8.2) \quad \forall i = 1, \dots, k, \quad a_i^{\psi} := [1 \otimes \hat{\alpha}_i], \quad b_i^{\psi} := [1 \otimes \hat{\beta}_i].$$

Then the space $\Lambda H_1^{\psi}(F_k, \star)$ can be identified to $\Lambda A_k^{\psi} \otimes \Lambda B_k^{\psi}$ where $A_k^{\psi} := \langle a_1^{\psi}, \dots, a_k^{\psi} \rangle$ and $B_k^{\psi} := \langle b_1^{\psi}, \dots, b_k^{\psi} \rangle$ are the subspaces of $H_1^{\psi}(F_k, \star)$ corresponding to meridians and parallels, respectively.

Lemma 8.1. *Let $\psi : H_1(C_k^0) \rightarrow G$ be a group homomorphism and let $\psi_- : H_1(F_k) \rightarrow G$ be the restriction of ψ to $F_k \subset \partial C_k^0$. Then the linear map*

$$R(C_k^0, \psi) : \Lambda H_1^{\psi_-}(F_k, \star) \longrightarrow \mathbb{F}$$

is trivial on $\Lambda^i A_k^{\psi_-} \otimes \Lambda^j B_k^{\psi_-}$ if $i \neq k$ or $j \neq 0$, and it sends $a_1^{\psi_-} \wedge \dots \wedge a_k^{\psi_-}$ to 1.

Proof. Set $N := C_k^0 \in \text{Cob}(k, 0)$. Since $R(N, \psi)$ has degree $-k$, it must be trivial on $\Lambda^r H_1^{\psi_-}(F_k, \star)$ for $r \neq k$. It remains to compute

$$(8.3) \quad R(N, \psi)(x_1 \wedge \dots \wedge x_k) = \mathcal{R}_N^{\psi}(n_-(x_1) \wedge \dots \wedge n_-(x_k))$$

for any $x_1, \dots, x_k \in H_1^{\psi_-}(F_k, \star)$. If one of the x_i 's belongs to $B_k^{\psi_-}$, the right-hand side of (8.3) is zero since, for all $j \in \{1, \dots, k\}$, β_j bounds a disk in N

so that $n_-(b_j^{\psi^-}) = 0$. So, we can assume that $x_1 \wedge \cdots \wedge x_k = a_1^{\psi^-} \wedge \cdots \wedge a_k^{\psi^-}$. In this case, we apply Lemma 5.1 to the obvious spine $X = X^+$ of N : the spine X is a wedge of circles whose 1-cells $\gamma_1, \dots, \gamma_k$ are obtained by “pushing” the curves $\alpha_1, \dots, \alpha_k$ in the interior of N . We deduce that the right-hand side of (8.3) is equal to 1. \square

Lemma 8.2. *Let $\psi : H_1(C_0^k) \rightarrow G$ be a group homomorphism and let $\psi_+ : H_1(F_k) \rightarrow G$ be the restriction of ψ to $F_k \subset \partial C_0^k$. Then the linear map*

$$R(C_0^k, \psi) : \mathbb{F} \longrightarrow \Lambda H_1^{\psi_+}(F_k, \star)$$

sends the scalar 1 to the multivector $a_1^{\psi_+} \wedge \cdots \wedge a_k^{\psi_+}$.

Proof. Set $(v_1, \dots, v_k, v_{k+1}, \dots, v_{2k}) := (a_1^{\psi_+}, \dots, a_k^{\psi_+}, b_1^{\psi_+}, \dots, b_k^{\psi_+})$ and let ω be the volume form on $H_1^{\psi_+}(F_k, \star)$ defined by $\omega(v_1 \wedge \cdots \wedge v_{2k}) = 1$. We denote $N := C_0^k \in \text{Cob}(0, k)$ and write

$$R(N, \psi)(1) = \sum_P z_P \cdot v_P \in \Lambda^k H_1^{\psi_+}(F_k, \star)$$

where P runs over k -element subsets of $\{1, \dots, 2k\}$ and v_P is the wedge of the v_p 's for all $p \in P$. For any k -element subset $P \subset \{1, \dots, 2k\}$, we have

$$(8.4) \quad \varepsilon_P \cdot z_P = \omega(R(N, \psi)(1) \wedge v_{\overline{P}}) = \mathcal{R}_N^{\psi}(\Lambda^k n_+(v_{\overline{P}}))$$

where \overline{P} is the complement of P and ε_P is the signature of the permutation $P\overline{P}$. To compute the right-hand side of (8.4), we apply Lemma 5.1 to the obvious spine $X = X^+$ of N : the spine X is a wedge of circles whose 1-cells $\gamma_1, \dots, \gamma_k$ are obtained by “pushing” the curves β_1, \dots, β_k in the interior of N . We obtain that $\mathcal{R}_N^{\psi}(\Lambda^k n_+(v_{\overline{P}}))$ is trivial except if $\overline{P} = \{k+1, \dots, 2k\}$, in which case it takes the value 1. We conclude that $z_P = 1$ if $P = \{1, \dots, k\}$ and $z_P = 0$ otherwise. \square

Lemma 8.3. *Let $f \in \mathcal{M}(F_k)$ and let $\psi_{\pm} : H_1(F_k) \rightarrow G$ be group homomorphisms such that $\psi_- = \psi_+ \circ f$. Denote by $\psi : H_1(F_k \times [-1, 1]) \rightarrow G$ the isomorphism $\psi_+ \circ \text{pr}$, where $\text{pr} : F_k \times [-1, 1] \rightarrow F_k$ is the cartesian projection. Then*

$$R(\mathbf{c}(f), \psi) : \Lambda H_1^{\psi_-}(F_k, \star) \longrightarrow \Lambda H_1^{\psi_+}(F_k, \star)$$

is induced by the isomorphism $f : H_1^{\psi-}(F_k, \star) \rightarrow H_1^{\psi+}(F_k, \star)$. Moreover, the matrix of this isomorphism in the bases $(a_1^{\psi\pm}, \dots, a_k^{\psi\pm}, b_1^{\psi\pm}, \dots, b_k^{\psi\pm})$ of $H_1^{\psi\pm}(F_k, \star)$ is

$$\psi_+ \begin{pmatrix} \frac{\partial f_*(\alpha_1)}{\partial \alpha_1} & \dots & \frac{\partial f_*(\alpha_k)}{\partial \alpha_1} & \frac{\partial f_*(\beta_1)}{\partial \alpha_1} & \dots & \frac{\partial f_*(\beta_k)}{\partial \alpha_1} \\ \vdots & & \vdots & \vdots & & \vdots \\ \frac{\partial f_*(\alpha_1)}{\partial \alpha_k} & \dots & \frac{\partial f_*(\alpha_k)}{\partial \alpha_k} & \frac{\partial f_*(\beta_1)}{\partial \alpha_k} & \dots & \frac{\partial f_*(\beta_k)}{\partial \alpha_k} \\ \frac{\partial f_*(\alpha_1)}{\partial \beta_1} & \dots & \frac{\partial f_*(\alpha_k)}{\partial \beta_1} & \frac{\partial f_*(\beta_1)}{\partial \beta_1} & \dots & \frac{\partial f_*(\beta_k)}{\partial \beta_1} \\ \vdots & & \vdots & \vdots & & \vdots \\ \frac{\partial f_*(\alpha_1)}{\partial \beta_k} & \dots & \frac{\partial f_*(\alpha_k)}{\partial \beta_k} & \frac{\partial f_*(\beta_1)}{\partial \beta_k} & \dots & \frac{\partial f_*(\beta_k)}{\partial \beta_k} \end{pmatrix}$$

where $f_* : \pi_1(F_k, \star) \rightarrow \pi_1(F_k, \star)$ is induced by f .

Proof. The first statement follows from (7.1). The second statement is well known. \square

8.3. Computation of \mathbf{A} with Heegaard splittings. Assume now that G is a finitely generated free abelian group and take $\mathbb{F} := Q(G)$. There are counterparts of Lemmas 8.1, 8.2 and 8.3 for the Alexander functor \mathbf{A} . These counterparts follow from the same lemmas using Theorem 5.4, or they can be proved independently using presentations of $\mathbb{Z}[G]$ -modules.

For $G = \{1\}$, we deduce that the functor \mathbf{A} is essentially the same thing as the TQFT constructed in [FNI]. (Compare the formulas given in [Ker1, §3] with the above lemmas.) However, there are a few technical differences: in particular, we have considered surfaces with circle boundary, whereas [FNI] works with closed surfaces.

9. Duality

We prove two duality properties for the Reidemeister functor. In this section, \mathbb{F} is a field where a multiplicative subgroup G is fixed, and we assume that \mathbb{F} is equipped with an involutive automorphism $f \mapsto \bar{f}$ such that $\bar{g} = g^{-1}$ for all $g \in G$.

9.1. Twisted intersection form. The first duality satisfied by \mathbf{R} involves the “twisted” intersection forms of oriented surfaces with boundary. We start by recalling this notion.

Let $k \geq 0$ be an integer and set $\pi := \pi_1(F_k, \star)$. The *homotopy intersection form* of F_k is the pairing $\lambda : \mathbb{Z}[\pi] \times \mathbb{Z}[\pi] \rightarrow \mathbb{Z}[\pi]$ defined by Turaev in [Tur7].

We also refer to Papakyriakopoulos' work [Pap] where this form is implicit, and to Perron's work [Per] where the same form λ is re-discovered (and is denoted there by ω).

The twisted homology group $H_1(F_k, \star; \mathbb{Z}[\pi])$ is identified (as a left $\mathbb{Z}[\pi]$ -module) to the augmentation ideal $I(\pi)$ of $\mathbb{Z}[\pi]$ in the following way: for any oriented loop $\gamma \subset F_k$ based at \star , let $\tilde{\gamma}$ be the unique lift of γ to the universal cover of F_k starting at the preferred lift $\tilde{\star}$, and identify $[1 \otimes \tilde{\gamma}] \in H_1(F_k, \star; \mathbb{Z}[\pi])$ to $[\gamma] - 1 \in I(\pi)$. Thus, by restricting λ to $I(\pi) \times I(\pi)$, we obtain a pairing

$$\langle -, - \rangle : H_1(F_k, \star; \mathbb{Z}[\pi]) \times H_1(F_k, \star; \mathbb{Z}[\pi]) \longrightarrow \mathbb{Z}[\pi].$$

The derivation properties of λ given in [Tur7, Per] imply that $\langle -, - \rangle$ is *sesquilinear* in the sense that

$$\langle ax + y, z \rangle = a \langle x, z \rangle + \langle y, z \rangle, \quad \langle z, ax + y \rangle = \langle z, x \rangle S(a) + \langle z, y \rangle$$

for all $a \in \mathbb{Z}[\pi]$ and $x, y, z \in H_1(F_k, \star; \mathbb{Z}[\pi])$. Here $S : \mathbb{Z}[\pi] \rightarrow \mathbb{Z}[\pi]$ is the *antipode*, i.e. the \mathbb{Z} -linear map defined by $S(a) = a^{-1}$ for all $a \in \pi$.

Let now $\psi : H_1(F_k) \rightarrow G$ be a group homomorphism: this induces a structure of right $\mathbb{Z}[\pi]$ -module on \mathbb{F} . By identifying $H_1^\psi(F_k, \star)$ to $\mathbb{F} \otimes_{\mathbb{Z}[\pi]} H_1(F_k, \star; \mathbb{Z}[\pi])$, we obtain a pairing

$$(9.1) \quad \langle -, - \rangle : H_1^\psi(F_k, \star) \times H_1^\psi(F_k, \star) \longrightarrow \mathbb{F}$$

defined by $\langle f_1 \otimes h_1, f_2 \otimes h_2 \rangle := f_1 \overline{f_2} \psi(\langle h_1, h_2 \rangle)$ for all $f_1, f_2 \in \mathbb{F}$ and $h_1, h_2 \in H_1(F_k, \star; \mathbb{Z}[\pi])$. This pairing is *sesquilinear* in the sense that

$$\langle fx + y, z \rangle = f \langle x, z \rangle + \langle y, z \rangle, \quad \langle z, fx + y \rangle = \overline{f} \langle z, x \rangle + \langle z, y \rangle$$

for all $f \in \mathbb{F}$ and $x, y, z \in H_1^\psi(F_k, \star)$. The pairing (9.1) can also be defined using Poincaré duality (with twisted coefficients) and the fact that $H_1^\psi(F_k, J) \simeq H_1^\psi(F_k, \star) \simeq H_1^\psi(F_k, J')$, where J, J' are two closed intervals such that $J \cup J' = \partial F_k$ and $J \cap J' = \partial J = \partial J'$. In particular, the pairing (9.1) is *non-singular* in the sense that $\langle x, - \rangle : H_1^\psi(F_k, \star) \rightarrow \text{Hom}(H_1^\psi(F_k, \star), \mathbb{F})$ is an isomorphism for any $x \in H_1^\psi(F_k, \star)$.

For any integer $r \geq 1$, the pairing (9.1) also induces a non-singular sesquilinear pairing $\langle -, - \rangle : \Lambda^r H_1^\psi(F_k, \star) \times \Lambda^r H_1^\psi(F_k, \star) \rightarrow \mathbb{F}$ defined by

$$\langle x_1 \wedge \cdots \wedge x_r, y_1 \wedge \cdots \wedge y_r \rangle = \det \begin{pmatrix} \langle x_1, y_1 \rangle & \cdots & \langle x_1, y_r \rangle \\ \vdots & \ddots & \vdots \\ \langle x_r, y_1 \rangle & \cdots & \langle x_r, y_r \rangle \end{pmatrix}$$

for all $x_1, \dots, x_r, y_1, \dots, y_r \in H_1^\psi(F_k, \star)$. For $r = 0$, we set $\langle x, y \rangle := x \overline{y}$ for all $x, y \in \mathbb{F}$.

Remark 9.1. The sesquilinear pairing (9.1) is not skew-hermitian. Instead, we have

$$(9.2) \quad \forall x, y \in H_1^\psi(F_k, \star), \quad \langle x, y \rangle = -\overline{\langle y, x \rangle} + \partial_*(x) \overline{\partial_*(y)}$$

where $\partial_* : H_1^\psi(F_k, \star) \rightarrow \mathbb{F}$ is the connecting homomorphism in the long exact sequence of the pair (F_k, \star) . This identity follows from a similar property for the homotopy intersection form λ : see [Tur7, Per].

9.2. First duality. Let $g_-, g_+ \geq 0$ be integers. The *dual* of an $M \in \text{Cob}(g_-, g_+)$ is the cobordism $\overline{M} \in \text{Cob}(g_+, g_-)$ obtained from M by reversing its orientation and by composing its boundary-parametrization $m : F(g_-, g_+) \rightarrow \partial M$ with the orientation-reversing homeomorphism between

$$\underbrace{-F_{g_+} \cup_{S^1 \times \{-1\}} (S^1 \times [-1, 1]) \cup_{S^1 \times \{1\}} F_{g_-}}_{F(g_+, g_-)}$$

and

$$\underbrace{-F_{g_-} \cup_{S^1 \times \{-1\}} (S^1 \times [-1, 1]) \cup_{S^1 \times \{1\}} F_{g_+}}_{F(g_-, g_+)},$$

which is given by “time-reversal” $(x, t) \mapsto (x, -t)$ on the annulus $S^1 \times [-1, 1]$ and by the identity on F_{g_+} and F_{g_-} .

Theorem 9.2. For any $(M, \varphi) \in \text{Cob}_G((g_-, \varphi_-), (g_+, \varphi_+))$ and for any $j \geq 0$, we have

$$(9.3) \quad \forall x \in \Lambda^j H_-, \quad \forall y \in \Lambda^{j+\delta g} H_+, \quad \langle \mathbb{R}(M, \varphi)(x), y \rangle = \langle x, \mathbb{R}(\overline{M}, \varphi)(y) \rangle$$

where $\delta g := g_+ - g_-$ and $H_\pm := H_1^{\varphi_\pm}(F_{g_\pm}, \star)$.

Of course, the identity (9.3) only holds true up to multiplication by a constant in $\pm G$ (independent of x and y). The pairing $\langle -, - \rangle$ denotes the twisted intersection form of H_+ (respectively, H_-) on the left-hand side (respectively, the right-hand side) of (9.3).

Proof of Theorem 9.2. Assume that $(M, \varphi) = (M', \varphi') \circ (M'', \varphi'')$ where (M', φ') and (M'', φ'') are two morphisms in Cob_G satisfying (9.3). Then the dual of M is $\overline{M}'' \circ \overline{M}'$, and it easily follows that (M, φ) also satisfies (9.3). Consequently, and following the discussion of §8, it is enough to prove (9.3) in the following three cases: (i) M is a mapping cylinder; (ii) M is a “stabilized” lower handlebody; (iii) M is a “stabilized” upper handlebody.

Case (i). Assume that $g_- = g_+ =: k$ and that $M = \mathbf{c}(f)$ is the mapping cylinder of an $f \in \mathcal{M}(F_k)$. Then $\overline{M} = \mathbf{c}(f^{-1})$. Since $\varphi_+ f = \varphi_- : H_1(F_k) \rightarrow G$ and since $f_* : \pi_1(F_k, \star) \rightarrow \pi_1(F_k, \star)$ preserves the homotopy intersection form, the isomorphism $f : H_- \rightarrow H_+$ induced by $f : F_k \rightarrow F_k$ preserves the pairings (9.1). Using the first statement of Lemma 8.3, we obtain (9.3) as follows:

$$\begin{aligned} \forall x \in \Lambda^j H_-, \quad \forall y \in \Lambda^j H_+, \quad \langle \mathbf{R}(M, \varphi)(x), y \rangle &= \langle \Lambda^j f(x), y \rangle \\ &= \langle x, \Lambda^j f^{-1}(y) \rangle \\ &= \langle x, \mathbf{R}(\overline{M}, \varphi)(y) \rangle. \end{aligned}$$

Interlude. In order to deal with cases (ii) and (iii), we need an explicit formula for the twisted intersection form $\langle -, - \rangle : H_1^\psi(F_k, \star) \times H_1^\psi(F_k, \star) \rightarrow \mathbb{F}$ defined by a group homomorphism $\psi : H_1(F_k) \rightarrow G$. For this, we fix a system of meridians and parallels $(\alpha_1, \dots, \alpha_k, \beta_1, \dots, \beta_k)$ on F_k as explained in §8.1, and we denote by $(a_1^\psi, \dots, a_k^\psi, b_1^\psi, \dots, b_k^\psi)$ the corresponding basis of $H_1^\psi(F_k, \star)$: see (8.2). For every $x, y \in H_1(F_k)$, set $P^\psi(x, y) := (1 - \psi(x))(1 - \psi(y)) \in \mathbb{F}$. Then, for an appropriate choice of meridians and parallels, the matrix of $\langle -, - \rangle$ in the basis $(a_1^\psi, \dots, a_k^\psi, b_1^\psi, \dots, b_k^\psi)$ is

$$J^\psi = \left(\begin{array}{c|c} J_{aa}^\psi & J_{ab}^\psi \\ \hline J_{ba}^\psi & J_{bb}^\psi \end{array} \right)$$

where $J_{aa}^\psi, J_{ab}^\psi, J_{ba}^\psi, J_{bb}^\psi$ are the following lower triangular matrices [Per, Lemma 2.4]:

$$(9.4) \quad J_{aa}^\psi = \begin{pmatrix} 1 - \psi(\alpha_1) & 0 & 0 & \cdots & 0 \\ P^\psi(\alpha_2, \alpha_1) & 1 - \psi(\alpha_2) & 0 & \cdots & 0 \\ P^\psi(\alpha_3, \alpha_1) & P^\psi(\alpha_3, \alpha_2) & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & 0 \\ P^\psi(\alpha_k, \alpha_1) & P^\psi(\alpha_k, \alpha_2) & \cdots & P^\psi(\alpha_k, \alpha_{k-1}) & 1 - \psi(\alpha_k) \end{pmatrix},$$

$$(9.5) \quad J_{ab}^\psi = \begin{pmatrix} \psi(\alpha_1)\overline{\psi(\beta_1)} & 0 & 0 & \cdots & 0 \\ P^\psi(\alpha_2, \beta_1) & \psi(\alpha_2)\overline{\psi(\beta_2)} & 0 & \cdots & 0 \\ P^\psi(\alpha_3, \beta_1) & P^\psi(\alpha_3, \beta_2) & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & 0 \\ P^\psi(\alpha_k, \beta_1) & P^\psi(\alpha_k, \beta_2) & \cdots & P^\psi(\alpha_k, \beta_{k-1}) & \psi(\alpha_k)\overline{\psi(\beta_k)} \end{pmatrix},$$

$$(9.6) \quad J_{ba}^\psi = \begin{pmatrix} 1 - \overline{\psi(\alpha_1)} - \psi(\beta_1) & 0 & 0 & \cdots & 0 \\ P^\psi(\beta_2, \alpha_1) & 1 - \overline{\psi(\alpha_2)} - \psi(\beta_2) & 0 & \cdots & 0 \\ P^\psi(\beta_3, \alpha_1) & P^\psi(\beta_3, \alpha_2) & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & 0 \\ P^\psi(\beta_k, \alpha_1) & P^\psi(\beta_k, \alpha_2) & \cdots & P^\psi(\beta_k, \alpha_{k-1}) & 1 - \overline{\psi(\alpha_k)} - \psi(\beta_k) \end{pmatrix},$$

$$(9.7) \quad J_{bb}^\psi = \begin{pmatrix} 1 - \overline{\psi(\beta_1)} & 0 & 0 & \cdots & 0 \\ P^\psi(\beta_2, \beta_1) & 1 - \overline{\psi(\beta_2)} & 0 & \cdots & 0 \\ P^\psi(\beta_3, \beta_1) & P^\psi(\beta_3, \beta_2) & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & 0 \\ P^\psi(\beta_k, \beta_1) & P^\psi(\beta_k, \beta_2) & \cdots & P^\psi(\beta_k, \beta_{k-1}) & 1 - \overline{\psi(\beta_k)} \end{pmatrix}.$$

Besides, the following notation will be useful in the sequel. Let $\varepsilon \in \{+, -\}$ be a sign. We denote by $(v_1^\varepsilon, \dots, v_{g_\varepsilon}^\varepsilon, v_{g_\varepsilon+1}^\varepsilon, \dots, v_{2g_\varepsilon}^\varepsilon) := (a_1^{\varphi_\varepsilon}, \dots, a_{g_\varepsilon}^{\varphi_\varepsilon}, b_1^{\varphi_\varepsilon}, \dots, b_{g_\varepsilon}^{\varphi_\varepsilon})$ the basis of $H_\varepsilon = H_1^{\varphi_\varepsilon}(F_{g_\varepsilon}, \star)$. For any s -element subset $P \subset \{1, \dots, 2g_\varepsilon\}$, let $v_P^\varepsilon \in \Lambda^s H_\varepsilon$ be the wedge of the vectors v_p^ε 's for all $p \in P$ and, when this makes sense, we shall also denote by $(v_P^\varepsilon)^{-\varepsilon} \in \Lambda^s H_{-\varepsilon}$ the multivector obtained from v_P^ε by the transformations $a_i^{\varphi_\varepsilon} \mapsto a_{i-\varepsilon\delta g}^{\varphi_{-\varepsilon}}$ and $b_i^{\varphi_\varepsilon} \mapsto b_{i-\varepsilon\delta g}^{\varphi_{-\varepsilon}}$.

Case (ii). Assume that $M = C_0^r \otimes \text{Id}_{g_-}$ where $r = \delta g$. Note that $\varphi_+(\alpha_i) = 1$ for all $i \in \{1, \dots, r\}$, so that (9.4) and (9.5) applied to $\psi := \varphi_+$ give

$$(9.8) \quad \forall i \in \{1, \dots, r\}, \forall j \in \{1, \dots, r + g_-\}, \quad \langle a_i^{\varphi_+}, a_j^{\varphi_+} \rangle = 0, \quad \langle a_i^{\varphi_+}, b_j^{\varphi_+} \rangle = \delta_{ij} \overline{\varphi_+(\beta_j)}$$

and, combining this with (9.2), we also obtain

$$(9.9) \quad \forall i \in \{1, \dots, r\}, \forall j \in \{1, \dots, r + g_-\}, \quad \langle a_j^{\varphi_+}, a_i^{\varphi_+} \rangle = 0, \quad \langle b_j^{\varphi_+}, a_i^{\varphi_+} \rangle = -\delta_{ij} \varphi_+(\beta_j).$$

Let $P \subset \{1, \dots, 2g_-\}$ with $|P| = j$ and let $Q \subset \{1, \dots, 2g_+\}$ with $|Q| = r + j$. It follows from Lemma 8.2 that

$$\langle \mathbb{R}(M, \varphi)(v_P^-), v_Q^+ \rangle = \langle a_1^{\varphi_+} \wedge \cdots \wedge a_r^{\varphi_+} \wedge (v_P^-)^+, v_Q^+ \rangle.$$

According to (9.8), this determinant is zero if the subset $B := \{g_+ + 1, \dots, g_+ + r\}$ is not contained in Q . If $B \subset Q$, then we get

$$\begin{aligned} \langle \mathbb{R}(M, \varphi)(v_P^-), v_Q^+ \rangle &= \varepsilon_B \langle a_1^{\varphi_+} \wedge \cdots \wedge a_r^{\varphi_+} \wedge (v_P^-)^+, v_B^+ \wedge v_{B^c}^+ \rangle \\ &= \varepsilon_B \langle a_1^{\varphi_+} \wedge \cdots \wedge a_r^{\varphi_+}, v_B^+ \rangle \langle (v_P^-)^+, v_{B^c}^+ \rangle \\ &= \varepsilon_B \overline{\varphi_+(\beta_1 \cdots \beta_r)} \langle (v_P^-)^+, v_{B^c}^+ \rangle \end{aligned}$$

where $B^c := Q \setminus B$ and ε_B is the signature of the permutation BB^c (where the elements of B in increasing order are followed by the elements of B^c in increasing order). We also deduce from (9.9) that $\langle (v_P^-)^+, v_{B^c}^+ \rangle = 0$ if B^c has a non-empty intersection with $A := \{1, \dots, r\}$, and it follows that $\langle \mathbb{R}(M, \varphi)(v_P^-), v_Q^+ \rangle = 0$ if $A \cap Q \neq \emptyset$.

Besides, it follows from Lemma 8.1 that $\mathbb{R}(\overline{M}, \varphi)(v_Q^+)$ is trivial if $A \cap Q \neq \emptyset$ or B is not contained in Q . If $A \cap Q = \emptyset$ and $B \subset Q$, we get

$$\langle v_P^-, \mathbb{R}(\overline{M}, \varphi)(v_Q^+) \rangle = \varepsilon_B \langle v_P^-, \mathbb{R}(\overline{M}, \varphi)(v_B^+ \wedge v_{B^c}^+) \rangle = \varepsilon_B \langle v_P^-, (v_{B^c}^+)^- \rangle.$$

We deduce that $\langle R(M, \varphi)(v_{\overline{P}}^-, v_Q^+) \rangle = \overline{\varphi_+(\beta_1 \cdots \beta_r)} \langle v_{\overline{P}}^-, R(\overline{M}, \varphi)(v_Q^+) \rangle$ for any P, Q . Since (9.3) is only required to hold true up to multiplication by a constant in $\pm G$, the theorem is proved in case (ii).

Case (iii). Assume now that $M = C_r^0 \otimes \text{Id}_{g_+}$ where $r = -\delta g$. Note that $\varphi_-(\beta_i) = 1$ for all $i \in \{1, \dots, r\}$, so that (9.7) and (9.5) applied to $\psi := \varphi_-$ give (9.10)

$$\forall i \in \{1, \dots, r + g_+\}, \forall j \in \{1, \dots, r\}, \quad \langle b_i^{\varphi_-}, b_j^{\varphi_-} \rangle = 0, \quad \langle a_i^{\varphi_-}, b_j^{\varphi_-} \rangle = \delta_{ij} \varphi_-(\alpha_i)$$

and, combining this with (9.2), we also obtain

$$(9.11) \quad \forall i \in \{1, \dots, r + g_+\}, j \in \{1, \dots, r\}, \quad \langle b_j^{\varphi_-}, b_i^{\varphi_-} \rangle = 0, \quad \langle b_j^{\varphi_-}, a_i^{\varphi_-} \rangle = -\delta_{ij} \overline{\varphi_-(\alpha_i)}.$$

Let $P \subset \{1, \dots, 2g_-\}$ with $|P| = j$ and let $Q \subset \{1, \dots, 2g_+\}$ with $|Q| = j - r$. By Lemma 8.1, $R(M, \varphi)(v_{\overline{P}}^-)$ is trivial if P does not contain $A := \{1, \dots, r\}$ or P has a non-empty intersection with $B := \{g_- + 1, \dots, g_- + r\}$. If $A \subset P$ and $P \cap B = \emptyset$, we obtain

$$\langle R(M, \varphi)(v_{\overline{P}}^-), v_Q^+ \rangle = \varepsilon_A \langle R(M, \varphi)(v_A^- \wedge v_{A^c}^-), v_Q^+ \rangle = \varepsilon_A \langle (v_{A^c}^-)^+, v_Q^+ \rangle$$

where $A^c := P \setminus A$ and ε_A is the signature of the permutation AA^c .

Besides, Lemma 8.2 gives

$$\langle v_{\overline{P}}^-, R(\overline{M}, \varphi)(v_Q^+) \rangle = \langle v_{\overline{P}}^-, b_1^{\varphi_-} \wedge \cdots \wedge b_r^{\varphi_-} \wedge (v_Q^+)^- \rangle$$

which, according to (9.10), is trivial if P does not contain A . If $A \subset P$, we get

$$\begin{aligned} \langle v_{\overline{P}}^-, R(\overline{M}, \varphi)(v_Q^+) \rangle &= \varepsilon_A \langle v_A^- \wedge v_{A^c}^-, b_1^{\varphi_-} \wedge \cdots \wedge b_r^{\varphi_-} \wedge (v_Q^+)^- \rangle \\ &= \varepsilon_A \langle v_A^-, b_1^{\varphi_-} \wedge \cdots \wedge b_r^{\varphi_-} \rangle \langle v_{A^c}^-, (v_Q^+)^- \rangle \\ &= \varepsilon_A \varphi_-(\alpha_1 \cdots \alpha_r) \langle v_{A^c}^-, (v_Q^+)^- \rangle. \end{aligned}$$

It follows from (9.11) that $\langle v_{A^c}^-, (v_Q^+)^- \rangle = 0$ if A^c has a non-empty intersection with B , so that $\langle v_{\overline{P}}^-, R(\overline{M}, \varphi)(v_Q^+) \rangle = 0$ if $P \cap B \neq \emptyset$. We deduce that $\langle R(M, \varphi)(v_{\overline{P}}^-), v_Q^+ \rangle = \overline{\varphi_-(\alpha_1 \cdots \alpha_r)} \langle v_{\overline{P}}^-, R(\overline{M}, \varphi)(v_Q^+) \rangle$ for any P, Q . This proves the theorem in case (iii). \square

Example 9.3. We consider the situation of §7.3: let $\psi : H_1(F_k) \rightarrow G$ be a group homomorphism and let $M \in \mathcal{C}^\psi(F_k)$ with $k \geq 1$. According to Proposition 7.2, $R(M, \psi)$ is determined by the relative Reidemeister torsion $\tau^\psi(M, \partial_+ M)$ and the Magnus representation $r^\psi(M) : H^\psi \rightarrow H^\psi$, where $H^\psi := H_1^\psi(F_k, \star)$. Specializing Theorem 9.2 to $j := 0$, we obtain the well-known duality theorem

$$(9.12) \quad \tau^\psi(M, \partial_+ M) = \overline{\tau^\psi(M, \partial_- M)} \in \mathbb{F} / \pm G,$$

see [Tur2, Appendix 3]. Next, specializing Theorem 9.2 successively to $j := 1$ and $j := 2$, we obtain the invariance property

$$\forall x, z \in H^\psi, \quad \langle r^\psi(M)(x), r^\psi(M)(z) \rangle = \langle x, z \rangle,$$

which is already observed in [Sak1, Theorem 2.4].

Example 9.4. We consider the situation of §3.3: let G be the infinite cyclic group generated by t and $\mathbb{F} := Q(G)$, let M_K be the exterior of an oriented knot K in an oriented homology 3-sphere and let $\varphi_K : H_1(M_K) \rightarrow G$ be the canonical isomorphism. There is a system of meridian and parallel (α, β) on F_1 and a boundary-parametrization $m : F(1, 0) \rightarrow \partial M_K$ such that

- (i) $m_-(\alpha)$ is the oriented meridian of K and $m_-(\beta)$ is the parallel of K that is null-homologous in M_K ,
- (ii) the matrix of $\langle -, - \rangle : H_- \times H_- \rightarrow \mathbb{F}$ in the corresponding basis $(a, b) := (a_1^{\varphi_K m_-}, b_1^{\varphi_K m_-})$ of $H_- := H_1^{\varphi_K m_-}(F_1, \star)$ is $\begin{pmatrix} 1-t & t \\ -t^{-1} & 0 \end{pmatrix}$.

According to Proposition 3.3, the map $R(M_K, \varphi_K)$ is determined by the Alexander polynomial $\Delta(K)$. By applying Theorem 9.2 successively to $x := a$ and $x := b$, we get

$$(9.13) \quad R(\overline{M}_K, \varphi_K)(1) = \overline{\Delta(K)} b \in H_-.$$

9.3. Second duality. The second duality satisfied by R does not involve the conjugation $f \mapsto \overline{f}$ of the field \mathbb{F} , and is an immediate consequence of the definitions.

Proposition 9.5. *For any $(M, \varphi) \in \text{Cob}_G((g_-, \varphi_-), (g_+, \varphi_+))$ and $j \geq 0$, we have $\forall x \in \Lambda^j H_-$, $\forall y \in \Lambda^{g-j} H_+$, $\omega(R(M, \varphi)(x) \wedge y) = (-1)^{jg} \cdot \omega(x \wedge R(\overline{M}, \varphi)(y))$ where $g := g_+ + g_-$, $H_\pm := H_1^{\varphi_\pm}(F_{g_\pm}, \star)$ and $\omega : \Lambda^{2g_\pm} H_\pm \rightarrow \mathbb{F}$ is an arbitrary integral volume form.*

Despite its simplicity, this proposition turns out to be interesting when it is combined with Theorem 9.2.

Example 9.6. We use the same notation as in Example 9.3. Let (z_1, \dots, z_{2k}) be a basis of H^ψ arising from of a basis of the free $\mathbb{Z}[H_1(F_k)]$ -module $H_1(F_k, \star; \mathbb{Z}[H_1(F_k)])$ and assume that ω is given by $\omega(z_1 \wedge \dots \wedge z_{2k}) = 1$. By applying Proposition 9.5 to $x := z_1 \wedge \dots \wedge z_{2k}$, we get $\tau^\psi(M, \partial_+ M) \cdot \det r^\psi(M) = \tau^\psi(M, \partial_- M)$. Combined with (9.12), this relation gives the symmetry

$$\tau^\psi(M, \partial_+ M) \cdot \det r^\psi(M) = \overline{\tau^\psi(M, \partial_+ M)} \in \mathbb{F} / \pm G$$

which is also observed in [Sak4, Theorem 5.3].

Example 9.7. We use the same notation as in Example 9.4. Let ω be the volume form on H_- defined by $\omega(a \wedge b) = 1$. By applying Proposition 9.5 successively to $x := a$ and $x := b$, we obtain $R(\overline{M_K}, \varphi_K)(1) = \Delta(K)b$. Combined with (9.13), we recover the classical symmetry of the Alexander polynomial:

$$\Delta(K) = \overline{\Delta(K)} \in \mathbb{Z}[G]/\pm G.$$

A. A short review of combinatorial torsions

We recall the definition and basic properties of the torsions of chain complexes. The reader is referred to [Mil2] and [Tur4] for further details and references. In this appendix, \mathbb{F} is a field.

A.1. Definition of the torsion. Given an \mathbb{F} -vector space V of finite dimension $n \geq 0$, an n -tuple $b = (b_1, \dots, b_n)$ of vectors in V and a basis $c = (c_1, \dots, c_n)$ of V , we denote by $[b/c] \in \mathbb{F}$ the determinant of the matrix expressing b in the basis c . Two bases b and c are said to be *equivalent* if $[b/c] = 1$.

Given a short exact sequence of \mathbb{F} -vector spaces $0 \rightarrow V' \rightarrow V \rightarrow V'' \rightarrow 0$ and some bases c' and c'' of V' and V'' respectively, we denote by $c'c''$ the equivalence class of bases of V obtained by juxtaposing (in this order) the image of c' in V and a lift of c'' to V .

By a *finite \mathbb{F} -chain complex of length $m \geq 1$* , we mean a chain complex C in the category of finite-dimensional \mathbb{F} -vector spaces and we assume that C is concentrated in degrees $0, \dots, m$:

$$C = \left(C_m \xrightarrow{\partial_m} C_{m-1} \longrightarrow \dots \xrightarrow{\partial_1} C_0 \right).$$

A *basis* of C is a family $c = (c_m, \dots, c_0)$ where c_i is a basis of C_i for all $i \in \{0, \dots, m\}$. A *homological basis* of C is a family $h = (h_m, \dots, h_0)$ where h_i is a basis of the i -th homology group $H_i(C)$ for all $i \in \{0, \dots, m\}$. If we have chosen a basis b_j of the space of j -dimensional boundaries $B_j(C) := \text{Im } \partial_{j+1}$ for all $j \in \{0, \dots, m-1\}$, then a homological basis h of C induces an equivalence class of bases of C_i for any i : specifically, we consider the basis $(b_i h_i) b_{i-1}$ of C_i obtained by juxtaposition in the following short exact sequences where we denote $Z_i(C) := \text{Ker } \partial_i$:

$$(A.1) \quad 0 \longrightarrow B_i(C) \longrightarrow Z_i(C) \longrightarrow H_i(C) \longrightarrow 0$$

$$(A.2) \quad \text{and } 0 \longrightarrow Z_i(C) \longrightarrow C_i \xrightarrow{\partial_i} B_{i-1}(C) \longrightarrow 0.$$

Definition A.1. The *torsion* of a finite \mathbb{F} -chain complex C of length m , equipped with a basis c and a homological basis h , is the scalar

$$\tau(C; c, h) := \prod_{i=0}^m [(b_i h_i) b_{i-1} / c_i]^{(-1)^{i+1}} \in \mathbb{F} \setminus \{0\}.$$

It is easily checked that this definition does not depend on the choice of b_0, \dots, b_m and, when C is acyclic, we set $\tau(C; c) := \tau(C; c, \emptyset)$.

The following lemma, which is well known, is a way of viewing the torsion as a function in homology.

Lemma A.2. *Let C be a finite \mathbb{F} -chain complex of length $m \geq 1$, let $k \in \{0, \dots, m\}$ and set $\beta := \dim H_k(C)$. Assume given a basis $c = (c_m, \dots, c_0)$ of C and a basis h_i of $H_i(C)$ for every $i \neq k$. Then there is a unique linear map $\ell : \Lambda^\beta H_k(C) \rightarrow \mathbb{F}$ such that*

$$\ell(v_1 \wedge \dots \wedge v_\beta) = \begin{cases} \tau(C; c, (h_m, \dots, h_{k+1}, v, h_{k-1}, \dots, h_0)) & \text{if } k \text{ is odd,} \\ \tau(C; c, (h_m, \dots, h_{k+1}, v, h_{k-1}, \dots, h_0))^{-1} & \text{if } k \text{ is even,} \end{cases}$$

for any basis $v = (v_1, \dots, v_\beta)$ of $H_k(C)$.

Proof. The unicity of ℓ is obvious and, clearly, we can assume that k is odd. Let $s : H_k(C) \rightarrow Z_k(C)$ and $t : B_{k-1}(C) \rightarrow C_k$ be \mathbb{F} -linear sections of (A.1) and (A.2), respectively. For any β -tuple $v = (v_1, \dots, v_\beta)$ of elements of $H_k(C)$, we set

$$\ell(v) := [b_k s(v) t(b_{k-1}) / c_k] \cdot \prod_{i \neq k} [(b_i h_i) b_{i-1} / c_i]^{(-1)^{i+1}} \in \mathbb{F}$$

where $b_k s(v) t(b_{k-1})$ denotes the family of vectors of C_k obtained by juxtaposing (in this order) b_k , $s(v)$ and $t(b_{k-1})$. The resulting map $\ell : H_k(C)^\beta \rightarrow \mathbb{F}$ is multilinear and alternate, hence it induces a map $\ell : \Lambda^\beta H_k(C) \rightarrow \mathbb{F}$ with the desired property. \square

A.2. Multiplicativity of the torsion. Consider a short exact sequence of finite \mathbb{F} -chain complexes of length $m \geq 1$:

$$(A.3) \quad 0 \longrightarrow C' \longrightarrow C \longrightarrow C'' \longrightarrow 0.$$

Let us assume that C', C, C'' are based by c', c, c'' respectively, and homologically based by h', h, h'' respectively. We further assume that the bases c', c, c'' are *compatible* in the sense that c_i is equivalent to $c'_i c''_i$ for every $i \in \{0, \dots, m\}$. The short exact sequence (A.3) induces a long exact sequence in homology:

$$\mathcal{H} := (H_m(C') \rightarrow H_m(C) \rightarrow H_m(C'') \rightarrow \cdots \rightarrow H_0(C') \rightarrow H_0(C) \rightarrow H_0(C'')).$$

We regard \mathcal{H} as an acyclic finite \mathbb{F} -chain complex based by

$$(h', h, h'') := (h'_m, h_m, h''_m, \dots, h'_0, h_0, h''_0).$$

The following formula is classical in the theory of combinatorial torsions: see [Mil2, Theorem 3.2] or [Tur2, Lemma 3.4.2].

Theorem A.3. *With the above notation, we have*

$$(A.4) \quad \tau(C; c, h) = \varepsilon \cdot \tau(C'; c', h') \cdot \tau(C''; c'', h'') \cdot \tau(\mathcal{H}; (h', h, h''))$$

where ε is a sign depending only on the dimensions of the \mathbb{F} -vector spaces C'_i, C_i, C''_i and $H_i(C'), H_i(C), H_i(C'')$ for all $i \in \{0, \dots, m\}$.

Example A.4. Assume that $C = C' \oplus C''$ and that the chain maps $C' \rightarrow C$ and $C \rightarrow C''$ in (A.3) are the natural inclusion and projection, respectively. For all $i \in \{0, \dots, m\}$, let c_i be the basis of $C_i = C'_i \oplus C''_i$ obtained by juxtaposing (in this order) some bases c'_i and c''_i of C'_i and C''_i , respectively; similarly, let h_i be the basis of $H_i(C) = H_i(C') \oplus H_i(C'')$ obtained by juxtaposing some bases h'_i and h''_i of $H_i(C')$ and $H_i(C'')$, respectively. We set $c := (c_m, \dots, c_0)$ and $h := (h_m, \dots, h_0)$. Then $\tau(C; c, h) = \varepsilon \cdot \tau(C'; c', h') \cdot \tau(C''; c'', h'')$.

References

- [Arc] J. ARCHIBALD, *The multivariable Alexander polynomial on tangles*. ProQuest LLC, Ann Arbor, MI, 2010. Thesis (Ph.D.)—University of Toronto (Canada). [MR 2941804](#)
- [Aty] M. ATIYAH, Topological quantum field theories. *Inst. Hautes Études Sci. Publ. Math.* **68** (1988), 175–186 (1989). [Zbl 0692.53053](#) [MR 1001453](#)
- [BCF] S. BIGELOW, A. CATTABRIGA and V. FLORENS, Alexander representation of tangles. *Acta Math. Vietnam.* **40** (2015), 339–352. [MR 3366173](#)
- [Bla] R.C. BLANCHFIELD, Intersection theory of manifolds with operators with applications to knot theory. *Ann. of Math. (2)* **65** (1957), 340–356. [Zbl 0080.16601](#) [MR 0085512](#)
- [Cha] T.A. CHAPMAN, Topological invariance of Whitehead torsion. *Amer. J. Math.* **96** (1974), 488–497. [Zbl 0358.57004](#) [MR 0391109](#)
- [CT1] D. CIMASONI and V. TURAEV, A Lagrangian representation of tangles. *Topology* **44** (2005), 747–767. [Zbl 1071.57006](#) [MR 2136533](#)
- [CT2] D. CIMASONI and V. TURAEV, A Lagrangian representation of tangles. II. *Fund. Math.* **190** (2006), 11–27. [Zbl 1106.57004](#) [MR 2232854](#)

- [Coh] M. M. COHEN, *A course in simple-homotopy theory*. Springer-Verlag, New York–Berlin, 1973. Graduate Texts in Mathematics, Vol. 10. [Zbl 0261.57009](#) [MR 0362320](#)
- [CY] L. CRANE and D. YETTER, On algebraic structures implicit in topological quantum field theories. *J. Knot Theory Ramifications* **8** (1999), 125–163. [Zbl 0935.57025](#) [MR 1687553](#)
- [FJR] S. FRIEDL, A. JUHÁSZ and J. RASMUSSEN, The decategorification of sutured Floer homology. *J. Topol.* **4**, 2 (2011), 431–478. [Zbl 1270.57041](#) [MR 2805998](#)
- [FN1] C. FROHMAN and A. NICAS, The Alexander polynomial via topological quantum field theory. In *Differential geometry, global analysis, and topology (Halifax, NS, 1990)*, vol. 12 of *CMS Conf. Proc.* Amer. Math. Soc., Providence, RI, 1991, pp. 27–40. [Zbl 0757.57013](#) [MR 1158467](#)
- [FN2] —, An intersection homology invariant for knots in a rational homology 3-sphere. *Topology* **33** (1994), 123–158. [Zbl 0822.57008](#) [MR 1259519](#)
- [Hab] K. HABIRO, Bottom tangles and universal invariants. *Algebr. Geom. Topol.* **6** (2006), 1113–1214 (electronic). [Zbl 1130.57014](#) [MR 2253443](#)
- [Ker1] T. KERLER, Homology TQFT’s and the Alexander-Reidemeister invariant of 3-manifolds via Hopf algebras and skein theory. *Canad. J. Math.* **55** (2003), 766–821. [Zbl 1083.57038](#) [MR 1994073](#)
- [Ker2] —, Towards an algebraic characterization of 3-dimensional cobordisms. In *Diagrammatic morphisms and applications (San Francisco, CA, 2000)*, vol. 318 of *Contemp. Math.* Amer. Math. Soc., Providence, RI, 2003, pp. 141–173. [Zbl 1168.57313](#) [MR 1973515](#)
- [KLW] P. KIRK, C. LIVINGSTON and Z. WANG, The Gassner representation for string links. *Commun. Contemp. Math.* **3** (2001), 87–136. [Zbl 0989.57005](#) [MR 1820015](#)
- [LDi] J.-Y. LE DIMET, Enlacements d’intervalles et représentation de Gassner. *Comment. Math. Helv.* **67** (1992), 306–315. [Zbl 0759.57010](#) [MR 1161287](#)
- [Les] C. LESCOP, A sum formula for the Casson-Walker invariant. *Invent. Math.* **133** (1998), 613–681. [Zbl 0917.57008](#) [MR 1645066](#)
- [MM] G. MASSUYEAU and J.-B. MEILHAN, Equivalence relations for homology cylinders and the core of the Casson invariant. *Trans. Amer. Math. Soc.* **365** (2013), 5431–5502. [Zbl 1283.57020](#) [MR 3074379](#)
- [Mat] S. MATVEEV, *Algorithmic topology and classification of 3-manifolds*, vol. 9 of *Algorithms and Computation in Mathematics*. Springer-Verlag, Berlin, 2003. [Zbl 1128.57001](#) [MR 2341532](#)
- [Mil1] J. MILNOR, A duality theorem for Reidemeister torsion. *Ann. of Math. (2)* **76** (1962), 137–147. [Zbl 0108.36502](#) [MR 0141115](#)
- [Mil2] —, Whitehead torsion. *Bull. Amer. Math. Soc.* **72** (1966), 358–426. [Zbl 0147.23104](#) [MR 0196736](#)
- [Pap] C. D. PAPAKYRIAKOPOULOS, Planar regular coverings of orientable closed surfaces. In *Knots, groups, and 3-manifolds (Papers dedicated to the memory of R. H. Fox)*. Princeton Univ. Press, Princeton, N.J., 1975, pp. 261–292. *Ann. of Math. Studies*, No. 84. [Zbl 0325.55002](#) [MR 0388396](#)

- [Per] B. PERRON, A homotopic intersection theory on surfaces: applications to mapping class group and braids. *Enseign. Math. (2)* **52** (2006), 159–186. [Zbl 1161.57009](#) [MR 2255532](#)
- [Sak1] T. SAKASAI, The symplecticity of the Magnus representation for homology cobordisms of surfaces. *Bull. Austral. Math. Soc.* **76** (2007), 421–431. [Zbl 1155.57002](#)
- [Sak2] —, The Magnus representation and higher-order Alexander invariants for homology cobordisms of surfaces. *Algebr. Geom. Topol.* **8** (2008), 803–848. [Zbl 1156.57001](#) [MR 2443097](#)
- [Sak3] —, A survey of Magnus representations for mapping class groups and homology cobordisms of surfaces. In *Handbook of Teichmüller theory. Volume III*, vol. 17 of *IRMA Lect. Math. Theor. Phys.* Eur. Math. Soc., Zürich, 2012, pp. 531–594. [Zbl 1272.57001](#) [MR 2952771](#)
- [Sak4] —, The Magnus representation and homology cobordism groups of homology cylinders. [arXiv:1108.6111](#), preprint 2011.
- [Tur1] V. TURAEV, The Alexander polynomial of a three-dimensional manifold. *Mat. Sb. (N.S.)* **97(139)**, 3(7) (1975), 341–359, 463. [Zbl 0315.57001](#)
- [Tur2] —, Reidemeister torsion in knot theory. *Uspekhi Mat. Nauk* **41** (1986), 97–147, 240.
- [Tur3] —, Euler structures, nonsingular vector fields, and Reidemeister-type torsions. *Izv. Akad. Nauk SSSR Ser. Mat.* **53** (1989), 607–643, 672. [Zbl 0692.57015](#) [MR 1013714](#)
- [Tur4] —, *Introduction to combinatorial torsions*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2001. Notes taken by Felix Schlenk. [Zbl 0970.57001](#) [MR 1809561](#)
- [Tur5] —, *Torsions of 3-dimensional manifolds*, vol. 208 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2002. [Zbl 1012.57002](#) [MR 1958479](#)
- [Tur6] —, *Homotopy quantum field theory*, vol. 10 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2010. Appendix 5 by M. MÜGER and Appendices 6 and 7 by A. VIRELIZIER. [Zbl 1243.81016](#) [MR 2674592](#)
- [Tur7] —, Intersections of loops in two-dimensional manifolds. *Mat. Sb.* **106(148)** (1978), 566–588. [MR 0507817](#)

(Reçu le 22 avril 2014)

Vincent FLORENS, LMA, Université de Pau & CNRS, Avenue de l'Université, 64000 Pau, France

e-mail: vincent.florens@univ-pau.fr

Gwénaél MASSUYEAU, IRMA, Université de Strasbourg & CNRS, 7 rue René Descartes, 67084 Strasbourg, France

e-mail: massuyeau@math.unistra.fr