

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 61 (2015)  
**Heft:** 3-4

**Artikel:** The exceptional set and the Green-Griffiths locus do not always coincide  
**Autor:** Diverio, Simone / Rousseau, Erwan  
**DOI:** <https://doi.org/10.5169/seals-630600>

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 13.01.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# The exceptional set and the Green–Griffiths locus do not always coincide

Simone DIVERIO and Erwan ROUSSEAU

**Abstract.** We give a very simple criterion for the Green–Griffiths locus of a projective manifold to be the whole manifold. Then, we use it to show that the Green–Griffiths locus of any projective manifold uniformized by a bounded symmetric domain of rank greater than one is the whole manifold. In particular, this clarifies an old undetailed example given by M. Green to S. Lang.

**Mathematics Subject Classification (2010).** Primary: 32Q45, 32M15; Secondary: 37F75.

**Keywords.** Bounded symmetric domain, Green–Griffiths locus, jet differentials, Kobayashi hyperbolicity, Green–Griffiths–Lang conjecture.

## 1. Introduction

Let  $X$  be a compact complex manifold. Then, by Brody's theorem,  $X$  is *Kobayashi hyperbolic* if and only if there is no non-constant holomorphic map  $f: \mathbb{C} \rightarrow X$  (such a map will be called *entire curve*). When  $X$  is a compact Riemann surface, the uniformization theorem together with Liouville's theorem imply that  $X$  is Kobayashi hyperbolic if and only if its geometric genus is at least two. Thus, the majority of compact Riemann surfaces are hyperbolic.

Being of genus greater than one can be also characterized in algebraic terms. To this effect, let us consider the canonical line bundle  $K_X$  of a compact hyperbolic Riemann surface  $X$ , as well as its tensor powers  $K_X^{\otimes m}$ . As usual, the canonical bundle is just the top exterior power of the holomorphic cotangent bundle of the manifold (thus, in this case, it is simply the holomorphic cotangent bundle itself). Denote by  $H^0(X, K_X^{\otimes m})$  the space of global holomorphic sections of the holomorphic line bundle  $K_X^{\otimes m}$ : these sections are called global pluricanonical forms. Then, by the Riemann–Roch theorem, the dimension of the space  $H^0(X, K_X^{\otimes m})$  grows linearly with  $m$ . On the other hand, if the genus of

$X$  is one (resp. zero), then these dimensions are constantly equal to one (resp. to zero).

A compact complex projective manifold is said to be of *general type* if there exists a positive constant  $C$  such that

$$C^{-1} m^{\dim X} \leq \dim H^0(X, K_X^{\otimes m}) \leq C m^{\dim X},$$

for all sufficiently large integers  $m > 0$ . For instance, a projective manifold with ample (i.e., positive) canonical bundle is easily seen to be of general type. In particular, a compact Riemann surface is hyperbolic if and only if it is of general type.

Since Kobayashi hyperbolicity obviously is a hereditary property when passing to subvarieties while being of general type is not, one cannot hope for a straightforward generalization in higher dimension of the above characterization of hyperbolicity in terms of pluricanonical sections. Still, one can impose such an hereditary property and conjecture:

**Conjecture 1.1** (Lang [Lan86]). *Let  $X$  be a compact complex projective manifold. Then,  $X$  is Kobayashi hyperbolic if and only if  $X$  together with all its subvarieties is of general type.*

This conjecture is almost completely open: a remarkable exception is the case of surfaces, where the necessity condition can be proved to be true by using the birational classification together with the non hyperbolicity of  $K3$  surfaces, while the sufficiency condition is shown in [McQ98] (see also [Bog77, Des79]) under the assumption of positivity of the second Segre number. The sufficiency part can be rephrased as follows.

**Conjecture 1.2** (Green–Griffiths, Lang [GG80, Lan86]). *Let  $X$  be a compact complex projective manifold of general type. Then, there should exist a proper subvariety  $Y \subsetneq X$  such that all entire curves of  $X$  are indeed contained in  $Y$ .*

A nowadays classical strategy to attack this kind of questions, i.e. algebraic degeneracy of entire curves, is to use jet differentials. Introduced (in their modern form) by Green and Griffiths [GG80], and subsequently refined by Demainly [Dem97], jet differentials are, roughly speaking, a particular kind of algebraic differential equations on a projective manifold which all entire curves must satisfy. In order to give an heuristic idea of this strategy, suppose  $X$  is a projective manifold and that we dispose of a certain family  $\{P_i\}_{i \in I}$  of such algebraic differential equations, say of order  $k$ . Then, if  $f: \mathbb{C} \rightarrow X$  is any entire curve, the general theory (see below for more details) tells us that for all  $i \in I$ ,

$$P_i(f, f', \dots, f^{(k)}) \equiv 0.$$

The hope is then that if we have enough *algebraically independent* differential equations on  $X$ , then we should be able to successively eliminate the variables  $f^{(k)}, f^{(k-1)}, \dots$ , and so forth, so that we end up with a non trivial algebraic equation for  $f$  only, say  $Q(f) \equiv 0$ . This would give the desired algebraic degeneracy for the entire curve. Roughly speaking, the Green–Griffiths locus of  $X$  (see right after Theorem 1.3 for the precise definition) is defined to be the intersection on  $X$  of the zero loci of all such algebraic equations  $Q$ . This is a closed algebraic subvariety of  $X$  (of course, a priori possibly equal to  $X$  itself) which thus contains all the non constant holomorphic images of the complex plane inside  $X$ , i.e., the so-called exceptional locus.

A first substantial obstacle is that it is already a challenge to find even one single such jet differential: for instance, it is known [Div08] (see also [Bro11]) that for  $X \subset \mathbb{P}^{n+1}$  a smooth projective hypersurface there is no non zero such algebraic differential operator of order less than  $n$ .

Anyway, given a complex manifold  $X$ , it turns out that for all positive integers  $k, m$ , there exists a holomorphic vector bundle  $E_{k,m}^{GG} T^* X \rightarrow X$  whose global holomorphic sections are precisely the algebraic differential equations (of order  $k$  and weighted degree  $m$ ) above. More generally, one can look at a holomorphic subbundle  $V \subset T_X$  (following [Dem97], we shall call the pair  $(X, V)$  a *directed manifold*) and consider only entire curves in  $X$  which are tangent to  $V$  (the typical situation is for instance when  $X$  is fibered over another manifold and one is interested in entire curves lying on the fibers). Then, there exists a corresponding theory of jet differentials acting on germs of holomorphic curves tangent to  $V$ , and the analogous vector bundle is then denoted by  $E_{k,m}^{GG} V^* \rightarrow X$ . For definitions and basic properties of these bundles we refer to [GG80] and [Dem97], as well as to Section 2 below.

The following theorem is the fundamental vanishing result and formalizes the above discussion.

**Theorem 1.3** (Green–Griffiths [GG80], Siu–Yeung [SY97], and Demailly [Dem97]). *Let  $(X, V)$  be a directed compact projective manifold and  $A \rightarrow X$  an ample line bundle. Then, for any entire curve  $f: \mathbb{C} \rightarrow X$  tangent to  $V$  and any  $P \in H^0(X, E_{k,m}^{GG} V^* \otimes A^{-1})$ , one has  $P(f, f', \dots, f^{(k)}) \equiv 0$ .*

Now, fix an ample line bundle  $A \rightarrow X$ . Define  $GG_A(X, V)$  to be the set of points  $x \in X$  such that for all integers  $k > 0$  there exists a  $k$ -jet of holomorphic curve  $\varphi_k: (\mathbb{C}, 0) \rightarrow (X, x)$  tangent to  $V$  with the property that for all integers  $m > 0$ , every global jet differential of order  $k$  and weighted degree  $m$  with values in  $A^{-1}$  vanishes whenever evaluated on (the  $k$ -jet defined by)  $\varphi_k$  at 0. Thus, if

$f: \mathbb{C} \rightarrow X$  is an entire curve tangent to  $V$ , Theorem 1.3 tells us immediately that  $f(\mathbb{C}) \subset GG_A(X, V)$ . We shall see in the next section that the locus  $GG_A(X, V)$  is indeed independent of the particular ample line bundle chosen.

**Definition 1.4.** Let  $X$  be a projective manifold and  $V \subset T_X$  a holomorphic subbundle. The *exceptional set*  $Exc(X, V)$  of  $(X, V)$  is defined to be the Zariski closure of the union of all the images of entire curves traced in  $X$  and tangent to  $V$ . The *Green–Griffiths locus*  $GG(X, V)$  of  $(X, V)$  is defined as

$$GG(X, V) = GG_A(X, V),$$

for some (and hence any) ample line bundle  $A \rightarrow X$ .

In the *absolute case*  $V = T_X$ , we shall simply call  $Exc(X, T_X) = Exc(X)$  and  $GG(X, T_X) = GG(X)$ .

By Theorem 1.3, one always has the inclusion:

$$Exc(X, V) \subset GG(X, V).$$

It is conjectured in [GG80] that if  $X$  is a  $n$  dimensional projective manifold of general type and  $V = T_X$ , then there should exist a large integer  $k = k(X)$  such that the growth rate of

$$m \mapsto \dim H^0(X, E_{k,m}^{GG} T_X^*)$$

would have to be maximal (that is asymptotic with  $m^{(k+1)n-1}$ ). This conjecture was proven in [GG80] for projective surfaces and therein supported in all dimensions by an Euler characteristic computation. It was then established in the special case of smooth projective hypersurfaces of general type in [Mer10] and was finally proven in full generality only very recently by J.-P. Demailly in [Dem11] by means of his holomorphic Morse inequalities combined with a delicate “probabilistic” curvature estimate. In particular, if  $X$  is a projective manifold of general type and  $A \rightarrow X$  an ample line bundle, then there always exist large integers  $k, m > 0$  such that  $H^0(X, E_{k,m}^{GG} T_X^* \otimes A^{-1}) \neq \{0\}$  and indeed this space is very big. Thus, this settles the above-mentioned problem of the existence of global differential operators satisfied by all entire curves in the case of projective manifold of general type. This abundance of high order jet differentials makes legitimate to ask if much more is true, namely that if  $X$  is a projective manifold of general type, then the Green–Griffiths locus is always a proper subvariety of  $X$ , i.e.

$$GG(X) \subsetneq X.$$

An affirmative answer to this would lead directly to the solution of Conjecture 1.2. One could even speculate more and ask whether  $Exc(X) = GG(X)$ , that is whether  $GG(X)$  provides an algebraic explicit description of the exceptional set. About this, let us quote S. Lang from his paper [Lan86], page 200:

*In particular, the exceptional set [...] is contained in the Green–Griffiths set [...]. I asked Green–Griffiths whether they might be equal. Green told me that the two sets are not equal in general. Certain Hilbert modular surfaces constructed by Shavel [Sha78], compact quotients of the product of the upper half-plane with itself, provide a counterexample which is hyperbolic, but such that the Green–Griffiths set is the whole variety. Thus the jet construction appears insufficient so far to characterize the exceptional set [...] completely algebraically.*

In [Lan86] there are no more details about that, nor in the subsequent literature as far as we know: we have somehow the impression that this paragraph has passed by unnoticed. Here, we take the opportunity to give a detailed account of the above key fact by proving indeed a much more general result, as well as several consequences.

Let  $\mathcal{W}$  be a coherent analytic sheaf on a complex manifold  $X$  and  $m > 0$  be an integer. We shall denote by  $\mathcal{W}^{[m]}$  the double dual of the  $m$ -th tensor power  $\mathcal{W}^{\otimes m}$  of  $\mathcal{W}$ . The starting point is the following criterion.

**Theorem 1.5.** *Let  $(X, V)$  be a complex projective directed manifold. Suppose that there exists a saturated coherent analytic subsheaf  $\mathcal{W} \subset \mathcal{O}_X(V)$  such that for some ample line bundle  $A \rightarrow X$  one has*

$$H^0(X, (\mathcal{W}^*)^{[m]} \otimes A^{-1}) = 0, \quad \forall m \geq 1.$$

*Then,  $GG(X, V) = X$ .*

Observe that we do not require any Frobenius integrability property for  $\mathcal{W}$ . Observe moreover, that by a very recent result of [CPI3], in the case where  $\mathcal{W} = T_X$ , the condition in the statement is satisfied if and only if  $X$  is not of general type. Thus:

**Corollary 1.6.** *If  $X$  is a projective manifold which is not of general type, then  $GG(X) = X$ .*

We now pass to direct and less direct consequences of the above criterion. First of all, we have the following result. It follows straightforwardly from Theorem 1.5 and a technical elementary lemma about growth rate of sections of tensor powers of pull-back of vector bundles (see Section 4 for more details and in fact a slightly more general result concerning finite surjective morphisms).

**Corollary 1.7.** *Let  $X \simeq Y \times Z$  be a complex projective manifold isomorphic to a non trivial product. Then,  $GG(X) = X$ .*

This already shows that in order to produce a counterexample in the spirit of Green–Lang it is not necessary to look so far away at Hilbert modular surfaces, but it is indeed sufficient to consider the product of two hyperbolic compact Riemann surfaces. Here, it is worthwhile mentioning also that a slight variation of the corollary above permits to show that certain surfaces constructed in [BCGP12] provide examples of simply connected surfaces of general type for which the Green–Griffiths locus is the entire surface but the exceptional locus is not known. Therefore, there exist surfaces whose hyperbolicity properties are unknown, and for which it is not possible to use the jet differentials techniques alone in order to deduce such properties (see Section 4 for more details about these examples).

Next, let  $\Omega \subset \mathbb{C}^n$  be a bounded symmetric domain,  $\Gamma \subset \text{Aut}(\Omega)$  a cocompact torsion free lattice and  $X = \Omega/\Gamma$ . The Bergman metric of  $\Omega$  is  $\text{Aut}(\Omega)$ -invariant and descends to a metric  $\omega$  on every smooth quotient  $X$ . Moreover, the cohomology class of  $\omega$  is  $-2\pi c_1(X)$ . Thus, the canonical bundle  $K_X = \det T_X^*$  is positive and in particular  $X$  is projective and of general type. Moreover, whenever  $X$  is a complex space whose universal cover is a bounded domain in  $\mathbb{C}^n$ , then every entire curve must be constant by Liouville’s theorem. Hence, the exceptional locus  $\text{Exc}(X)$  is empty. If  $\Omega$  has rank one or, equivalently, if it is isomorphic to the unit ball  $\Omega \simeq \mathbb{B}^n$ , then it is well known that  $X$  has ample cotangent bundle. In particular, since jet differentials of order one are simply symmetric holomorphic forms, already by looking at order one jet differentials one finds that  $GG(X) = \emptyset$ . The situation changes drastically as soon as the rank of  $\Omega$  becomes bigger than one.

**Theorem 1.8.** *Let  $X$  be a projective manifold uniformized by a bounded symmetric domain  $\Omega$ . Then, either  $\Omega \simeq \mathbb{B}^n$  and thus  $GG(X) = \emptyset$ , or  $GG(X) = X$ .*

This result should be compared to the “all or nothing” principle on the Lang locus of rational points in Shimura varieties established in [UY10] (see also Section 6.1 for more details about this correspondence). One should also relate it to the following fact brought to our attention several times in the last years by Michael McQuillan (cf. also the recent preprint [McQ14]): rational curves on bi–disc quotients which are not the product of curves are Zariski dense in mixed characteristic. This implies, from the mathematical logic point of view, that there cannot exist a proof of the Green–Griffiths conjecture in the first order theory of algebraically closed fields of characteristic zero.

Theorem 1.8 provides a very large class of projective manifolds of general type for which the worst possible situation occurs, as far as a description of the exceptional locus in terms of the Green–Griffiths locus is concerned:

$$\emptyset = \text{Exc}(X) \subsetneq GG(X) = X.$$

This includes of course also the particular Hilbert modular surfaces mentioned above.

Among these locally symmetric manifolds, it is worth distinguishing the two subclasses of locally reducible and of locally irreducible ones. The proof of Theorem 1.8 will be quite different in the two cases. Indeed, in the locally reducible case it is immediate to identify (after a finite unramified covering) a foliation – and thus a subsheaf of the tangent sheaf – to work with in order to apply directly Theorem 1.5. On the other hand, in the locally irreducible case, there is no natural foliation at our disposal. To overcome this difficulty, one possibility is to use the arithmeticity of the lattice (since the rank is at least 2) and the theory of Shimura varieties (see the beginning of Section 6 for details). Alternatively, it is possible to avoid the use of the arithmeticity of the lattice using the theory of characteristic bundles [Mok89]. This shall provide a holomorphic fiber bundle over  $X$ , endowed with a holomorphic foliation by curves given by liftings of minimal disks (we are grateful to N. Mok for suggesting to us several things about this approach). This foliation will turn out to be of negative Kodaira dimension and this will be sufficient to obtain the desired conclusion (see Section 6 for the details).

## 2. Proof of Theorem 1.5

Let us start by recalling some basic facts from [Dem97] about jet differentials in the general framework of directed manifolds (for more details see the cited references).

So, let  $(X, V)$  be a directed manifold, that is a complex manifold  $X$  together with a holomorphic subbundle  $V \subset T_X$ , non necessarily integrable, of the holomorphic tangent bundle  $T_X$  of  $X$ . We call  $J_k V$  the bundle of  $k$ -jets of holomorphic curves  $\varphi: (\mathbb{C}, 0) \rightarrow X$  which are tangent to  $V$ , that is  $\varphi'(t) \in V_{\varphi(t)}$  for all  $t$ , together with the projection  $\varphi \mapsto \varphi(0)$  onto  $X$ . It is a holomorphic fiber bundle, which is naturally a subbundle of the bundle  $J_k T_X$  of  $k$ -jets of holomorphic curves with values in  $X$  with no further restrictions. Moreover, there is a canonically defined fiber-wise  $\mathbb{C}^*$ -action on  $J_k V$  given by the reparametrization of a  $k$ -jet tangent to  $V$  by the homotheties corresponding to elements of  $\mathbb{C}^*$ . Of course, this action is compatible with restriction to smaller subbundles.

Let  $E_{k,m}^{GG} V^* \rightarrow X$  be the holomorphic vector bundle over  $X$  whose fibers are complex valued polynomials on the fibers of  $J_k V$  of weighted degree  $m$  with respect to the above-defined  $\mathbb{C}^*$ -action. If  $W \subset V \subset T_X$  are holomorphic subbundles, the inclusions

$$J_k W \subset J_k V \subset J_k T_X$$

induce surjective arrows

$$E_{k,m}^{GG} T_X^* \rightarrow E_{k,m}^{GG} V^* \rightarrow E_{k,m}^{GG} W^*,$$

so that  $E_{k,m}^{GG} W^*$  can be regarded as a quotient of  $E_{k,m}^{GG} V^*$ , the quotient map being given by evaluating jet differentials tangent to  $V$  only on  $k$ -jets tangent to  $W$ .

**Remark 2.1.** In particular, we see that if every global section of  $E_{k,m}^{GG} W^*$  vanishes at some point  $j_k \in J_k W \subset J_k V$ , then so does every global section of  $E_{k,m}^{GG} V^*$ .

Now, considering the highest monomials (with respect to the reverse lexicographic order) gives a natural filtration on weighted homogeneous polynomials. Such filtration defines an intrinsic filtration on the bundles  $E_{k,m}^{GG} V^*$ , whose graded series are given by

$$\text{Gr}^\bullet E_{k,m}^{GG} V^* = \bigoplus_{\ell_1 + 2\ell_2 + \dots + k\ell_k = m} S^{\ell_1} V^* \otimes \dots \otimes S^{\ell_k} V^*.$$

Therefore, it follows that if for all  $k$ -tuples of non negative integers  $(\ell_1, \dots, \ell_k)$  such that  $\ell_1 + 2\ell_2 + \dots + k\ell_k = m$ , we have

$$H^0(X, S^{\ell_1} V^* \otimes \dots \otimes S^{\ell_k} V^*) = 0,$$

then

$$H^0(X, E_{k,m}^{GG} V^*) = 0,$$

as well.

Before entering into the proof of Theorem 1.5, let us show as promised that the locus  $GG_A(X, V)$  is independent of the ample line bundle  $A \rightarrow X$ .

**Lemma 2.2.** *The set  $GG_A(X, V)$  does not depend on the ample line bundle  $A \rightarrow X$ .*

*Proof.* Let  $A, B \rightarrow X$  two ample line bundles and  $\ell > 0$  be a positive integer such that  $B^{\otimes \ell} \otimes A^{-1}$  is globally generated. We shall show that

$$GG_A(X, V) \subset GG_B(X, V),$$

and the equality will follow by interchanging the roles of  $A$  and  $B$ .

Let  $x \notin GG_B(X, V)$ . Then, there exist integers  $k, m > 0$ , a global section  $P \in H^0(X, E_{k,m}^{GG} V^* \otimes B^{-1})$  and a  $k$ -jet of holomorphic curve  $\varphi_k: (\mathbb{C}, 0) \rightarrow (X, x)$

tangent to  $V$  such that  $P(\varphi_k) \neq 0$ . Let  $\sigma \in H^0(X, B^{\otimes \ell} \otimes A^{-1})$  be a global section such that  $\sigma(x) \neq 0$ . Then,

$$P^\ell \otimes \sigma \in H^0(X, E_{k,\ell m}^{GG} V^* \otimes A^{-1})$$

and

$$(P^\ell \otimes \sigma)(\varphi_k) = P^\ell(\varphi_k) \cdot \sigma(x) \neq 0,$$

so that  $x \notin GG_A(X, V)$ .  $\square$

*Proof of Theorem 1.5.* Let us first suppose that  $\mathcal{W} = \mathcal{O}_X(W)$  is the sheaf of germs of holomorphic sections of a holomorphic subbundle  $W \subset V$ . Let  $A \rightarrow X$  be any ample line bundle and  $k, m > 0$  be two integers. For all  $k$ -tuples of non-negative integers  $(\ell_1, \dots, \ell_k)$  such that  $\ell_1 + 2\ell_2 + \dots + k\ell_k = m$  we have

$$S^{\ell_1} W^* \otimes \dots \otimes S^{\ell_k} W^* \otimes A^{-1} \subset (W^*)^{\otimes |\ell|} \otimes A^{-1},$$

where  $|\ell| = \sum \ell_j$ . Thus,

$$H^0(X, S^{\ell_1} W^* \otimes \dots \otimes S^{\ell_k} W^* \otimes A^{-1}) = 0$$

and, since taking the tensor product of  $E_{k,m} W^*$  with  $A^{-1}$  affects the corresponding graded bundle just tensoring by  $A^{-1}$ , we have

$$H^0(X, E_{k,m}^{GG} W^* \otimes A^{-1}) = 0.$$

Moreover, we have of course surjective morphisms

$$E_{k,m}^{GG} V^* \otimes A^{-1} \rightarrow E_{k,m}^{GG} W^* \otimes A^{-1}.$$

Therefore, by Remark 2.1, in such a situation every global jet differential tangent to  $V$  of order  $k$  and weighted degree  $m$  with values in  $A^{-1}$  must vanish when evaluated on  $k$ -jets which are tangent to  $W$ . Since at each point  $x \in X$  we have such jets, it follows that  $GG(X, V) = X$ .

Now, take  $\mathcal{W}$  as in the hypotheses. Since  $\mathcal{W}$  is saturated,  $\mathcal{O}_X(V)/\mathcal{W}$  is torsion-free and thus locally free in codimension two. In other words, there exists a proper subvariety  $Y \subsetneq X$ ,  $\text{codim}_X Y \geq 2$ , and a holomorphic vector bundle

$$W \rightarrow U := X \setminus Y$$

on the dense open subset  $U$  such that  $\mathcal{O}_U(W) \simeq \mathcal{W}|_U$  and  $W$  is actually a subbundle of the restriction  $V|_U$  of  $V$  to  $U$ . Of course, over  $U$ , we also have  $\mathcal{W}^*|_U \simeq \mathcal{O}_U(W^*)$ . Since  $(\mathcal{W}^*)^{[\ell]}$  is reflexive by definition and thus normal, we have a surjection

$$H^0(X, (\mathcal{W}^*)^{[\ell]} \otimes A^{-1}) \rightarrow H^0(U, (\mathcal{W}^*)^{[\ell]} \otimes A^{-1}) = H^0(U, (W^*)^{\otimes |\ell|} \otimes A^{-1}),$$

so that  $H^0(U, (W^*)^{\otimes |\ell|} \otimes A^{-1}) = \{0\}$ . From

$$S^{\ell_1} W^* \otimes \cdots \otimes S^{\ell_k} W^* \otimes A^{-1}|_U \subset (W^*)^{\otimes |\ell|} \otimes A^{-1}|_U,$$

we conclude that

$$H^0(U, \text{Gr}^\bullet E_{k,m}^{GG} W^* \otimes A^{-1}) = \{0\}$$

and thus  $H^0(U, E_{k,m}^{GG} W^* \otimes A^{-1}) = \{0\}$ . Then, since each section of  $E_{k,m}^{GG} V^* \otimes A^{-1}$  over  $U$  extends to a section of  $H^0(X, E_{k,m}^{GG} V^* \otimes A^{-1})$ , we have  $U \subset GG(X, V)$  and since  $GG(X, V)$  is a closed set, then  $GG(X, V) = X$ .  $\square$

**2.1. Theorem 1.5 for Demainly–Semple jets.** In this subsection we would like to give a somehow more geometric proof of (a special case of) Theorem 1.5 in the case of invariant jet differentials, even if this case is of course included in what is proved above. For the sake of simplicity we shall assume that  $L \subset T_X$  is a holomorphic line subbundle and not just an injection of sheaves of arbitrary rank. In this case, the hypothesis on  $L$  simply become that  $L$  is not big, that is its Kodaira–Iitaka dimension is not maximal. Invariant jet differentials were introduced in [Dem97] as a refined version of Green–Griffiths’ ones: they are constructed taking invariant holomorphic functions on the  $k$ -jet space by a larger group, namely the full group of  $k$ -jets of biholomorphisms of  $(\mathbb{C}, 0)$ , instead of merely homotheties. The vector bundle  $E_{k,m} V^* \rightarrow X$  of invariant jet differential of order  $k$  and weighted degree  $m$  acting on germs of holomorphic curves tangent to  $V \subset T_X$  can be obtained as a direct image of an invertible sheaf on a tower of projective spaces, as follows.

Start with a directed manifold  $(X, V)$  and define the new directed manifold  $(\widetilde{X}, \widetilde{V})$  to be

$$\widetilde{X} = P(V), \quad \widetilde{V} = \pi_*^{-1}(\mathcal{O}_{P(V)}(-1)) \subset T_{P(V)},$$

where  $\pi: P(V) \rightarrow X$  is the projectivized bundle of lines of  $V$  and  $\mathcal{O}_{P(V)}(-1)$ , which is in turn a subbundle of  $\pi^* V$ , the tautological line bundle of  $P(V)$ . Now, set  $(P_0(V), V_0) = (X, V)$  and define inductively

$$(P_k(V), V_k) = (\widetilde{P_{k-1}(V)}, \widetilde{V_{k-1}})$$

together with the total projection  $\pi_{0,k}: P_k(V) \rightarrow X$ . The functoriality of this construction shows that if  $(Y, W) \subset (X, V)$  is a directed submanifold (i.e.  $Y$  is a smooth submanifold of  $X$ , possibly the whole  $X$ , and  $W \subset T_Y \subset T_X|_Y$  is a holomorphic subbundle of  $V|_Y$ ), then for all positive integers  $k$  we have

$$P_k(W) \subset P_k(V), \quad W_k \subset V_k|_{P_k(W)} \quad \text{and} \quad \mathcal{O}_{P_k(V)}(-1)|_{P_k(W)} = \mathcal{O}_{P_k(W)}(-1),$$

and moreover the projection maps are of course compatible.

If  $\varphi: (\mathbb{C}, 0) \rightarrow X$  is a germ of holomorphic curve tangent to  $V$  then we can define a projectivized lifting  $\varphi_{[k]}: (\mathbb{C}, 0) \rightarrow P_k(V)$ . Such a lifting is tangent to  $V_k$  and satisfies  $\pi_{0,k} \circ \varphi_{[k]} = \varphi$ .

It turns out that for any positive integer  $m$  there is an isomorphism of sheaves

$$(\pi_{0,k})_* \mathcal{O}_{P_k(V)}(m) \simeq \mathcal{O}_X(E_{k,m} V^*).$$

By (the version in the invariant case [Dem97] of) Theorem 1.3, for any positive line bundle  $A \rightarrow X$  and for all positive integers  $k, m$ , we have that

$$f_{[k]}(\mathbb{C}) \subset \text{Bs}(\mathcal{O}_{P_k(V)}(m) \otimes \pi_{0,k}^* A^{-1}),$$

whenever  $f: \mathbb{C} \rightarrow X$  is an entire curve tangent to  $V$  (here,  $\text{Bs}(\bullet)$  stands for the base locus of a line bundle).

With this in mind, we define the *Demailly–Semple locus*  $DS(X, V)$  to be

$$DS(X, V) = DS_A(X, V),$$

where

$$DS_A(X, V) = \bigcap_{k \geq 1} \pi_{0,k} \left( \bigcap_{m \geq 1} \text{Bs}(\mathcal{O}_{P_k(V)}(m) \otimes \pi_{0,k}^* A^{-1}) \setminus P_k(V)^{\text{sing}} \right).$$

Here,  $P_k(V)^{\text{sing}}$  is the complement of the set of values  $\varphi_{[k]}(0)$  reached by all regular (i.e. with non-zero first derivative) germs of curves  $\varphi$ . The fact that  $DS_A(X, V)$  is independent of the ample line bundle  $A \rightarrow X$  can be shown in the same way as in Lemma 2.2.

**Remark 2.3.** Observe that  $P_1(V)^{\text{sing}} = \emptyset$ , and  $P_k(V)^{\text{sing}} \subset \text{Bs}(\mathcal{O}_{P_k(V)}(m))$  for all integers  $m$  (see [Dem97]). Moreover,  $\pi_{0,k}(P_k(V)^{\text{sing}}) = X$ , for  $k \geq 2$ , but if  $f: \mathbb{C} \rightarrow X$  is an entire curve tangent to  $V$  such that  $f_{[k]}(\mathbb{C}) \subset P_k(V)^{\text{sing}}$ , then  $f$  is in fact constant. Thus, it is necessary to remove  $P_k(V)^{\text{sing}}$  from the base loci of the (anti)tautological bundles in order to get a really useful notion of Demailly–Semple locus (compare with the analogous definition in [Dem97, beginning of §13] and [DT10, Introduction], where this minor point was not noticed).

The relation

$$\text{Exc}(X, V) \subset GG(X, V) \subset DS(X, V)$$

is immediate from definitions and Theorem 1.3. We do not know whether  $GG(X, V) = DS(X, V)$ , for instance, under the natural hypothesis that  $\det V^*$  is big.

*Proof of Theorem 1.5 for Demainly–Semple jets.* Fix any ample line bundle  $A \rightarrow X$ . We shall construct for each integer  $k > 0$  a smooth manifold  $X_k \subset P_k(V)$  which projects biholomorphically onto  $X$  via  $\pi_{0,k}$  and such that for all integers  $m > 0$  we have

$$X_k \subset \text{Bs}(\mathcal{O}_{P_k(V)}(m) \otimes \pi_{0,k}^* A^{-1}).$$

The starting datum is a directed manifold  $(X, V)$  together with a rank one holomorphic subbundle  $L \subset V \subset T_X$  such that  $\kappa(L^{-1}) < \dim X$ . We define  $X_k$  to be

$$X_k := P_k(L) \subset P_k(V).$$

Since at each step we are always projectivizing a rank one vector bundle, all  $X_k$  are isomorphic to the starting  $X$ , the isomorphism being given by the projections  $\pi_{0,k}$ . Moreover, since as we have seen

$$\mathcal{O}_{P_k(V)}(-1)|_{P_k(L)} = \mathcal{O}_{P_k(L)}(-1)$$

and, on the other hand,

$$\mathcal{O}_{P_k(L)}(-1) \simeq \pi_{0,k}^* L \simeq L,$$

we deduce that the restriction of  $\mathcal{O}_{P_k(V)}(m) \otimes \pi_{0,k}^* A^{-1}$  to  $X_k = P_k(L)$  has no non-zero holomorphic sections for all positive integers  $m$  by Kodaira's lemma, being isomorphic to  $L^{-\otimes m} \otimes A^{-1}$ . But then, it follows that the base locus of  $\mathcal{O}_{P_k(V)}(m) \otimes \pi_{0,k}^* A^{-1}$  must necessarily contain  $X_k$ , and we are done.  $\square$

We want to underline that we have indeed the much stronger property that not only  $X_k$  is contained in the base locus but also that the restriction of  $\mathcal{O}_{P_k(V)}(m) \otimes \pi_{0,k}^* A^{-1}$  to  $X_k$  has itself no non-zero sections.

### 3. Finite coverings and Green–Griffiths locus

In this section we shall describe the behavior of the Green–Griffiths locus with respect to finite maps. We begin with the following result.

**Proposition 3.1.** *Let  $\rho: X' \rightarrow X$  be a finite surjective morphism of smooth projective manifolds and let  $B \subsetneq X$  be its branch locus. Then, there is an inclusion  $\rho(GG(X')) \subseteq GG(X) \cup B$ .*

*In particular, if  $GG(X') = X'$ , then  $GG(X) = X$  and if  $\rho$  is étale then  $\rho(GG(X')) \subseteq GG(X)$ .*

*Proof.* Fix an ample line bundle  $A \rightarrow X$  and suppose  $x \in X \setminus (GG(X) \cup B)$ . Since  $x \notin GG(X)$ , there exists an integer  $k_0 > 0$  such that for all non constant  $k_0$ -jets

$\varphi_{k_0}: (\mathbb{C}, 0) \rightarrow (X, x)$  there exist a positive integer  $m$  and a global holomorphic section  $P \in H^0(X, E_{k_0, m}^{GG} T_X^* \otimes A^{-1})$  such that  $P(\varphi_{k_0}(0)) \neq 0$ . Next, let  $y \in X'$  be such that  $\rho(y) = x$ . Since  $x \notin B$ , then  $y$  is not a ramification point and thus any non constant  $k_0$ -jet  $\psi_{k_0}: (\mathbb{C}, 0) \rightarrow (X', y)$  gives by composition with  $\rho$  a non constant  $k_0$ -jet  $\rho \circ \psi_{k_0}: (\mathbb{C}, 0) \rightarrow (X, x)$  for which therefore there exist an integer  $m > 0$  and a global holomorphic section  $P \in H^0(X, E_{k_0, m}^{GG} T_X^* \otimes A^{-1})$  such that  $P((\rho \circ \psi_{k_0})(0)) \neq 0$ . Thus,  $\rho^* P$  is a global holomorphic section  $\rho^* P \in H^0(X', E_{k_0, m} T_{X'}^* \otimes \rho^* A^{-1})$  such that  $(\rho^* P)(\psi_{k_0}(0)) \neq 0$  (notice that  $\rho^* A$  is ample since  $\rho$  is a finite morphism), and therefore  $y \notin GG(X')$ .

The last assertion is clear since both  $GG(X)$  and  $B$  are Zariski closed sets.  $\square$

**Corollary 3.2.** *Let  $\Phi \in \text{Aut}(X)$  be an automorphism. Then,  $\Phi(GG(X)) = GG(X)$ .*

**Remark 3.3.** Suppose that we have a surjective morphism between two projective manifolds  $f: Y \rightarrow X$ , and let  $V \subset T_Y$  be a vector subbundle such that  $df|_V: V \rightarrow f^* T_X$  is injective (we only mention the case of vector subbundles and not that of general subsheaves for the sake of simplicity). It is then immediate to check just following exactly the same proof of the proposition above that, more generally, we have  $f(GG(Y, V)) \subset GG(X)$ . This stronger version will be used in Section 6.

For the converse, we suppose that the morphism is étale. We shall see later that this hypothesis is in fact necessary.

**Proposition 3.4.** *If  $\rho: X' \rightarrow X$  is a finite étale cover of smooth projective manifolds, then  $\rho^{-1}(GG(X)) \subseteq GG(X')$ .*

*In particular, if  $GG(X) = X$ , then  $GG(X') = X'$ .*

*Proof.* Let us first reduce to the case when the finite étale cover is moreover Galois: this is done as follows. We claim that there exists a finite Galois étale cover  $X'' \rightarrow X$  which factorizes through  $X'$ . Once the proposition is known for Galois covers, just apply it to  $X'' \rightarrow X$  and then Proposition 3.1 to  $X'' \rightarrow X'$  to get the desired result as follows. Let  $\mu: X'' \rightarrow X' \rightarrow X$  be a Galois cover factorizing through  $\nu: X'' \rightarrow X'$ ,  $x \in GG(X)$  and  $z \in \rho^{-1}(x) \subset X'$ . Then, there exists  $y \in \mu^{-1}(x) \subset X''$  such that  $\nu(y) = z$ . Since  $\mu^{-1}(x) \subset GG(X'')$  then, by Proposition 3.1,  $\nu(y) \in GG(X')$  for any  $y \in \mu^{-1}(x)$ , and thus  $z \in GG(X')$ .

The claim is a consequence of the correspondence between subgroups of the fundamental group and étale covers and the following elementary lemma.

**Lemma 3.5.** *Let  $H \leq \Gamma$  be a subgroup of finite index in  $\Gamma$ . Then, there exists a normal subgroup  $N \trianglelefteq \Gamma$  of finite index in  $\Gamma$  and such that  $N$  is contained in  $H$ .*

*Proof.* The group  $\Gamma$  acts on the set of, say, left cosets  $G/H$  by permutation. Then, we have a group homomorphism  $\Gamma \rightarrow \mathfrak{S}(G/H)$  from  $\Gamma$  to the finite group of permutations of  $G/H$ . It is straightforward to verify that its kernel  $N$  has the required properties.  $\square$

So, let  $\rho: X' \rightarrow X$  be a finite Galois étale cover, say of degree  $d$ , and let  $\text{Aut}(\rho)$  be its deck transformation group, which acts by biholomorphisms freely and transitively on each fiber, by hypothesis. Given an ample line bundle  $A \rightarrow X$ , and any global holomorphic section  $Q \in H^0(X', E_{k,m}^{GG} T_{X'}^* \otimes \rho^* A^{-1})$ , we can recover, by “averaging”  $Q$  with the deck transformations, a global holomorphic section

$$\bigotimes_{\Phi \in \text{Aut}(\rho)} \Phi^* Q =: \tilde{P} \in H^0(X', E_{k,dm}^{GG} T_{X'}^* \otimes \rho^* A^{-d})$$

invariant by  $\text{Aut}(\rho)$ , which thus descends to a global holomorphic section  $P \in H^0(X, E_{k,dm}^{GG} T_X^* \otimes A^{-d})$ .

If  $y \in X' \setminus GG(X')$ , take  $k_0 > 0$  such that for all non constant  $k_0$ -jets  $\psi_{k_0}: (\mathbb{C}, 0) \rightarrow (X', y)$  there exist an integer  $m > 0$  and a global holomorphic section  $Q \in H^0(X, E_{k_0,m}^{GG} T_X^* \otimes \rho^* A^{-1})$  such that  $Q(\psi_{k_0}(0)) \neq 0$ .

Let  $\{y_1 := y, \dots, y_d\}$  be the orbit of  $y$  under  $\text{Aut}(\rho)$ . Then, by Corollary 3.2,  $y_i \in X' \setminus GG(X')$  for  $i = 1, \dots, d$ .

Next, consider a  $k_0$ -jet  $\varphi_{k_0}: (\mathbb{C}, 0) \rightarrow (X, x)$ , where  $x = \rho(y)$ , and its (unique) lifting  $\tilde{\varphi}_{k_0}^i$  to  $X'$  such that  $\tilde{\varphi}_{k_0}^i(0) = y_i$ . Thus, we can find a  $Q_i \in H^0(X, E_{k_0,m}^{GG} T_X^* \otimes \rho^* A^{-1})$  such that  $Q_i(\tilde{\varphi}_{k_0}^i(0)) \neq 0$ . Considering sections of the form  $Q := Q_1 + \epsilon_2 Q_2 + \dots + \epsilon_d Q_d$  we see easily that choosing the scalars  $\epsilon_i$  sufficiently small, we obtain a section  $Q \in H^0(X, E_{k_0,m}^{GG} T_X^* \otimes \rho^* A^{-1})$  such that  $Q(\tilde{\varphi}_{k_0}^i(0)) \neq 0$  for  $i = 1, \dots, d$ . Indeed, by induction on  $d$ , suppose that  $\epsilon_2, \dots, \epsilon_{d-1}$  are chosen in such a way that  $S := Q_1 + \epsilon_2 Q_2 + \dots + \epsilon_{d-1} Q_{d-1}$  satisfies  $S(\tilde{\varphi}_{k_0}^i(0)) \neq 0$  for  $i = 1, \dots, d-1$ . If  $S(\tilde{\varphi}_{k_0}^d(0)) \neq 0$  we are done taking  $\epsilon_d = 0$ . Otherwise  $S(\tilde{\varphi}_{k_0}^d(0)) = 0$  and we choose  $\epsilon_d \neq 0$  sufficiently small so that  $S(\tilde{\varphi}_{k_0}^i(0)) + \epsilon_d Q_d(\tilde{\varphi}_{k_0}^i(0)) \neq 0$  for  $i = 1, \dots, d-1$ .

Finally, the averaging process above applied to this particular  $Q$  then gives a  $P \in H^0(X, E_{k_0,dm}^{GG} T_X^* \otimes A^{-d})$  such that  $P(\varphi_{k_0}(0)) \neq 0$ , as it is straightforwardly checked, and  $x \notin GG(X)$ .  $\square$

**Remark 3.6.** Moreover, the averaging process above shows that in the Galois situation if  $H^0(X', E_{k,m}^{GG} T_{X'}^*) \neq 0$  for some  $k, m > 0$ , then  $H^0(X, E_{k,dm}^{GG} T_X^*) \neq 0$  where  $d$  is the degree of the covering.

The next example, together with the remark above, shows that the étale assumption is necessary.

**Example 3.7.** Let  $X \subset \mathbb{P}^n$  a smooth hypersurface of degree  $d$ , cut out by the single equation  $P(z_0, \dots, z_n) = 0$ . Consider the smooth hypersurface  $X' \subset \mathbb{P}^{n+1}$  cut out by the equation  $z_{n+1}^d - P(z_0, \dots, z_n) = 0$ . The map  $\pi: X' \rightarrow \mathbb{P}^n$  given by the projection onto the first  $n+1$  coordinates is a branched Galois cover whose automorphism group is  $\mathbb{Z}/d\mathbb{Z}$  (acting by multiplying  $z_{n+1}$  by powers of a primitive  $d$ th root of unity), ramified along  $X$ . By [Div09], we know that if  $d$  is large enough, then  $H^0(X', E_{k,m}^{GG} T_{X'}^*) \neq 0$  for  $m \gg k \gg 1$ . On the other hand, since  $T_{\mathbb{P}^n}$  is ample, for all  $k$ -tuples of integers  $\ell_1, \dots, \ell_k > 0$ , the graded terms  $H^0(\mathbb{P}^n, S^{\ell_1} T_{\mathbb{P}^n}^* \otimes \dots \otimes S^{\ell_k} T_{\mathbb{P}^n}^*) = 0$  and thus  $H^0(\mathbb{P}^n, E_{k,m}^{GG} T_{\mathbb{P}^n}^*) = 0$  for all  $k, m > 0$ .

Let us finally remark that one knows from [DMR10] that generic projective hypersurfaces of high degree  $X \subset \mathbb{P}^n$  have  $GG(X) \neq X$ .

#### 4. Green–Griffiths locus for product manifolds

In this section we will show the following consequence of Theorem 1.5 (compare with Corollary 1.7).

**Corollary 4.1.** *Let  $X$  be a complex projective manifold and suppose that  $X$  splits, up to finite (possibly branched) coverings, into a nontrivial product  $Y \times Z$ . Then,  $GG(X) = X$ .*

To prove Corollary 4.1, we will need the following two elementary lemmata.

**Lemma 4.2.** *Let  $E \rightarrow X$  be a rank  $r$  holomorphic vector bundle on a projective manifold  $X$  of dimension  $n$ , and let  $A \rightarrow X$  be an ample line bundle. If, for some integer  $m_0 > 0$ ,  $H^0(X, S^{m_0} E \otimes A^{-1}) \neq 0$ , then there exists a constant  $C > 0$  such that for all sufficiently large integers  $m > 0$  we have*

$$C^{-1} m^{n+r-1} \leq h^0(X, S^m E) \leq C m^{n+r-1}.$$

Moreover, this is the maximum possible growth rate for the dimension of the spaces of global sections of symmetric powers of  $E$ .

*Proof.* Let  $\pi: \mathbb{P}(E) \rightarrow X$  be the projective bundle of hyperplanes of  $E$  and  $\mathcal{O}_E(1) \rightarrow \mathbb{P}(E)$  the tautological quotient line bundle associated to  $E$ . We have that  $\dim \mathbb{P}(E) = n + r - 1$  and, using the projection formula, it is well known that

$$(1) \quad H^0(\mathbb{P}(E), \mathcal{O}_E(m) \otimes \pi^*G) \simeq H^0(X, S^m E \otimes G),$$

for any line bundle  $G \rightarrow X$ . Since  $\mathcal{O}_E(1)$  is  $\pi$ -ample, there exists a positive integer  $\ell_0$  such that  $\mathcal{O}_E(1) \otimes \pi^*A^{\ell_0}$  is ample. On the other hand, by hypothesis and (1), we have that  $\mathcal{O}_E(m_0) \otimes \pi^*A^{-1}$  is effective. Now, fix any ample line bundle  $F \rightarrow \mathbb{P}(E)$  and let  $k_0$  be a positive integer such that  $\mathcal{O}_E(k_0) \otimes \pi^*A^{\ell_0 k_0} \otimes F^{-1}$  is effective. Then,

$$\mathcal{O}_E(k_0(m_0\ell_0 + 1)) \otimes F^{-1} \simeq \mathcal{O}_E(m_0\ell_0 k_0) \otimes \pi^*A^{-\ell_0 k_0} \otimes \mathcal{O}_E(k_0) \otimes \pi^*A^{\ell_0 k_0} \otimes F^{-1}$$

is effective.

Thus, by Kodaira's lemma,  $\mathcal{O}_E(1)$  is big and there exists a constant  $C > 0$  such that for all sufficient large integers  $m > 0$  we have

$$C^{-1} m^{n+r-1} \leq h^0(\mathbb{P}(E), \mathcal{O}_E(m)) \leq C m^{n+r-1},$$

and this is the maximum possible growth rate of the dimension of the spaces of global sections of powers of  $\mathcal{O}_E(1)$ . Thanks to (1), the same holds for symmetric powers of  $E$  on  $X$ .  $\square$

**Lemma 4.3.** *Let  $f: X \rightarrow Y$  be a surjective morphism with connected fibers of smooth projective manifolds and suppose that  $n = \dim X > \dim Y = m$ . If  $E \rightarrow Y$  is any holomorphic vector bundle, then for any ample line bundle  $A \rightarrow X$  we have  $H^0(X, f^*E \otimes A^{-1}) = 0$ .*

*Proof.* Suppose that  $E$  is of rank  $r$ . If, by contradiction,  $H^0(X, f^*E \otimes A^{-1}) \neq 0$ , then, by Lemma 4.2, there exists a positive constant  $C$  such that

$$C^{-1} k^{n+r-1} \leq h^0(X, f^*S^k E) \leq C k^{n+r-1},$$

for all sufficiently large integers  $k$ . On the other hand, by the projection formula,  $H^0(X, f^*V) \simeq H^0(Y, V)$ , for any holomorphic vector bundle  $V \rightarrow Y$ . But then,

$$C^{-1} k^{n+r-1} \leq h^0(X, f^*S^k E) = h^0(Y, S^k E) \leq B k^{m+r-1},$$

for some positive constant  $B$  and all sufficiently large  $k$ , contradiction.  $\square$

We are now in a good shape to give a short proof of Corollary 4.1.

*Proof of Corollary 4.1.* Let  $\rho: \tilde{X} \rightarrow X$  be a finite surjective morphism such that  $\tilde{X} \simeq Y \times Z$  and let  $A \rightarrow \tilde{X}$  be an ample line bundle. Then,  $0 < \dim Y < \dim \tilde{X}$  and  $\text{pr}_1^* T_Y$  is a holomorphic subbundle of  $T_{\tilde{X}}$ , where  $\text{pr}_1: \tilde{X} \rightarrow Y$  is the first projection. For any integer  $m > 0$ , by Lemma 4.3 applied to  $(T_Y^*)^{\otimes m}$ , we have that  $H^0(\tilde{X}, (\text{pr}_1^* T_Y^*)^{\otimes m} \otimes A^{-1}) = 0$ , so that by Theorem 1.5 we have  $GG(\tilde{X}) = \tilde{X}$ . Therefore, by Proposition 3.1,  $GG(X) = X$ .  $\square$

**Examples.** Using the preceding results it is already possible to produce many examples of projective manifolds of general type covered by their Green–Griffiths locus. Let us consider the case of surfaces, although easy generalizations to higher dimensions can be given.

The easiest application of Corollary 4.1 is certainly to take  $X = C_1 \times C_2$  a product of 2 compact Riemann surfaces of genus  $g_i \geq 2$  which gives an example of general type and even hyperbolic, as observed in the Introduction.

Starting from these examples and using Proposition 3.1, one can construct more interesting examples taking quotient by finite groups.

First, let us consider symmetric products of curves. Given a smooth projective curve  $C$  of genus  $g$ , take  $X := C^{(2)} = C \times C / \mathfrak{S}_2$ , the quotient by the symmetric group. If  $g \geq 3$  then  $C^{(2)}$  is of general type, hyperbolic [SZ00] and satisfies  $GG(X) = X$ .

One can also consider the algebraic surfaces whose canonical models arise as quotients  $X = (C_1 \times C_2)/G$  of the product  $C_1 \times C_2$  of two curves of genera  $g_1 := g(C_1), g_2 := g(C_2) \geq 2$ , by the action of a finite group  $G$  (in other words,  $X$  has only rational double points as singularities). The minimal resolution  $S$  of the quotient  $X = (C_1 \times C_2)/G$  is called a product–quotient surface and has been intensively studied in [BCGPI2]. This gives examples of surfaces of general type  $S$  such that  $GG(S) = S$  which are not hyperbolic whenever the action of  $G$  is not free (i.e. whenever  $X$  is singular) and can have in some cases finite fundamental group.

In particular considering the universal cover, one obtains simply connected examples of surfaces of general type covered by their Green–Griffiths locus. Moreover, unlike previous examples, the distribution of rational, elliptic or entire curves on these surfaces seems not to be known.

## 5. Green–Griffiths locus of locally reducible quotients of bounded symmetric domains

For basic results and notation about hermitian symmetric domains, we refer to [Mok89]. Let  $\Omega = G_0/K$  be a bounded symmetric domain, where  $G_0 = \text{Aut}_0(\Omega)$  is the connected component of the identity of the automorphisms group of  $\Omega$ , and let  $\Gamma \subset \text{Aut}(\Omega)$  be a lattice. The quotient  $X = \Omega/\Gamma$  is said to be *locally irreducible* (resp. *locally reducible*) if  $\Omega$  is irreducible (resp. reducible) as a hermitian symmetric space.

**Definition 5.1.** We say that  $X = \Omega/\Gamma$  is *reducible* if there exists a subgroup  $\Gamma_0 \subset \Gamma$  of finite index and a decomposition  $\Omega \simeq \Omega_1 \times \Omega_2$  into a product

of bounded symmetric domains  $\Omega_1$  and  $\Omega_2$  such that  $\Gamma_0 \simeq \Gamma_1 \times \Gamma_2$  and  $\Gamma_i \subset \text{Aut}(\Omega_i)$ ,  $i = 1, 2$ . Otherwise,  $X$  is said to be *irreducible*.

In particular, a reducible quotient is by definition locally reducible. Geometrically, the reducibility of  $X$  means that, up to a finite covering,  $X$  can be decomposed isometrically in a non trivial way. We now concentrate on the locally reducible case.

**Theorem 5.2.** *Let  $X$  be a complex projective manifold uniformized by a reducible bounded symmetric domain. Then,  $GG(X) = X$ .*

*Proof.* The proof splits naturally into two parts: the reducible and the irreducible case. We first treat the reducible case.

**The case of reducible quotients.** Let  $X$  be a reducible quotient. By definition, there exists a decomposition  $\Omega \simeq \Omega_1 \times \Omega_2$  into a product of bounded symmetric domains and a finite étale covering  $\tilde{X}$  of  $X$  which decomposes into a product  $\tilde{X} \simeq Y \times Z$  whose factors are uniformized respectively by  $\Omega'$  and  $\Omega''$ . Since  $\tilde{X}$  is a product, then, by Corollary 4.1,  $GG(X) = X$ .

**The case of irreducible quotients.** Let us treat first the particular case of an irreducible quotient of a polydisk, in order to give the flavor of the proof in a very explicit context.

Fix a co-compact subgroup  $\Gamma \subset \text{PSL}(2, \mathbb{R})^n$  acting freely and properly discontinuously on  $\Delta^n$ . Suppose that  $\Delta^n / \Gamma$  is irreducible, so that by the so-called Density Lemma [Rag72, Cor. (5.21) and Thm. (5.22), p. 86] the projection

$$\Gamma \rightarrow \text{PSL}(2, \mathbb{R})$$

onto, say, the first factor has dense image  $\Gamma_1$  (it is indeed dense onto every factor). Call  $z \in \Delta$  the complex coordinate of the first factor and  $w = (w_1, \dots, w_{n-1}) \in \Delta^{n-1}$  the complex coordinates of the last  $n-1$  factors. We begin with a preliminary elementary lemma.

**Lemma 5.3** (Compare with [SB95]). *Let  $\eta = f(z, w) (dz)^{\otimes m}$  be a symmetric differential of degree  $m$  on  $\Delta_z \times \Delta_w^{n-1}$ . Suppose that  $\eta$  is  $\Gamma$ -invariant. Then,  $\eta$  vanishes identically.*

**Remark 5.4.** This lemma should be regarded as a consequence of the following classical result for automorphic forms. Let  $\Gamma \subset \mathrm{PSL}(2, \mathbb{R})^n$  be an irreducible discrete subgroup with compact quotient  $\Delta^n / \Gamma$ , and  $f$  a  $\Gamma$ -automorphic form of weight  $2r$ ,  $r = (r_1, \dots, r_n) \in \mathbb{Z}^n$ . If  $r_1 \cdots r_n = 0$ , then  $f$  must vanish identically. The same statement holds true for instance in the following more general setting: the quotient  $\Delta^n / \Gamma$  is not necessarily compact but  $\Gamma$  is commensurable with the Hilbert modular group (for these and related statements see for instance [Fre90]). This latter version will be used in Section 6.2.

We shall return to the existence of such groups  $\Gamma$  in Subsection 5.1.1, and now we give an elementary proof of the above lemma.

*Proof.* Consider the smooth real function given by taking the Poincaré norm of  $\eta$ :

$$(z, w) \mapsto |f(z, w)|(1 - |z|^2)^m.$$

This is a  $\Gamma$ -invariant smooth function, thus defined on the compact quotient  $X = \Delta^n / \Gamma$ . Let  $p \in X$  be a point where this function attains its maximum and consider the discrete set of points  $\{(z_{(j)}, w_{(j)})\}_{j \in J} \in \Delta \times \Delta^{n-1}$  which are in the preimage of  $p$  by the quotient map. For each  $j \in J$ , the holomorphic map defined on the polydisk  $\{z_{(j)}\} \times \Delta_w$  by

$$w \mapsto f(z_{(j)}, w)$$

attains a maximum at the interior point  $w_{(j)}$  and so it is constant. So, all the  $w_\lambda$ -derivatives vanish:

$$\partial f / \partial w_\lambda(z_{(j)}, w) = 0, \quad \text{for all } j \in J, \lambda = 1, \dots, n-1, \text{ and } w \in \Delta^{n-1}.$$

By the density of  $\Gamma_1$ , the set  $\{z_{(j)}\}_{j \in J}$  is dense in  $\Delta$ , so  $\partial f / \partial w_\lambda \equiv 0$ ,  $\lambda = 1, \dots, n-1$ , and  $f$  does not depend on  $w$ . Therefore  $\eta$  does only depend on  $z$  and can thus be regarded as a symmetric differential of degree  $m$  on  $\Delta_z$ , invariant by the action of  $\Gamma_1$  on  $\mathrm{Aut}(\Delta_z)$  which has, once again by hypothesis, dense image. But then,  $\eta \equiv 0$ .  $\square$

Now, consider the projective  $n$ -dimensional complex manifold  $X = \Delta^n / \Gamma$ . The holomorphic foliation by disks  $\tilde{\mathcal{F}}$  on  $\Delta^n$  generated by  $\partial / \partial z$  descends to a smooth foliation by curves  $\mathcal{F}$  on  $X$ . Consider its tangent bundle  $T_{\mathcal{F}}$ : it is a rank one holomorphic subbundle of the tangent bundle  $T_X$  of  $X$ .

**Proposition 5.5** (Compare for instance with [Bru04]). *The canonical bundle  $K_{\mathcal{F}} = T_{\mathcal{F}}^*$  of  $\mathcal{F}$  has negative Kodaira–Iitaka dimension.*

*Proof.* We have to show that  $H^0(X, K_{\mathcal{F}}^{\otimes \ell}) = \{0\}$  for all integers  $\ell > 0$ . The latter space of global sections identifies canonically with the space of  $\Gamma$ -invariant global holomorphic sections of  $K_{\mathcal{F}}^{\otimes \ell}$  over  $\Delta^n$ . Since  $\widetilde{\mathcal{F}}$  is generated by  $\partial/\partial z$ , these sections are exactly of the form considered in Lemma 5.3, and therefore they vanish identically.  $\square$

In particular  $K_{\mathcal{F}}$  is not big and, by Kodaira's lemma, for any ample line bundle  $A \rightarrow X$  and for any integer  $m > 0$ , we have  $H^0(X, K_{\mathcal{F}}^{\otimes m} \otimes A^{-1}) = 0$ , so Theorem 1.5 applies.

Now, we pass to the general case. Let  $\Omega = \Omega_1 \times \cdots \times \Omega_k$ ,  $k \geq 2$ , be the decomposition of  $\Omega$  into irreducible components. A classical theorem of Cartan (which can be found in [Nar71, Chap. 5]) states that all automorphisms of  $\Omega$  are given by automorphisms of individual irreducible factors and by permutation of isomorphic factors. It follows that for any lattice  $\Gamma \subset \text{Aut}(\Omega)$ , there is a subgroup  $\Gamma_0 \subset \Gamma$  of finite index such that  $\Gamma_0 \subset \text{Aut}_0(\Omega) = \text{Aut}(\Omega_1) \times \cdots \times \text{Aut}(\Omega_k)$ . Thanks to Proposition 3.1, we can thus suppose without loss of generality that the fundamental group of  $X$ , seen as a lattice inside  $\text{Aut}(\Omega)$ , is contained in the connected component of the identity  $\text{Aut}_0(\Omega)$ . In particular we can suppose that the action of  $\Gamma$  preserves the factors and thus we have a corresponding splitting

$$T_X \simeq V_1 \oplus \cdots \oplus V_k$$

of the tangent bundle of  $X$  such that  $\pi^* V_i = T_{\Omega_i}$ , where  $\pi: \Omega \rightarrow X$  is the quotient map.

Let  $K_X$  be the canonical bundle of  $X$ . Since it is ample there exists an integer  $\ell > 0$  such that  $K_X^\ell$  is effective. We shall show that for any  $i = 1, \dots, k$ , and for any integer  $m > 0$ , we have  $H^0(X, (V_i^*)^{\otimes m} \otimes K_X^{-\ell}) = 0$ , so that  $GG(X) = X$  by Theorem 1.5 (and in fact, more generally, that  $GG(X, V) = X$ , for any subbundle  $V \subset T_X$  containing one of the  $V_i$ 's). For  $i = 1, \dots, k$ , let

$$\Omega_i = G_i / K_i, \quad G_i = \text{Aut}_0(\Omega_i) \quad \text{and} \quad \mathfrak{g}_i = \mathfrak{k}_i + \mathfrak{m}_i,$$

where  $\mathfrak{g}_i$  and  $\mathfrak{k}_i$  are respectively the Lie algebras of  $G_i$  and  $K_i$  and  $\mathfrak{m}_i \simeq \mathfrak{g}_i/\mathfrak{k}_i = T_{\Omega_i, eK_i}$  (here  $eK_i$  is the identity coset). Next, let

$$\varrho_i: K_i \rightarrow \text{GL}(\mathfrak{m}_i)$$

be the isotropy representation of  $K_i$  on  $\mathfrak{m}_i$ , so that the corresponding homogeneous vector bundle is the tangent bundle  $T_{\Omega_i}$ . The irreducible locally homogeneous vector bundle  $V_i \rightarrow X = \Omega/\Gamma$  is thus the one associated to the irreducible representation

$$\begin{aligned} \sigma_i: K_1 \times \cdots \times K_i \times \cdots \times K_k &\rightarrow \text{GL}(\mathfrak{m}_i) \\ (g_1, \dots, g_i, \dots, g_k) &\mapsto \varrho_i(h_i). \end{aligned}$$

Now, fix any integer  $m > 0$  and let  $(\sigma_i^*)^{\otimes m}$  be the representation defining  $(V_i^*)^{\otimes m}$ . Since  $K_i$  is compact, all its representations completely decompose into a direct sum of irreducible ones, thus:

$$(\mathfrak{m}_i^*)^{\otimes m} \simeq E_1 \oplus \cdots \oplus E_N$$

and we have a corresponding decomposition of  $(V_i^*)^{\otimes m}$  into a direct sum of irreducible locally homogeneous vector bundles

$$(V_i^*)^{\otimes m} \simeq W_1 \oplus \cdots \oplus W_N.$$

Now, they all have a natural induced hermitian metric coming from the Bergman metric on  $\Omega_i$ , and the decomposition is as hermitian vector bundles. Moreover, observe that each of the  $W_j$ 's comes from a homogeneous vector bundle on  $\Omega$  which is in fact a pull-back of a homogeneous vector bundle on  $\Omega_i$ . Therefore, the Chern curvature of the  $W_j$ 's is zero whenever evaluated (in its  $(1, 1)$ -form part) on tangent vectors lying on the orthogonal complement of  $V_i$ . In particular, the  $W_j$ 's cannot be of strictly positive Chern curvature in the sense of Griffiths.

To conclude, we use the following deep vanishing theorem which we rephrase in a slightly different way.

**Theorem 5.6** (See [Mok89, Corollary 1' on page 212]). *Let  $\Omega$  be a bounded symmetric domain of complex dimension  $\geq 2$  and  $X = \Omega/\Gamma$  be an irreducible quotient of finite volume of  $\Omega$  by a torsion free discrete group  $\Gamma$  of automorphisms. Suppose  $V$  is an irreducible locally homogeneous Hermitian vector bundle on  $X$  which is not of strictly positive Chern curvature in the sense of Griffiths. Then,  $H^0(X, V) = 0$  unless  $V$  is trivial.*

Now, if  $W_j$  is non trivial then, by Theorem 5.6,  $H^0(X, W_j) = 0$  and a fortiori  $H^0(X, W_j \otimes K_X^{-\ell}) = 0$  since  $K_X^\ell$  is effective. If, on the other hand,  $W_j$  is trivial, then

$$W_j \otimes K_X^{-\ell} \simeq (K_X^{-\ell})^{\oplus \text{rk } W_j}$$

and thus  $H^0(X, W_j \otimes K_X^{-\ell}) \simeq H^0\left(X, (K_X^{-\ell})^{\oplus \text{rk } W_j}\right) = 0$ , since  $K_X$  is ample.  $\square$

**Remark 5.7.** Although we were not able to prove it, we strongly expect that in the proof above the case  $W_j$  trivial should not exist. This would give the stronger statement that for any  $i = 1, \dots, k$ , and for any integer  $m > 0$ ,

$$H^0(X, (V_i^*)^{\otimes m}) = 0.$$

**5.1. The case of surfaces.** Let us now speculate more on the case of surfaces. In this case  $\Omega$  is necessarily the bidisk  $\Delta^2$ . We begin with a few words about the existence of irreducible co-compact subgroups  $\Gamma \subset \mathrm{PSL}(2, \mathbb{R})^2$  acting freely and properly discontinuously on  $\Delta^2$ , which is however quite classical (see for instance in [Sha78]).

**5.1.1. Construction of  $\Gamma$ .** Take a quaternion algebra  $A$  which is division and whose center is a totally real quadratic number field  $k$  (for an excellent reference about quaternion algebras and Fuchsian groups see [Kat92]). Assume that

$$A \otimes_{\mathbb{Q}} \mathbb{R} = M(2, \mathbb{R})^2,$$

that is,  $A$  is unramified at the two places corresponding to the two different embeddings of  $k$  into  $\mathbb{R}$ . If  $\mathfrak{D}$  is a maximal order in  $A$ , denote by  $\Gamma(1)$  the group of units in  $\mathfrak{D}$  with reduced norm 1, and identify it with its isomorphic image in  $\mathrm{SL}(2, \mathbb{R})^2$ . Next, call  $\Gamma' = \Gamma(1)/\{\pm 1\}$  the image of  $\Gamma(1)$  in  $\mathrm{PSL}(2, \mathbb{R})^2$ . Then, it is well known that the action of  $\Gamma'$  on  $\Delta^2$  is irreducible and properly discontinuous (see [Sha78, Shi94]); moreover, the fact that  $A$  is division classically implies that this action is co-compact.

Now, since the action is co-compact,  $\Gamma'$  is a group of finite type. By Selberg's theorem, a finitely generated linear group over a field of zero characteristic is virtually torsion-free, i.e. it has some finite index subgroup  $\Gamma$  which is torsion-free. Thus,  $\Gamma$  acts freely on  $\Delta^2$ . Moreover, being of finite index in  $\Gamma'$ , it is straightforward to see that it is again irreducible and with a properly discontinuous, co-compact action.

**5.1.2. Surfaces of general type with a holomorphic foliation by curves.** Let  $S$  be a surface of general type and  $L \subset T_S$  a holomorphic line subbundle, i.e.  $S$  is endowed with a smooth holomorphic foliation  $\mathcal{F}$  by curves whose tangent bundle  $T_{\mathcal{F}}$  is  $L$ . Suppose that the canonical bundle  $K_{\mathcal{F}}$  is not big.

**Proposition 5.8.** *Let  $(S, \mathcal{F})$  be as above. Then,  $S$  is a quotient of the bidisk  $\Delta^2$ .*

*Proof.* Following [Bru97], the existence of such a smooth foliation on the surface of general type  $S$  implies that  $K_S$  is ample and so, by Aubin–Yau,  $S$  is Kähler–Einstein and hence  $T_S$  is  $K_S$ -semistable. The semistability inequality reads

$$(2) \quad c_1(T_{\mathcal{F}}) \cdot c_1(S) \geq \frac{1}{2} c_1(S)^2 > 0.$$

Since  $\mathcal{F}$  is smooth, the Baum–Bott formulae give

$$(3) \quad \begin{aligned} c_2(S) - c_1(T_{\mathcal{F}}) \cdot c_1(S) + c_1(T_{\mathcal{F}})^2 &= 0 \\ c_1(S)^2 - 2 c_1(T_{\mathcal{F}}) \cdot c_1(S) + c_1(T_{\mathcal{F}})^2 &= 0. \end{aligned}$$

If  $T_S$  is stable, the first inequality in (2) is strict and using the second of (3) we obtain  $c_1(T_{\mathcal{F}})^2 > 0$ . Thus,  $T_{\mathcal{F}}$  or its dual  $K_{\mathcal{F}}$  must be big. Since, by (2),  $c_1(K_{\mathcal{F}}) \cdot c_1(K_S) > 0$ , we get that  $K_{\mathcal{F}}$  is big, contradiction.

Therefore,  $T_S$  is polystable but not stable, that is  $T_S = L_1 \oplus L_2$  is a direct sum of two line bundles and by Beauville–Yau’s uniformization theorem [Bea00, Yau93], the universal cover  $\widetilde{S}$  of  $S$  splits as a product of simply connected Riemann surfaces, the decomposition of the tangent bundle lifts and the fundamental group of  $S$  acts diagonally on  $\widetilde{S}$ . Of course, the only possibility for  $\widetilde{S}$  is to be the product of two disks.  $\square$

It is known by [Lu96] that surfaces as above satisfy the Green–Griffiths conjecture since they verify the Chern numbers inequality  $c_1(S)^2 - 2c_2(S) \geq 0$  (to see this just inject the difference of the two identities in (3) into the first inequality in (2)).

Next, assume that  $S$  is a smooth projective surface of general type and  $\mathcal{F}$  a (possibly singular, with at most isolated singularities) holomorphic foliation by curves whose canonical bundle  $K_{\mathcal{F}}$  is not big. By Seidenberg’s theorem, we can suppose without loss of generality that  $\mathcal{F}$  has reduced (or canonical) singularities.

Thus, the birational classification of foliations developed by Brunella and McQuillan (see [Bru97, McQ08] and [Bru04]) tells us that  $\mathcal{F}$  is necessarily of the following two types.

- A Hilbert modular foliation, and thus  $S$  is a Hilbert modular surface, if  $\kappa(K_{\mathcal{F}}) = -\infty$ .
- An isotrivial fibration of genus  $\geq 2$ , if  $\kappa(K_{\mathcal{F}}) = 1$ .

This gives the “singular” analogue of Proposition 5.8. Then, for instance, Hilbert modular surfaces which are minimal resolution of surfaces with cusps give examples of surfaces  $S$  with  $c_1(S)^2 < 2c_2(S)$  such that  $GG(S) = S$ .

Finally, what if  $K_{\mathcal{F}}$  is big? Here is a natural question which was asked to us by M. McQuillan in a private communication.

**Question 5.9.** *Let  $S$  be a surface admitting a holomorphic foliation by curves  $\mathcal{F}$  with canonical singularities. Suppose that  $K_{\mathcal{F}}$  is big. Is it then true that  $GG(S, T_{\mathcal{F}}) \subsetneq S$ ? If moreover  $S$  is of general type, what can be said about  $GG(S)$  in this case?*

## 6. Green–Griffiths locus of locally irreducible quotients of bounded symmetric domains

In this section, we will prove the following theorem.

**Theorem 6.1.** *Let  $X$  be a complex projective manifold uniformized by an irreducible bounded symmetric domain of rank greater than or equal to two. Then,  $GG(X) = X$ .*

Let us explain the idea to prove such a result. Recall that in the locally reducible case we used in an essential way the existence of natural holomorphic foliations on the manifold coming from the irreducible factors of the universal cover. At first glance, it could be therefore tempting to think that the irreducibility of the tangent bundle could be somehow an obstruction for the Green–Griffiths locus to cover the whole manifold. The theorem above tells us that this is indeed not the case.

To prove Theorem 6.1 we need thus to find something which replaces the existence of natural foliations. The idea is somehow to use, instead of the trivial foliations on the universal cover  $\Omega$  of  $X$ , the existence of many polydisks nicely embedded in  $\Omega$ .

**6.1. The Green–Griffiths locus of Shimura varieties: An “all or nothing” principle à la Ullmo–Yafaev.** It is not easy in general to construct totally geodesic subvarieties in arbitrary compact quotients of symmetric hermitian domains. Interesting examples are given by the theory of Shimura varieties. Consider a connected semisimple and simply connected  $\mathbb{Q}$ -anisotropic linear algebraic group  $\mathbb{G}$  over  $\mathbb{Q}$  with associated Lie group  $G = \mathbb{G}(\mathbb{R})$ . Then, for any congruence subgroup  $\Gamma$  which acts without fixed point on  $D = G/K$ , the quotient space  $X = D/\Gamma$  is a smooth projective variety.

Any  $g \in \mathbb{G}_{\mathbb{Q}}$  gives rise to a Hecke correspondence  $T_g$  on  $X$  as follows. Let  $\Gamma_g = \Gamma \cap g^{-1}\Gamma g$ : this is a congruence subgroup of finite index in  $\Gamma$ . The variety  $Y = D/\Gamma_g$  is smooth projective and admits two finite étale maps to  $X$ , given by  $p_1(\Gamma_g x) = \Gamma x$ ,  $p_2(\Gamma_g x) = \Gamma g x$ , for any  $x \in D$ . The action on cycles of  $X$  is given by  $p_2 \circ p_1^*$ .

A direct application of Propositions 3.1 and 3.4 gives the following result which should be seen as a geometric counterpart of the “all or nothing” principle of [UY10] and as a preliminary version of Theorem 6.1.

**Theorem 6.2.** *Let  $X$  be a compact Shimura variety associated to a semisimple, connected and simply connected  $\mathbb{Q}$ -anisotropic algebraic group over  $\mathbb{Q}$ . Then  $GG(X) = \emptyset$  or  $GG(X) = X$ .*

*Proof.* Since the group  $G$  is connected,  $\mathbb{G}_{\mathbb{Q}}$  is dense in  $G = \mathbb{G}(\mathbb{R})$  [PR94]. It follows that if  $Y \subset X$  is a subvariety of  $X$ , the irreducible components of  $T_g(Y)$  for  $g \in \mathbb{G}_{\mathbb{Q}}$  are dense in  $X$ .

We conclude using Propositions 3.1 and 3.4 which give that if  $Y \subset GG(X)$  then  $T_g(Y) \subset GG(X)$ .  $\square$

Now, we give a criterion to decide on which side of the alternative we are.

**Corollary 6.3.** *Let  $X$  be a compact Shimura variety associated to a semisimple, connected and simply connected  $\mathbb{Q}$ -anisotropic algebraic group over  $\mathbb{Q}$ . Suppose that  $X$  contains a subvariety  $Y$  of positive dimension such that  $GG(Y) = Y$ . Then,  $GG(X) = X$ .*

*Proof.* Fix an ample line bundle  $A \rightarrow X$ . First we observe that if  $GG(Y) = Y$  then  $Y \subset GG(X)$ . Indeed, if  $y \in Y$ , for all integer  $k > 0$  there exists a  $k$ -jet  $\varphi_k: (\mathbb{C}, 0) \rightarrow (Y, y)$  such that for all integer  $m > 0$  and all  $Q \in H^0(Y, E_{k,m}^{GG} T_Y^* \otimes A^{-1}|_Y)$ , one has  $Q(\varphi_k(0)) = 0$ . Given any  $P \in H^0(X, E_{k,m}^{GG} T_X^* \otimes A^{-1})$ , restricting  $P$  to  $Y$  we obtain  $P(\varphi_k(0)) = 0$ . But then,  $y \in GG(X)$ .

Finally, since then  $GG(X) \neq \emptyset$ , thanks to Theorem 6.2 we obtain  $GG(X) = X$ .  $\square$

Although we have considered the compact case, the same strategy can be applied in the isotropic case. An interesting and very explicit case is the one of Siegel modular varieties which we shall discuss now in some detail.

**6.2. The Green–Griffiths locus of (compactifications of) Siegel modular varieties.** Let  $n \geq 2$  and

$$\Omega = D_n^{III} := \{Z \in M(n, \mathbb{C}) \mid Z = {}^\dagger Z, \quad \text{Id}_n - {}^\dagger \overline{Z} \cdot Z > 0\}$$

be the classical bounded symmetric domain of type  $III$ , which is holomorphically equivalent to the Siegel upper half–space  $\mathbb{H}_n = \{\tau \in M(n, \mathbb{C}) \mid \tau = {}^\dagger \tau, \text{Im} \tau > 0\}$ . It has complex dimension  $n(n + 1)/2$  and rank  $n$ . The group  $\text{Sp}(2n, \mathbb{R})$  acts transitively on  $\Omega$  and we have indeed a presentation

$$\Omega = \text{Sp}(2n, \mathbb{R}) / U(n)$$

as a homogeneous space. Now, let  $\Gamma \subset \text{Sp}(2n, \mathbb{R})$  be a lattice commensurable with  $\text{Sp}(2n, \mathbb{Z})$ , and consider the quotient manifold  $X = \Omega / \Gamma$  and any smooth compactification  $\overline{X}$  of  $X$ .

**Proposition 6.4.** *The Green–Griffiths locus  $GG(\overline{X})$  of  $\overline{X}$  is the whole manifold.*

*Proof.* There is a totally geodesic polydisk  $\Delta^n \hookrightarrow \Omega$  given by

$$\Delta^n \ni z = (z_1, \dots, z_n) \mapsto z^* = \text{diag}(z_1, \dots, z_n),$$

where  $\text{diag}(z_1, \dots, z_n)$  is the diagonal matrix with  $(z_1, \dots, z_n)$  as entries along the diagonal. This corresponds to the embedding

$$\begin{aligned} \text{SL}(2, \mathbb{R})^n &\hookrightarrow \text{Sp}(2n, \mathbb{R}) \\ M = (M_1, \dots, M_n) &\mapsto M^* = \begin{pmatrix} a^* & b^* \\ c^* & d^* \end{pmatrix} \end{aligned}$$

where, for  $i = 1, \dots, n$ ,

$$M_i = \begin{pmatrix} a_i & b_i \\ c_i & d_i \end{pmatrix} \in \text{SL}(2, \mathbb{R}),$$

$a^* = \text{diag}(a_1, \dots, a_n)$  and similarly for  $b^*$ ,  $c^*$  and  $d^*$ .

More generally (the following construction is taken from [Fre79]), given  $A \in \text{GL}(n, \mathbb{R})$ , one can consider the map  $\Delta^n \hookrightarrow \Omega$ , given by  $\Delta^n \ni z = (z_1, \dots, z_n) \mapsto A^t z^* A$ . In order to take quotients, one defines

$$\Gamma_A := \left\{ M \in \text{SL}(2, \mathbb{R})^n \text{ such that } \begin{pmatrix} A^t & 0 \\ 0 & A^{-1} \end{pmatrix} M^* \begin{pmatrix} A^t & 0 \\ 0 & A^{-1} \end{pmatrix}^{-1} \in \Gamma \right\}.$$

Indeed, it is straightforward to see that we thus have a well-defined map

$$\phi_A: \Delta^n / \Gamma_A \rightarrow X.$$

Again following [Fre90], consider a totally real number field  $\mathbb{K}$  of degree  $n$  together with its embedding with dense image

$$\begin{aligned} \mathbb{K} &\hookrightarrow \mathbb{R}^n \\ \alpha &\mapsto (\alpha^{(1)}, \dots, \alpha^{(n)}). \end{aligned}$$

Next, fix a basis  $\Lambda = \{\alpha_1, \dots, \alpha_n\}$  of  $\mathbb{K}$  as a  $\mathbb{Q}$ -vector space. Then, the matrices

$$A_\Lambda = (\alpha_i^{(j)})_{i,j=1,\dots,n} \in \text{GL}(n, \mathbb{K})$$

have the property that the corresponding  $\Gamma_{A_\Lambda}$  is commensurable with the Hilbert modular group associated to  $\mathbb{K}$ . Moreover, such matrices  $A_\Lambda$  are clearly dense in  $\text{GL}(n, \mathbb{R})$  as  $\Lambda$  runs through all possible bases.

Now, take a global jet differential of order  $k$  and weighted degree  $m$  over  $\overline{X}$ ,  $P \in H^0(\overline{X}, E_{k,m}^{GG} T_{\overline{X}}^*)$ . Taking the pull-back  $\phi_{A_\Lambda}^* P$ , we obtain a  $k$ -jet differential on  $\Delta^n / \Gamma_{A_\Lambda}$ . Therefore, by Remark 5.4 and an evident variation in the non-compact case of the subsequent proposition,  $\phi_{A_\Lambda}^* P$  must vanish on jets tangent to the directions given by the foliations defined by the factors. Thus, we have that  $\phi_{A_\Lambda}(\Delta^n / \Gamma_A) \subset GG(\overline{X})$ . By density, we finally get  $GG(\overline{X}) = \overline{X}$ .  $\square$

**The characteristic bundle [Mok89, Chapter 6].** In order to “globalize” this polydisk approach, the right tool turns out to be the characteristic bundle introduced by Mok. We recall below one possible construction as well as some basic features of this important object.

So, let  $\Omega = G_0/K$  be an irreducible bounded symmetric domain of rank  $\geq 2$  endowed with its Bergman metric  $\omega$  and  $X = \Omega/\Gamma$  a quotient of  $\Omega$  by a torsion-free discrete group of automorphisms. By a slight abuse of notation, we will still call  $\omega$  the induced metric on  $X$ . Now, consider the projectivized bundle  $\pi: P(T_X) \rightarrow X$  of lines of  $T_X$  and the corresponding tautological line bundle  $\mathcal{O}_{P(T_X)}(-1) \subset \pi^*T_X \rightarrow P(T_X)$  with the natural hermitian metric  $h$  induced by  $\omega$ .

Next, let  $\Theta_\omega(T_X)$  be the Chern curvature of  $T_X$  with respect to the metric  $\omega$ . For each  $x \in X$  and  $v \in T_{X,x} \setminus \{0\}$ , consider the hermitian form on  $T_{X,x}$  defined by

$$(4) \quad T_{X,x} \times T_{X,x} \ni (\xi, \eta) \mapsto \frac{1}{\|v\|_\omega^2} \omega(\Theta_\omega(T_X)(v, \bar{v}) \cdot \xi, \eta).$$

It is the hermitian form associated to the Griffiths curvature of  $(T_X, \omega)$  in the direction given by  $v$ , which is therefore semi-negative. Call  $\mathcal{N}_{x,[v]} \subset T_{X,x}$  its zero eigenspace: it is clearly neither zero nor the whole space. We say that  $[v_0]$  is a *characteristic direction* at the point  $x \in X$  if  $\dim \mathcal{N}_{x,[v_0]}$  is maximum among  $\dim \mathcal{N}_{x,[v]}$ , for  $[v] \in T_{X,x} \setminus \{0\}$ . This maximal dimension does depend only on  $\Omega$ : we call it  $n(\Omega)$ , the *null dimension* of  $\Omega$ .

**Definition 6.5.** The set  $\mathcal{S} = \mathcal{S}(X) \subset P(T_X)$  of characteristic directions together with the induced projection  $\pi|_{\mathcal{S}}: \mathcal{S} \rightarrow X$  onto  $X$  is called the *characteristic bundle*.

The characteristic directions are, equivalently [Mok89, Proposition 1 on page 242], the directions which minimize the holomorphic sectional curvature

$$T_{X,x} \setminus \{0\} \ni v \mapsto \frac{1}{\|v\|_\omega^4} \omega(\Theta_\omega(T_X)(v, \bar{v}) \cdot v, v).$$

In algebraic terms, the characteristic vectors (i.e. the non zero vectors which define a characteristic direction) are given by the highest weights of the isotropy representation on the holomorphic tangent space.

It is a remarkable fact that  $\pi|_{\mathcal{S}}: \mathcal{S} \rightarrow X$  is indeed a holomorphic fiber bundle and that  $\mathcal{S}$  is a smooth closed complex submanifold of  $P(T_X)$  of dimension

$$\dim \mathcal{S} = \dim P(T_X) - n(\Omega) = 2 \dim X - 1 - n(\Omega).$$

Even more remarkable, we have the following

**Proposition 6.6** (Mok [Mok89, Proposition 4 on page 262]). *Let  $X = \Omega/\Gamma$  be a compact quotient of an irreducible bounded symmetric domain of rank  $\geq 2$ . Then, for any integer  $m > 0$  and any  $\sigma \in H^0(P(T_X), \mathcal{O}_{P(T_X)}(m))$ , we have that  $\sigma$  vanishes identically on  $\mathcal{S}$ .*

**Remark 6.7.** We stated here a weaker form of the proposition above, which gives in fact – in its full strength – a precise description of the stable base locus of  $\mathcal{O}_{P(T_X)}(1)$  in terms of higher characteristic bundles, see [Mok89] for more details.

In the sequel, we shall need the following slightly refined version of Proposition 6.6, which deals with global sections of the *restriction* of the (anti)tautological bundle to  $\mathcal{S}$ .

**Proposition 6.8** (Compare with [Mok89, Chapter 6, (3.1) and (3.2)]). *Let  $X = \Omega/\Gamma$  be a compact quotient of an irreducible bounded symmetric domain of rank  $\geq 2$ . Then, for any integer  $m > 0$  we have*

$$H^0(\mathcal{S}, \mathcal{O}_{\mathcal{S}}(m)) = \{0\},$$

where  $\mathcal{O}_{\mathcal{S}}(m)$  is the restriction  $\mathcal{O}_{P(T_X)}(m)|_{\mathcal{S}}$ .

In other words, the Kodaira–Iitaka dimension of  $\mathcal{O}_{\mathcal{S}}(1)$  is negative.

*Proof.* The proof relies essentially on the fact that the argument used to prove [Mok89, Proposition 4 on page 262] goes through in this stronger version. We sketch it anyway for the reader’s convenience.

We proceed by contradiction and suppose there exists an integer  $m > 0$  and a non zero section  $\sigma \in H^0(\mathcal{S}, \mathcal{O}_{\mathcal{S}}(m))$ . By a slight abuse of notation, we still call  $h$  the restriction of the natural hermitian metric  $h$  on  $\mathcal{O}_{P(T_X)}(-1)$  to  $\mathcal{O}_{\mathcal{S}}(-1)$ . Then,  $g := (h^m + \sigma \otimes \bar{\sigma})^{1/m}$  defines a hermitian metric on  $\mathcal{O}_{\mathcal{S}}(-1)$  which still has semi-negative curvature, since  $\sigma$  is holomorphic.

**Claim 6.9** (Compare with Mok’s hermitian metric rigidity theorem). In this situation,  $g = C \cdot h$  for some positive constant  $C$ .

This claim implies that  $\|\sigma\|_{h^{-m}}$  is constant. If it were a non zero constant this would imply that  $\sigma$  never vanishes and thus  $\mathcal{O}_{\mathcal{S}}(m)$  is holomorphically trivial. But this is impossible since

**Claim 6.10.** The following inequality holds:

$$\int_{\mathcal{S}} c_1(\mathcal{O}_{P(T_X)}(1)) \wedge \nu^{\dim \mathcal{S}-1} > 0,$$

where  $\nu$  is the Kähler form on  $P(T_X)$  given by  $c_1(\mathcal{O}_{P(T_X)}(1), h^{-1}) + \pi^* \omega$ .

But then  $||\sigma||_{h^{-m}} \equiv 0$  and  $\sigma$  is identically zero, too.  $\square$

It remains to prove the two claims above.

*Proof of Claim 6.9.* We follow almost word-by-word the alternative proof of Mok’s hermitian metric rigidity theorem which makes use of Moore’s ergodicity theorem [Mok89, Chapter 6, §3]: we just need to check that his argument goes through when everything lives on  $\mathcal{S}$  and does not come necessarily from the whole  $P(T_X)$ .

So, write  $g = e^u \cdot h$ , where  $u$  is a smooth real function on  $\mathcal{S}$  and write  $q := n(\Omega)$ . By [Mok89, Chapter 6, formula (2) on page 116], we have that

$$i \partial u \wedge \bar{\partial} u \wedge c_1(\mathcal{O}_{P(T_X)}(-1), h)^{2n-2q-1} \equiv 0 \quad \text{on } \mathcal{S}.$$

This follows from a quite straightforward integral formula over  $\mathcal{S}$ . Now, we lift everything on  $\Omega$ , where the characteristic bundle is trivial

$$\mathcal{S}(\Omega) \simeq \Omega \times \mathcal{S}_0 \subset \Omega \times \mathbb{P}^{n-1} = P(T_\Omega).$$

Here,  $\mathcal{S}_o$  is the set of characteristic direction over a base point  $o \in \Omega$ . In particular we have an identification

$$T_{\mathcal{S}(\Omega), (o, [v])} \simeq T_{\mathcal{S}_o, [v]} \oplus T_{\Omega, o}.$$

From the explicit expression of the curvature  $\Theta_h(\mathcal{O}_{P(T_X)}(-1))$  it is immediate to check that the non-negative  $(1, 1)$ -form  $i \partial u \wedge \bar{\partial} u$  must vanish on  $\mathcal{N}_{o, [v]} \subset T_{\Omega, o}$ . Since  $u$  is real, it follows that

$$(5) \quad du|_{\mathcal{N}_{o, [v]}} \equiv 0.$$

Let  $U(T_X)$  the  $\omega$ -unitary tangent bundle to  $X$  and  $U(\mathcal{S}) \subset U(T_X)$  the subspace of unitary characteristic vectors. Of course,  $u$  pulls-back to a ( $S^1$ -invariant) function on  $U(\mathcal{S})$ , which we still call  $u$  by abuse of notation. Fix a unit characteristic vector  $v$  and let  $L \subset G_0$  be its stabilizer, so that  $U(\mathcal{S}) = \Gamma \backslash G_0 / L$  has a locally homogeneous space (it is more convenient to write  $\Gamma$  on the left for this proof). By pulling-back again we get a positive  $\Gamma$ -invariant and  $L$ -invariant smooth function  $\hat{u}$  on  $G_0$  which we will show to be invariant under the right action of a closed non-compact subgroup  $H \subset G_0$ . Now, by Moore’s ergodicity theorem, every  $H$ -invariant subset of  $\Gamma \backslash G_0$  is either of zero or of full measure, since  $G_0$  is simple and  $\Gamma$  a lattice. If  $\hat{u}: \Gamma \backslash G_0 \rightarrow \mathbb{R}$  were not constant then we could find two positive real numbers  $a < b$  such that the set

$$V_{a,b} := \{y \in \Gamma \backslash G_0 \mid a < \hat{u}(y) < b\}$$

is neither of zero nor of full measure. But by the  $H$ -invariance of  $\hat{u}$ ,  $V_{a,b}$  would be  $H$ -invariant, contradiction. It then follows that  $u$  itself is constant and therefore  $g$  is a constant multiple of  $h$ .

We now come back to the existence of such a closed non-compact subgroup  $H$ : it is obtained as a one-parameter subgroup of transvections as follows. Take a non zero vector  $w \in \mathcal{N}_{o,[v]}$  and let  $\gamma$  be the geodesic on  $\Omega$  determined by  $w$ . Without loss of generality, we can suppose that the characteristic vector  $v$  is unitary and form the curve  $\gamma^* \subset U(\mathcal{S}(\Omega))$  obtained by parallel transport of  $v$  along  $\gamma$  (here we use that  $\mathcal{S}$  and hence  $U(\mathcal{S})$  are invariant by parallel transport). By construction, the curve  $t \mapsto \gamma^*(t)$  is pointwise tangent to  $\mathcal{N}_{\gamma(t),[\gamma^*(t)]}$ , so that  $u$  is constant along  $\gamma^*$  by (5). By the same argument,  $u$  is also constant along the image  $\varphi(\gamma^*)$  of  $\gamma^*$  by any isometry  $\varphi \in G_0$ . By the general theory,  $\gamma^*$  is the orbit  $H(v) = H(e \bmod L)$  of  $v$  under a non-compact one-parameter family  $H \subset G_0$  of isometries. But then, for  $\varphi \in G_0$ ,  $u$  is constant along the  $H$ -orbit  $\varphi H \bmod L = \phi(\gamma^*)$ . This means exactly that  $\hat{u}$  on  $\Gamma \backslash G_0$  is invariant under the action of  $H$ .  $\square$

*Proof of Claim 6.10.* The Chern curvature  $i \Theta_h(\mathcal{O}_{P(T_X)}(-1))$  of  $h$  at a point  $(x, [v]) \in P(T_X)$  is roughly given by a “vertical” term, which is nothing but minus the Fubini–Study metric, and a “horizontal” term, which is given by the  $(1, 1)$ -form associated to the hermitian form in (4). In particular, if  $[v]$  is a characteristic direction, at  $(x, [v])$  it is semi-negative of rank  $2n - 1 - n(\Omega)$ . So, the restriction of  $i \Theta_h(\mathcal{O}_{P(T_X)}(-1))$  to  $\mathcal{S}$  is also semi-negative of rank at least  $2n - 1 - 2n(\Omega) > 0$ . Therefore, along  $\mathcal{S}$ , the trace with respect to  $v$

$$c_1(\mathcal{O}_{P(T_X)}(1), h^{-1}) \wedge v^{\dim \mathcal{S}-1} = -c_1(\mathcal{O}_{P(T_X)}(-1), h) \wedge v^{\dim \mathcal{S}-1}$$

is strictly positive and thus so is its integral over  $\mathcal{S}$ .  $\square$

**Foliating the characteristic bundle by minimal disks.** Let  $\Omega = G_0/K$  be a bounded symmetric domain of rank  $\geq 2$  and  $X = \Omega/\Gamma$  a compact quotient. We shall recall how to construct a natural (smooth) holomorphic foliation by curves  $\mathcal{M}$  on  $\mathcal{S}$ . The foliation in question is the one associated to minimal disks in  $\Omega$ . Let us describe how one can construct it. Let  $\Omega^\vee = G_c/K$  be the compact dual of  $\Omega$ , where  $G_c$  is a compact real form of the complexification of  $G_0$ . Write  $\Omega \subset \subset \mathbb{C}^{\dim \Omega} \subset \Omega^\vee$  for the Harish–Chandra and Borel embeddings. As in the non compact case, one defines the notion of characteristic vectors on  $\Omega^\vee$  as highest weights of the isotropy representations on the holomorphic tangent space. Since the isotropy representations for  $\Omega$  and  $\Omega^\vee$  are the same at points of  $\Omega$ , a tangent vector to  $\Omega$  is characteristic for  $\Omega$  if and only if it is characteristic for  $\Omega^\vee$ . Next, there is a notion of *minimal rational curve* in  $\Omega^\vee$ : these are rational curves representing the positive generator of  $H_2(\Omega^\vee, \mathbb{Z}) \simeq \mathbb{Z}$ .

**Proposition 6.11** (Mok [Mok89, Proposition 1 on page 143]). *For any  $x \in \Omega^\vee$  and any characteristic vector  $v \in T_{\Omega^\vee, x}$  there exists a unique minimal rational curve  $C$  passing through  $x$  such that  $T_{C, x} = [v]$ . Furthermore, all minimal rational curves are obtained in this way.*

Now, it can be shown that the intersection of any minimal rational curve with  $\Omega$  is biholomorphic to a disk. Such disks in  $\Omega$  are called *minimal disks*. We shall still call minimal disk the image in  $X$  of a minimal disk in  $\Omega$  under the uniformization map  $\Omega \rightarrow X$ .

Consider the projectivized bundle  $P(T_X) \rightarrow X$ . For each non singular germ of holomorphic map  $\varphi: (\mathbb{C}, 0) \rightarrow X$ , one can define a projectivized lifting  $\tilde{\varphi}: (\mathbb{C}, 0) \rightarrow P(T_X)$  simply by writing  $\tilde{\varphi}(\zeta) = (\varphi(\zeta), [\varphi'(\zeta)])$ ; here, we think at point of  $P(T_X)$  as pairs  $(x, [v])$ , where  $x \in X$  and  $v \in T_{X, x} \setminus \{0\}$ . Tautologically, the restriction of  $\mathcal{O}_{P(T_X)}(-1)$  to the image  $\tilde{\varphi}((\mathbb{C}, 0))$  of  $\tilde{\varphi}$  is isomorphic to the tangent bundle to  $\tilde{\varphi}((\mathbb{C}, 0))$ . Now, as we saw, for each point  $(x, [v]) \in \mathcal{S}$  there exists a unique minimal disk  $f: \Delta \rightarrow X$  passing through  $x$  and whose tangent space at  $x$  is  $[v]$ . Moreover, by uniqueness, for each  $f(\zeta)$  one has that  $(f(\zeta), [f'(\zeta)]) \in \mathcal{S}$ , i.e.  $\tilde{f}(\Delta) \subset \mathcal{S}$ . The collection of all  $\tilde{f}(\Delta)$ , when  $f$  runs among all minimal disks in  $X$ , thus endows  $\mathcal{S}$  with a holomorphic foliation by curve  $\mathcal{M}$ , whose tangent bundle  $T_{\mathcal{M}}$  is naturally isomorphic to the restriction of  $\mathcal{O}_{P(T_X)}(-1)$  to  $\mathcal{S}$ .

The proof of Theorem 6.1 is now almost immediate.

*Proof of Theorem 6.1.* The idea is to use the characteristic bundle  $\pi: \mathcal{S} \rightarrow X$  together with Remark 3.3.

The (smooth) holomorphic foliation  $\mathcal{M}$  on  $\mathcal{S}$  constructed above is such that the canonical bundle  $K_{\mathcal{M}}$  to this foliation is isomorphic to  $\mathcal{O}_{P(T_X)}(1)|_{\mathcal{S}}$ . By Proposition 6.8,  $K_{\mathcal{M}}$  has negative Kodaira–Iitaka dimension. Thus, by Theorem 1.5, we obtain that  $GG(\mathcal{S}, T_{\mathcal{M}}) = \mathcal{S}$ .

Next, the tangent bundle  $T_{\mathcal{M}} \subset T_{\mathcal{S}} \subset T_{P(T_X)}|_{\mathcal{S}}$  to  $\mathcal{M}$  is transverse to the kernel of the differential  $d\pi$ . In fact given a direction tangent to  $\mathcal{M}$ , it is of the form  $\tilde{f}'$  for some minimal disk  $f$  in  $X$ . Thus, since  $\pi \circ \tilde{f} = f$ , we have

$$d\pi(\tilde{f}') = f' \neq 0.$$

Then, Remark 3.3 tells us that  $\pi(GG(\mathcal{S}, T_{\mathcal{M}})) \subset GG(X)$  and, since we have  $GG(\mathcal{S}, T_{\mathcal{M}}) = \mathcal{S}$ , Theorem 6.1 follows. □

## 7. Final remarks on the Green–Griffiths–Lang conjecture

The final step for the solution of the Green–Griffiths–Lang conjecture using only (invariant) jet bundles requires to show that at least one of the base loci

$$B_k := \bigcap_{m \geq 1} \text{Bs}(\mathcal{O}_{P_k(V)}(m) \otimes \pi_{0,k}^* A^{-1}) \setminus P_k(V)^{\text{sing}} \subset P_k(V)$$

projects down to a proper algebraic subvariety  $Y_k := \pi_{0,k}(B_k)$  in  $X$ . The results in this paper show that, unfortunately, this is hopeless without any further assumption on  $X$  besides that of being of general type.

Nevertheless, one could think of a less demanding property, namely that for all irreducible subvarieties  $Z \subset P_k(V)$  such that  $\pi_{0,k}(Z) = X$ , the restriction  $\mathcal{O}_{P_k(V)}(1)|_Z$  could be big.

This is exactly what is proved in a theorem of Lu and Yau [LY90] for order one jets, for  $X$  a surface of general type with  $c_1(X)^2 - 2c_2(X) > 0$ , achieving thus the proof of the Green–Griffiths–Lang conjecture in this case. But we cannot expect even this less demanding property to be true in general since we have seen at the end of Section 2 that there exist projective manifolds  $X$  of general type such that for each  $k > 0$  there is a smooth submanifold  $X_k \subset P_k(V)$  which projects biholomorphically onto  $X$  via  $\pi_{0,k}$  and such that for all integers  $m > 0$  we have

$$H^0(X_k, (\mathcal{O}_{P_k(V)}(m) \otimes \pi_{0,k}^* A^{-1})|_{X_k}) = 0.$$

Thus, we see that Lu and Yau’s result is sharp in this sense and that  $c_1^2(X) = 2c_2(X)$  is somehow a threshold for projective surfaces of general type as far as the solution to the Green–Griffiths–Lang conjecture using only base loci of sections of  $k$ -jet bundles is concerned.

On the other hand, McQuillan’s celebrated work on surfaces [McQ98] shows that the conjecture is true whenever the second Segre number  $c_1^2 - c_2$  of the surface is positive (this being the case for the compact free quotients of the bidisc thanks to the Hirzebruch proportionality principle, see [Hir58]). Observe that his proof relies upon a combination of the theories of jet differentials and that of foliations: jet differentials of order one (the existence of which is assured by the assumption on the Segre number) produce (multi)foliations such that every entire curve must be tangent to. The algebraic degeneracy is then obtained in this situation as a consequence of deep results about parabolic leaves of holomorphic foliations on surfaces of general type rather than trying to control the Green–Griffiths locus (which, *a posteriori* would not give the desired result, also in view of what is described in these lines).

As far as we know, at present, the only cases where an explicit control of the Green–Griffiths locus is known are given by generic projective hypersurfaces

of high degree: it is shown in [DMR10] that if  $X \subset \mathbb{P}^{n+1}$ ,  $n \geq 2$ , is a generic projective hypersurface of degree  $\deg X \geq 2^{n^5}$ , then  $GG(X)$  is contained in a proper subvariety of  $X$ . This result is further refined in [DT10] where, under the same hypotheses, it is proved that  $GG(X)$  is contained in a proper subvariety of codimension at least two. The reader may consult the survey [DR11] for more details about hyperbolicity of projective hypersurfaces.

**Acknowledgments.** The first-named author would like to warmly thank Nicolas Bergeron, Olivier Biquard, Sébastien Boucksom, Fabrizio Catanese, Gilles Courtois, Jean–Pierre Demailly, Antonio J. Di Scala, Bruno Klingler and Roberto Pignatelli for very interesting discussions about several themes concerning different aspects of this paper.

The second-named author thanks Ziyang Gao, Carlo Gasbarri, Steven Lu, Michael McQuillan and Ngaiming Mok for stimulating discussions. He also thanks CNRS for the opportunity to spend a semester in Montreal at the UMI CNRS-CRM and the hospitality of UQAM-Cirget where part of this work was done.

## References

- [BCGP12] I. BAUER, F. CATANESE, F. GRUNEWALD and R. PIGNATELLI, Quotients of products of curves, new surfaces with  $p_g = 0$  and their fundamental groups. *Amer. J. Math.* **134** (2012), 993–1049. Zbl 1258.14043 MR 2956256
- [Bea00] A. BEAUVILLE, Complex manifolds with split tangent bundle. In *Complex analysis and algebraic geometry*, pages 61–70. de Gruyter, Berlin, 2000. Zbl 1001.32010 MR 1760872
- [Bog77] F. A. BOGOMOLOV, Families of curves on a surface of general type. *Dokl. Akad. Nauk SSSR* **236** (1977), 1041–1044. Zbl 0415.14013 MR 0457450
- [Bro11] D. Brotbek, Differential equations as embedding obstructions and vanishing theorems. *ArXiv e-prints*, November 2011.
- [Bru97] M. BRUNELLA, Feuilletages holomorphes sur les surfaces complexes compactes. *Ann. Sci. École Norm. Sup. (4)* **30** (1997), 569–594. Zbl 0893.32019 MR 1474805
- [Bru04] M. BRUNELLA, *Birational geometry of foliations*. Publicações Matemáticas do IMPA. [IMPA Mathematical Publications]. Instituto de Matemática Pura e Aplicada (IMPA), Rio de Janeiro, 2004. Zbl 1082.32022 MR 2114696
- [CP13] F. CAMPANA and M. PĂUN, Orbifold generic semi-positivity: an application to families of canonically polarized manifolds. *Ann. Inst. Fourier* **65** (2015), 835–861. Zbl 06496595 MR 3449168

[Dem97] J.-P. DEMAILLY, Algebraic criteria for Kobayashi hyperbolic projective varieties and jet differentials. In *Algebraic geometry – Santa Cruz 1995*, volume 62 of *Proc. Sympos. Pure Math.*, pages 285–360. Amer. Math. Soc., Providence, RI, 1997. Zbl 0919.32014 MR 1492539

[Dem11] J.-P. DEMAILLY, Holomorphic Morse inequalities and the Green-Griffiths-Lang conjecture. *Pure Appl. Math. Q.* **7** (Special Issue: In memory of Eckart Viehweg) (2011), 1165–1207. Zbl 1316.32014 MR 2918158

[Des79] M. DESCHAMPS, Courbes de genre géométrique borné sur une surface de type général [d’après F. A. Bogomolov]. In *Séminaire Bourbaki, 30e année (1977/78)*, volume 710 of *Lecture Notes in Math.*, pages Exp. No. 519, pp. 233–247. Springer, Berlin, 1979. Zbl 0412.14013 MR 0554224

[Div08] S. DIVERIO, Differential equations on complex projective hypersurfaces of low dimension. *Compos. Math.* **144** (2008), 920–932. Zbl 1193.32013 MR 2441250

[Div09] S. DIVERIO, Existence of global invariant jet differentials on projective hypersurfaces of high degree. *Math. Ann.* **344** (2009), 293–315. Zbl 1166.32013 MR 2495771

[DMR10] S. DIVERIO, J. MERKER and E. ROUSSEAU, Effective algebraic degeneracy. *Invent. Math.* **180** (2010), 161–223. Zbl 1192.32014 MR 2593279

[DR11] S. DIVERIO and E. ROUSSEAU, *A survey on hyperbolicity of projective hypersurfaces*. Publicações Matemáticas do IMPA. [IMPA Mathematical Publications]. Instituto de Matemática Pura e Aplicada (IMPA), Rio de Janeiro, 2011. [On the title page: A survey on hiperbolicity of projective hypersurfaces]. Zbl 1250.32001 MR 2933164

[DT10] S. DIVERIO and S. TRAPANI, A remark on the codimension of the Green-Griffiths locus of generic projective hypersurfaces of high degree. *J. Reine Angew. Math.* **649** (2010), 55–61. Zbl 1211.32015 MR 2746466

[Fre79] E. FREITAG, Ein Verschwindungssatz für automorphe Formen zur Siegelschen Modulgruppe. *Math. Z.* **165** (1979), 11–18. Zbl 0393.10027 MR 0521517

[Fre90] E. FREITAG, *Hilbert modular forms*. Springer-Verlag, Berlin, 1990. Zbl 0702.11029 MR 1050763

[GG80] M. GREEN and P. GRIFFITHS, Two applications of algebraic geometry to entire holomorphic mappings. In *The Chern Symposium 1979 (Proc. Internat. Sympos., Berkeley, Calif., 1979)*, pages 41–74. Springer, New York, 1980. Zbl 0508.32010 MR 0609557

[Hir58] F. HIRZEBRUCH, Automorphe Formen und der Satz von Riemann-Roch. In *Symposium internacional de topología algebraica International symposium on algebraic topology*, pages 129–144. Universidad Nacional Autónoma de México and UNESCO, Mexico City, 1958. Zbl 0129.29801 MR 0103280

[Kat92] S. KATOK, *Fuchsian groups*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1992. Zbl 0753.30001 MR 1177168

- [Lan86] S. LANG, Hyperbolic and Diophantine analysis. *Bull. Amer. Math. Soc. (N.S.)* **14** (1986), 159–205. Zbl 0602.14019 MR 0828820
- [Lu96] S. SHIN-YI LU, On hyperbolicity and the Green–Griffiths conjecture for surfaces. In *Geometric complex analysis (Hayama, 1995)*, pages 401–408. World Sci. Publ., River Edge, NJ, 1996. Zbl 0941.32024 MR 1453621
- [LY90] S. SHIN-YI LU and S.-T. YAU, Holomorphic curves in surfaces of general type. *Proc. Nat. Acad. Sci. U.S.A.* **87** (1990), 80–82. Zbl 0702.32015 MR 1031949
- [McQ98] M. MCQUILLAN, Diophantine approximations and foliations. *Inst. Hautes Études Sci. Publ. Math.* **87** (1998), 121–174. Zbl 1006.32020 MR 1659270
- [McQ08] M. MCQUILLAN, Canonical models of foliations. *Pure Appl. Math. Q.* **4** (2008), 877–1012. Zbl 1166.14010 MR 2435846
- [McQ14] M. MCQUILLAN, Curves on surfaces of mixed characteristic. Preprint, 2014.
- [Mer10] J. MERKER, Complex projective hypersurfaces of general type: toward a conjecture of Green and Griffiths. *ArXiv e-prints*, May 2010.
- [Mok89] N. MOK, *Metric rigidity theorems on Hermitian locally symmetric manifolds*, volume 6 of *Series in Pure Mathematics*. World Scientific Publishing Co. Inc., Teaneck, NJ, 1989. Zbl 0912.32026 MR 1081948
- [Nar71] R. NARASIMHAN, *Several complex variables*. The University of Chicago Press, Chicago, Ill.-London, 1971. Chicago Lectures in Mathematics. Zbl 0223.32001 MR 0342725
- [PR94] V. PLATONOV and A. RAPINCHUK, *Algebraic groups and number theory*, volume 139 of *Pure and Applied Mathematics*. Academic Press Inc., Boston, MA, 1994. Translated from the 1991 Russian original by Rachel Rowen. Zbl 0841.20046 MR 1278263
- [Rag72] M. S. RAGHUNATHAN, *Discrete subgroups of Lie groups*. Springer-Verlag, New York, 1972. *Ergebnisse der Mathematik und ihrer Grenzgebiete*, Band 68. Zbl 0254.22005 MR 0507234
- [SB95] N. I. SHEPHERD-BARRON, Infinite generation for rings of symmetric tensors. *Math. Res. Lett.* **2** (1995), 125–128. Zbl 0839.53048 MR 1324696
- [Sha78] I. H. SHABEL, A class of algebraic surfaces of general type constructed from quaternion algebras. *Pacific J. Math.* **76** (1978), 221–245. Zbl 0422.14022 MR 0572981
- [Shi94] G. SHIMURA, *Introduction to the arithmetic theory of automorphic functions*, volume 11 of *Publications of the Mathematical Society of Japan*. Princeton University Press, Princeton, NJ, 1994. Reprint of the 1971 original, Kanô Memorial Lectures, 1. Zbl 0872.11023 MR 1291394
- [SY97] Y.-T. SIU and S.-K. YEUNG, Defects for ample divisors of abelian varieties, Schwarz lemma, and hyperbolic hypersurfaces of low degrees. *Amer. J. Math.* **119** (1997), 1139–1172. Zbl 0947.32012 MR 1473072
- [SZ00] B. SHIFFMAN and M. ZAIDENBERG, Two classes of hyperbolic surfaces in  $\mathbf{P}^3$ . *Internat. J. Math.* **11** (2000), 65–101. Zbl 1070.14518 MR 1757892

[UY10] E. ULLMO and A. YAFAEV, Points rationnels des variétés de Shimura: un principe du “tout ou rien”. *Math. Ann.* **348** (2010), 689–705. Zbl 1264.11052 MR 2677900

[Yau93] S.-T. YAU, A splitting theorem and an algebraic geometric characterization of locally Hermitian symmetric spaces. *Comm. Anal. Geom.* **1** (1993), 473–486. Zbl 0842.53035 MR 1266476

(Reçu le 11 septembre 2014)

Simone DIVERIO, CNRS, Institut de Mathématiques de Jussieu–Paris Rive Gauche, UMR7586, Sorbonne Universités, UPMC Univ Paris 06, Univ Paris Diderot, Sorbonne Paris Cité, 75005, Paris, France

*e-mail:* simone.diverio@imj-prg.fr

Erwan Rousseau, Aix Marseille Université, CNRS, Centrale Marseille, I2M, UMR 7373, 13453 Marseille, France

*e-mail:* erwan.rousseau@univ-amu.fr