

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 62 (2016)
Heft: 1-2

Artikel: Note on the distortion of $(2,q)$ -torus knots
Autor: Studer, Luca
DOI: <https://doi.org/10.5169/seals-685357>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 29.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Note on the distortion of $(2, q)$ -torus knots

LUCA STUDER

Abstract. We show that the distortion of the $(2, q)$ -torus knot is not bounded linearly from below.

Mathematics Subject Classification (2010). Primary: 57M25.

Keywords. Distortion, torus knots.

1. Introduction

The notion of distortion was introduced by Gromov [GPL]. If γ is a rectifiable simple closed curve in \mathbf{R}^3 , then its distortion δ is defined as

$$\delta(\gamma) = \sup_{v, w \in \gamma} \frac{d_\gamma(v, w)}{|v - w|},$$

where $d_\gamma(v, w)$ denotes the length of the shortest arc connecting v and w in γ and $|\cdot|$ denotes the euclidean norm on \mathbf{R}^3 . For a knot K , its distortion $\delta(K)$ is defined as the infimum of $\delta(\gamma)$ over all rectifiable curves γ in the isotopy class K . Gromov [Gro] asked in 1983 if every knot K has distortion $\delta(K) \leq 100$. The question was open for almost three decades until Pardon gave a negative answer. His work [Par] presents a lower bound for the distortion of simple closed curves on closed PL embedded surfaces with positive genus. Pardon showed that the minimal intersection number of such a curve with essential discs of the corresponding surface bounds the distortion of the curve from below. In particular for the (p, q) -torus knot he showed that $\delta(T_{p, q}) \geq \min(p, q)/160$. By considering a standard embedding of $T_{p, p+1}$ into a torus of revolution one obtains $\delta(T_{p, p+1}) \leq \text{const} \cdot p$, hence for $q = p + 1$ Pardon's result is sharp up to a constant.

An alternative proof for the existence of families with unbounded distortion was given by Gromov and Guth [GG]. In both works the answer to Gromov's question was obtained by estimating the conformal length, which is up to a

constant a lower bound for the distortion of rectifiable closed curves. However, the conformal length is in general not a good estimate for the distortion. For example, one finds easily an embedding of the $(2, q)$ -torus knot with conformal length ≤ 100 and distortion $\geq q$ by looking at standard embeddings into a torus of revolution with suitable dimensions. In particular, neither Pardon's nor Gromov and Guth's arguments yield lower bounds for $\delta(T_{2,q})$. While Pardon conjectures that $\lim_{q \rightarrow \infty} \delta(T_{2,q}) = \infty$ and that there are to his knowledge no known embeddings of $T_{2,q}$ with sublinear distortion [Par, p. 638], Gromov and Guth [GG, p. 2588] write that the distortion of $T_{2,q}$ appears to be approximately q . In this article we show that the growth rate of $\delta(T_{2,q})$ is in fact sublinear in q .

Theorem. *Let $q \geq 50$. Then $\delta(T_{2,q}) \leq 7q/\log q$. In particular the distortion of the $(2, q)$ -torus knot is not bounded linearly from below.*

With the same technique as used in this article and somewhat more effort one can give an embedding γ_q of $T_{2,q}$ with $\delta(\gamma_q) \sim \frac{\pi}{2} \frac{q}{\log q}$. Moreover, a more technical proof yields that this asymptotical upper bound for $\delta(T_{2,q})$ is sharp for those embeddings of $T_{2,q}$ that project orthogonally onto a standard knot diagram. This leads to the following question.

Question. *Is $\delta(T_{2,q})$ up to a constant asymptotically equal to $q/\log q$? And if yes, is the constant equal to $\pi/2$?*

2. Proof of the Theorem

In order to prove the Theorem we need to give for every odd integer $q \geq 50$ an embedding γ of the $(2, q)$ -torus knot with distortion smaller or equal to $7q/\log q$. The idea is to use a logarithmic spiral. Let S be a logarithmic spiral of unit length starting at its center $0 \in \mathbf{R}^3$ and ending at some $u \in \mathbf{R}^3$. An elementary calculation shows that its distortion is equal to $1/|u|$. For another path $\alpha \subset \mathbf{R}^3$ of unit length and diameter $\leq 2|u|$ with endpoints $\{v, w\} = \partial\alpha$ we get

$$\delta(\alpha) \geq \frac{d_\alpha(v, w)}{|v - w|} = \frac{1}{|v - w|} \geq \frac{1}{2|u|} = \frac{\delta(S)}{2}.$$

Hence up to at most a factor 2 the logarithmic spiral has the smallest distortion among all paths for a prescribed pathlength-pathdiameter-ratio. It seems therefore natural to pack the q windings of the $(2, q)$ -torus knot into a logarithmic spiral in order to minimize distortion.




FIGURE 1
The embedding γ for $q = 7$

We define the embedding γ as the union of a segment of the logarithmic spiral S with slope $k = \log(q)/2\pi q$ and a piecewise linear part L , see Figure 1. The segment of the logarithmic spiral S is contained in the vertical (x, z) plane and parametrized by

$$\varphi : [0, \pi q] \rightarrow \mathbf{R}^2, \quad \varphi(s) = e^{ks} \cdot \begin{pmatrix} \cos(s) \\ \sin(s) \end{pmatrix},$$

see Figures 1 and 2. The segment of the piecewise linear part L is in the horizontal (x, y) plane, see Figures 1 and 3. Note that

$$|\varphi(\pi q)| = e^{k\pi q} = \sqrt{q} \quad \text{and} \quad |\varphi(0)| = 1,$$

hence the lengths defining L in Figure 3 are chosen such that the union γ of S and L is the simple closed curve illustrated in Figure 1. The linear segments L_1 and L_2 indicated in Figure 3 are named because of their special role in the following computations.

To see that the obtained curve is an embedded $(2, q)$ -torus knot, we perturb γ , see Figure 4. This simple closed curve is ambient isotopic in \mathbf{R}^3 to γ and if we project it onto the (x, y) plane, we see a well known diagram of the $(2, q)$ -torus knot, see Figure 5.

We now estimate the distortion of γ . One has to show that

$$\frac{d_\gamma(v, w)}{|v - w|} \leq \frac{7q}{\log q}$$




FIGURE 2
The logarithmic spiral S in the (x, z) plane




FIGURE 3
The linear part L in the (x, y) plane

for all pairs of points $v, w \in \gamma$. An easy computation shows that

$$\frac{1}{k} \cdot \sqrt{2k^2 + 1} = \frac{2\pi q}{\log q} \cdot \sqrt{2(\log q/2\pi q)^2 + 1} \leq \frac{7q}{\log q}$$

for all integers $q \geq 2$. Therefore, it suffices to show that

$$\frac{d_\gamma(v, w)}{|v - w|} \leq \frac{\sqrt{2k^2 + 1}}{k}.$$

In order to do this, we distinguish four cases.

Case I: $v, w \in S$. Let $0 \leq s \leq t \leq \pi q$, $v = \varphi(s)$, $w = \varphi(t)$. From

$$|\varphi'(r)| = \left| \begin{pmatrix} \cos(r) & -\sin(r) \\ \sin(r) & \cos(r) \end{pmatrix} \begin{pmatrix} ke^{kr} \\ e^{kr} \end{pmatrix} \right| = \left| \begin{pmatrix} ke^{kr} \\ e^{kr} \end{pmatrix} \right| = \sqrt{k^2 + 1} \cdot e^{kr},$$

we get




FIGURE 4
Perturbation of γ




FIGURE 5
Projection onto the (x, y) plane

$$\begin{aligned}
 d_\gamma(v, w) &\leq d_S(v, w) \\
 &= \int_s^t |\varphi'(r)| dr \\
 &= \sqrt{k^2 + 1} \int_s^t e^{kr} dr \\
 &= \frac{\sqrt{k^2 + 1}}{k} \cdot (e^{kt} - e^{ks}) \\
 &= \frac{\sqrt{k^2 + 1}}{k} \cdot (|\varphi(t)| - |\varphi(s)|) \\
 &= \frac{\sqrt{k^2 + 1}}{k} \cdot (|w| - |v|).
 \end{aligned}$$

Since $|w - v| \geq |w| - |v|$, we conclude that




FIGURE 6
Visualization of Case 2

$$\frac{d_\gamma(v, w)}{|v - w|} \leq \frac{\sqrt{k^2 + 1}}{k} \cdot \frac{(|w| - |v|)}{(|w| - |v|)} = \frac{\sqrt{k^2 + 1}}{k}.$$

Case 2: $v \in L_1 \cup L_2$, $w \in S$. We consider the case where $v \in L_1$. The idea is to find the maximum of $d_\gamma(v, w)/|v - w|$ for fixed w and varying v . Let $t = |v - \varphi(0)|$, $a = |\varphi(0) - w|$, and $b = d_S(\varphi(0), w)$, see Figure 6. Note that $|v - w| = \sqrt{t^2 + a^2}$ and $d_\gamma(v, \varphi(0)) = |v - \varphi(0)| = t$. We get

$$\frac{d_\gamma(v, w)}{|v - w|} \leq \frac{d_\gamma(v, \varphi(0)) + d_S(\varphi(0), w)}{|v - w|} = \frac{t + b}{\sqrt{t^2 + a^2}} =: f(t).$$

Deriving f with respect to t yields a unique critical point at $t = a^2/b$:

$$0 = f'(t) = \frac{a^2 - bt}{(a^2 + t^2)^{3/2}} \iff t = a^2/b.$$

Since a^2/b is the only critical point, $f(\infty) = 1 \leq b/a = f(0)$ and

$$f(0) = \frac{b}{a} \leq \frac{\sqrt{a^2 + b^2}}{a} = \frac{\frac{a^2}{b} + b}{\sqrt{(\frac{a^2}{b})^2 + a^2}} = f(a^2/b),$$

a^2/b must be a global maximum. Consequently we get

$$\begin{aligned} \frac{d_\gamma(v, w)}{|v - w|} &\leq \frac{\sqrt{a^2 + b^2}}{a} \\ &= \sqrt{1 + \left(\frac{b}{a}\right)^2} \\ &= \sqrt{1 + \left(\frac{d_S(\varphi(0), w)}{|\varphi(0) - w|}\right)^2} \end{aligned}$$

$$\begin{aligned} &\stackrel{\text{Case 1}}{\leq} \sqrt{1 + \left(\frac{\sqrt{k^2+1}}{k}\right)^2} \\ &= \frac{\sqrt{2k^2+1}}{k}. \end{aligned}$$

In the case where $v \in L_2$, we make the estimate with the path that connects v with w through $\varphi(\pi q)$. It works exactly the same and yields the same estimate.

Case 3: $v, w \in L$. Consider Figure 3 and note that all pairs of points $v, w \in L$ of euclidean distance < 1 are either on the same linear segment or on neighboring linear segments of L . It is easy to see that such pairs of points cannot cause distortion $> \sqrt{2}$. For the pairs of points $v, w \in L$ of euclidean distance ≥ 1 we get

$$\frac{d_Y(v, w)}{|v - w|} \leq d_L(\varphi(0), \varphi(\pi q)) = 11\sqrt{q} + 1.$$

A direct calculation shows that

$$11\sqrt{q} + 1 \leq \frac{2\pi q}{\log q} = \frac{1}{k}$$

for $q \geq 50$.

Case 4: $v \in L \setminus (L_1 \cup L_2), w \in S$. Note that for these pairs of points we have $|v - w| \geq |w|$. We estimate $d_Y(v, w)$ using results of Cases 1 and 3:

$$\begin{aligned} d_Y(v, w) &\leq d_L(v, \varphi(0)) + d_S(\varphi(0), w) \\ &\leq \frac{1}{k} + \frac{\sqrt{k^2+1}}{k} \cdot (|w| - 1) \\ &\leq \frac{\sqrt{k^2+1}}{k} \cdot |w|. \end{aligned}$$

We conclude that

$$\frac{d_Y(v, w)}{|v - w|} \leq \frac{\frac{\sqrt{k^2+1}}{k} \cdot |w|}{|w|} = \frac{\sqrt{k^2+1}}{k},$$

which finishes the proof.

Acknowledgment. First and foremost, I would like to thank Sebastian Baader and Pierre Dehornoy for their inspiring introduction to the topic. I would like to thank John Pardon for useful comments on the presented example and Peter Feller, Filip Misev, Johannes Josi and Livio Liehti for many mathematical discussions. I also thank Paul Frischknecht for the pictures.

References

- [GPL] M. GROMOV, J. LAFONTAINE and P. PANSU, *Structures métriques pour les variétés riemanniennes*. Cedic/Fernand Nathan, Paris, 1981. Zbl 0509.53034 MR 0682063
- [GG] M. GROMOV and L. GUTH, Generalizations of the Kolmogorov-Barzdin embedding estimates. *Duke Math. J.* **161** (2012), 2549–2603. Zbl 1261.53041 MR 2988903
- [Gro] M. GROMOV, Filling Riemannian manifolds. *J. Differential Geom.* **18** (1983), 1–147. Zbl 0515.53037 MR 0697984
- [Par] J. PARDON, On the distortion of knots on embedded surfaces. *Ann. of Math. (2)* **174** (2011), 637–646. Zbl 1227.57013 MR 2811613

(Reçu le 27 mars 2015)

Luca STUDER, Institute of Mathematics, University of Bern, Alpeneggstrasse 22,
3012 Bern, Switzerland

e-mail: luca.studer@math.unibe.ch