

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 63 (2017)
Heft: 1-2

Artikel: The Cartan-Hadamard Theorem for metric spaces with local geodesic bicomblings
Autor: Miesch, Benjamin
DOI: <https://doi.org/10.5169/seals-760290>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The Cartan–Hadamard Theorem for metric spaces with local geodesic bicomblings

Benjamin MIESCH

Abstract. We prove the Cartan–Hadamard Theorem for spaces which are not necessarily uniquely geodesic but locally possess a suitable selection of geodesics, a so-called convex geodesic bicombling.

Furthermore, we deduce a local-to-global theorem for injective (or hyperconvex) metric spaces, saying that under certain conditions a complete, simply-connected, locally injective metric space is injective. A related result for absolute 1-Lipschitz retracts follows.

Mathematics Subject Classification (2010). Primary: 53C22, 53C23; Secondary: 51F99, 54E35.

Keywords. Local-to-global, geodesic bicomblings, injective metric spaces, hyperconvex metric spaces.

1. Introduction

Local-to-global principles are spread all-around in mathematics. The classical Cartan–Hadamard Theorem from Riemannian geometry was generalized by W. Ballmann [Bal] for metric spaces with non-positive curvature, and by S. Alexander and R. Bishop [AB] for locally convex metric spaces, i.e., for spaces that locally satisfy the Busemann property, meaning that $d \circ (\gamma_1, \gamma_2)$ is convex for all constant speed geodesics γ_1, γ_2 . As a normed vector space satisfies the Busemann property if and only if its norm is strictly convex, this property is not preserved under limit processes. This motivates to look at an even weaker notion of non-positive curvature, where we only request convexity for a certain choice of geodesics, compare [Kle, Section 10].

We use the convention of [ABr, Lan] to call such a collection of paths a bicombling, a term originally coined by W. Thurston in the context of geometric group theory. The following definition is for instance satisfied by the linear segments $(1-t)x + ty$ in a normed vector space. In a metric space (X, d) ,

a *geodesic bicombing* is a selection of a geodesic between each pair of points. This is a map $\sigma: X \times X \times [0, 1] \rightarrow X$ such that, for all $x, y \in X$, the path $\sigma_{xy} := \sigma(x, y, \cdot)$ is a geodesic from x to y , i.e. $\sigma_{xy}(0) = x$, $\sigma_{xy}(1) = y$ and $d(\sigma_{xy}(s), \sigma_{xy}(t)) = |s - t|d(x, y)$ for all $s, t \in [0, 1]$. Moreover, we assume that this choice is consistent in the sense that $\sigma_{pq}([0, 1]) \subset \sigma_{xy}([0, 1])$ for all $p, q \in \sigma_{xy}([0, 1])$ with $d(x, p) \leq d(x, q)$. A geodesic bicombing σ is called *convex* if the function $t \mapsto d(\sigma_{xy}(t), \sigma_{\bar{x}\bar{y}}(t))$ is convex for all $x, y, \bar{x}, \bar{y} \in X$. Furthermore, we say that σ is *reversible* if $\sigma_{yx}([0, 1]) = \sigma_{xy}([0, 1])$ for all $x, y \in X$. A metric space admits a *local geodesic bicombing*, if such a selection exists in a neighborhood $U(x, r_x)$ of each point $x \in X$, see Section 2 for the exact definition.

Metric spaces with a geodesic bicombing resemble hyperbolic spaces after U. Kohlenbach [Koh, KL], which specify W -convex metric spaces considered by W. Takahashi [Tak] and S. Itoh [Ito]. Geodesic bicomblings were recently studied by D. Descombes and U. Lang in [Des, DL1, DL2] and also by G. Basso in [Bas], where they show that several results for CAT(0) and Busemann spaces carry over to spaces with convex geodesic bicomblings. Here we will contribute to these studies by proving the following Cartan–Hadamard Theorem.

Theorem 1.1. *Let X be a complete, simply-connected metric space with a convex local geodesic bicombing σ . Then the induced length metric on X admits a unique convex geodesic bicombing $\tilde{\sigma}$ which is consistent with σ . As a consequence, X is contractible. Moreover, if the local geodesic bicombing σ is reversible, then $\tilde{\sigma}$ is reversible as well.*

As we show in a subsequent paper joint with G. Basso [BM], Theorem 1.1 leads to a uniqueness result for convex geodesic bicomblings on convex subsets of certain Banach spaces.

Important examples of spaces with convex geodesic bicomblings are given by injective metric spaces, which include the real line, \mathbb{R} -trees and $l_\infty(I)$ for any index set I . Recall that every metric space X possesses an injective hull, i.e., a smallest injective metric space into which X embeds [Isb]. Injective metric spaces play a crucial role in the theory of mapping extensions [AP] and fixed point theory [Sin, Soa], see also [EK] and the references therein.

A metric space X is *injective* if for all metric spaces A, B with $A \subset B$ and every 1-Lipschitz map $f: A \rightarrow X$, there is a 1-Lipschitz extension $\bar{f}: B \rightarrow X$, i.e. $\bar{f}|_A = f$. In fact, D. Descombes and U. Lang show in their work that every proper, injective metric space of finite combinatorial dimension admits a (unique) convex geodesic bicombing [DL1, Theorem 1.2]. Such spaces occur, for instance, as injective hulls of hyperbolic groups [Lan, Theorem 1.4] and therefore, every

hyperbolic group acts properly and cocompactly by isometries on a space with a convex geodesic bicombing [DL1, Theorem 1.3].

Recall that injective metric spaces are complete, geodesic and contractible. Now, knowing that under the above conditions injective metric spaces possess a convex geodesic bicombing, we deduce the following local-to-global theorem for injective metric spaces.

Theorem 1.2. *Let X be a complete, locally compact, simply-connected, locally injective length space with locally finite combinatorial dimension. Then X is an injective metric space.*

It is well known that injective metric spaces are the same as absolute 1-Lipschitz retracts. For Lipschitz retracts, the weaker notion of absolute Lipschitz uniform neighborhood retracts is common, see Section 4 for more details. Absolute 1-Lipschitz uniform neighborhood retracts are locally injective but the converse is not true as we will see in Example 4.2. In fact, it turns out that the following holds.

Theorem 1.3. *Let X be a locally compact absolute 1-Lipschitz uniform neighborhood retract with locally finite combinatorial dimension. Then X is an absolute 1-Lipschitz retract.*

This paper is organized as follows. We start Section 2 by studying spaces with local geodesic bicomblings, establish an appropriate exponential map and finally prove Theorem 1.1. In Section 3, we first show that every uniformly locally injective metric space with a reversible, convex geodesic bicombing is injective. Afterwards, we describe how to construct a reversible, convex local geodesic bicombing on locally injective metric spaces, which extends to a convex geodesic bicombing by Theorem 1.1. Thereby we establish Theorem 1.2. Finally in Section 4, we then investigate absolute 1-Lipschitz neighborhood retracts and prove Theorem 1.3.

2. Local geodesic bicomblings

Let us first fix some notation. In a metric space X , we denote by

$$U(x, r) := \{y \in X : d(x, y) < r\}$$

the open ball of radius r around $x \in X$ and by

$$B(x, r) := \{y \in X : d(x, y) \leq r\}$$

the closed one.

Let X be a metric space and $\gamma: [0, 1] \rightarrow X$ a continuous curve. The *length* of γ is given by

$$L(\gamma) := \sup \left\{ \sum_{k=1}^n d(\gamma(t_{k-1}), \gamma(t_k)) : 0 = t_0 < \dots < t_n = 1 \right\}.$$

Then

$$\bar{d}(x, y) := \inf \{ L(\gamma) : \gamma: [0, 1] \rightarrow X, \gamma(0) = x, \gamma(1) = y \}$$

defines a metric on X , called the *induced length metric*. If we have $d = \bar{d}$, we say that (X, d) is a *length space*.

For a metric space X , let $\mathcal{G}(X) := \{c: [0, 1] \rightarrow X\}$ be the set of all geodesics in X , i.e. continuous maps $c: [0, 1] \rightarrow X$ with $d(c(s), c(t)) = |s-t| \cdot d(c(0), c(1))$ for all $s, t \in [0, 1]$. Note that geodesics need not be parametrised by arc-length. We equip $\mathcal{G}(X)$ with the metric

$$D(c, c') := \sup_{t \in [0, 1]} d(c(t), c'(t)).$$

Let $c \in \mathcal{G}(X)$ and $0 \leq a \leq b \leq 1$, then we denote by $c_{[a, b]}$ the reparametrized geodesic given by $c_{[a, b]}: [0, 1] \rightarrow X$ with $c_{[a, b]}(t) := c((1-t)a + tb)$.

Definition 2.1. A *local geodesic bicombing* on a metric space X is a local selection of geodesics, i.e., a map $\sigma: U \times [0, 1] \rightarrow X$, $(y, z, t) \mapsto \sigma_{yz}(t)$, with $U \subset X \times X$ open and the following properties:

- (i) For all $x \in X$, there is some $r_x > 0$ such that, for all $y, z \in U(x, r_x)$, there is a geodesic $\sigma_{yz}: [0, 1] \rightarrow U(x, r_x)$ from y to z , and

$$U = \{(y, z) \in X \times X : y, z \in U(x, r_x) \text{ for some } x\}.$$

- (ii) The selection is consistent with taking subsegments of geodesics, i.e.,

$$\sigma_{\sigma_{yz}(s_1)\sigma_{yz}(s_2)}(t) = \sigma_{yz}((1-t)s_1 + ts_2)$$

for $(y, z) \in U$, $0 \leq s_1 \leq s_2 \leq 1$ and $t \in [0, 1]$.

We call a local geodesic bicombing σ *convex* if it is locally convex, i.e. for $y, z, y', z' \in U(x, r_x)$, it holds that

$$t \mapsto d(\sigma_{yz}(t), \sigma_{y'z'}(t))$$

is a convex function. Furthermore, σ is *reversible* if

$$\sigma_{zy}(t) = \sigma_{yz}(1-t)$$

for all $(y, z) \in U$ and $t \in [0, 1]$.

Remark. Observe that, by property (ii), a local geodesic bicombing is convex if and only if

$$d(\sigma_{yz}(t), \sigma_{y'z'}(t)) \leq (1-t)d(y, y') + td(z, z')$$

for all $y, z, y', z' \in U(x, r_x)$ and $t \in [0, 1]$.

A (local) geodesic $c: [0, 1] \rightarrow X$ is *consistent* with the local geodesic bicombing σ if

$$c_{[a,b]}(t) = \sigma_{c(a)c(b)}(t)$$

for all $0 \leq a \leq b \leq 1$ with $(c(a), c(b)) \in U$.

To prove Theorem 1.1, we roughly follow the structure of Chapter II.4 in [BH]. Adapting the methods of S. Alexander and R. Bishop [AB], we can prove the following key lemma.

Lemma 2.2. *Let X be a complete metric space with a convex local geodesic bicombing σ and let c be a local geodesic from x to y which is consistent with σ . Then, there is some $\epsilon > 0$ such that, for all $\bar{x}, \bar{y} \in X$ with $d(x, \bar{x}), d(y, \bar{y}) < \epsilon$, there is a unique local geodesic \bar{c} from \bar{x} to \bar{y} with $D(c, \bar{c}) < \epsilon$ which is consistent with σ . Moreover, we have*

$$L(\bar{c}) \leq L(c) + d(x, \bar{x}) + d(y, \bar{y})$$

and if \tilde{c} is another consistent geodesic from \tilde{x} to \tilde{y} with $D(c, \tilde{c}) < \epsilon$, then

$$t \mapsto d(\tilde{c}(t), \bar{c}(t))$$

is convex.

Proof. Let $\epsilon > 0$ be such that $\sigma|_{U(c(t), 2\epsilon) \times U(c(t), 2\epsilon) \times [0, 1]}$ is a convex geodesic bicombing for all $t \in [0, 1]$. Now, let $P(A)$ be the following statement:

$P(A)$: For all $a, b \in [0, 1]$ with $0 \leq b - a \leq A$ and for all $p, q \in X$ with $d(c(a), p), d(c(b), q) < \epsilon$, there is a unique local geodesic $\bar{c}_{pq}: [0, 1] \rightarrow X$ from p to q with $D(c_{[a,b]}, \bar{c}_{pq}) < \epsilon$ which is consistent with σ . Moreover, for all such local geodesics the map $t \mapsto d(\bar{c}_{pq}(t), \bar{c}_{p'q'}(t))$ is convex.

By our assumption, $P(\frac{\epsilon}{L(c)})$ holds. Therefore, let us show $P(A) \Rightarrow P(\frac{3A}{2})$.

Given $a, b \in [0, 1]$ with $0 \leq b - a \leq \frac{3A}{2}$, define $p_0 := c(\frac{2}{3}a + \frac{1}{3}b)$ and $q_0 := c(\frac{1}{3}a + \frac{2}{3}b)$. Then, by $P(A)$, there are consistent local geodesics c_1 from p to q_0 and c'_1 from p_0 to q . Inductively, we set $p_n := c_n(\frac{1}{2})$ and $q_n := c'_n(\frac{1}{2})$, where c_n is a consistent local geodesic from p to q_{n-1} and c'_n from p_{n-1} to q . Observe that, by convexity of the c_n, c'_n , we have $d(p_{n-1}, p_n), d(q_{n-1}, q_n) < \frac{\epsilon}{2^n}$ and hence the sequences $(p_n)_n$ and $(q_n)_n$ converge to some p_∞ and q_∞ ,

respectively, and we have $d(p_\infty, p_0), d(q_\infty, q_0) < \epsilon$. Furthermore, by convexity, the c_n, c'_n converge to the consistent local geodesics c_∞ from p to q_∞ and c'_∞ from p_∞ to q , which coincide between $p_\infty = c_\infty(\frac{1}{2})$ and $q_\infty = c'_\infty(\frac{1}{2})$. Hence, they define a new local geodesic c_{pq} from p to q which is consistent with σ and $p_\infty = c_{pq}(\frac{1}{3}), q_\infty = c_{pq}(\frac{2}{3})$.

Now, given two local geodesics c_{pq} and $c_{p'q'}$ with $D(c_{[a,b]}, c_{pq}) < \epsilon$ and $D(c_{[a,b]}, c_{p'q'}) < \epsilon$, set $s := d(p, p'), t := d(q, q'), s' := d(c_{pq}(\frac{1}{3}), c_{p'q'}(\frac{1}{3}))$ and $t' := d(c_{pq}(\frac{2}{3}), c_{p'q'}(\frac{2}{3}))$. Then we have $s' \leq \frac{s+t'}{2}, t' \leq \frac{s'+t}{2}$ and therefore $s' \leq \frac{s}{2} + \frac{s'}{4} + \frac{t}{4}$, i.e. $s' \leq \frac{2s+t}{3}$ and similarly $t' \leq \frac{s+2t}{3}$ follows. Hence, we get convexity of $t \mapsto d(c_{pq}(t), c_{p'q'}(t))$ and therefore also uniqueness follows.

It remains to prove that $L(\tilde{c}) \leq L(c) + d(x, \bar{x}) + d(y, \bar{y})$. Let \tilde{c} be the unique consistent local geodesic from x to \bar{y} with $D(c, \tilde{c}) < \epsilon$. For t small enough we have

$$\begin{aligned} tL(\tilde{c}) &= d(\tilde{c}(0), \tilde{c}(t)) = d(c(0), \tilde{c}(t)) \\ &\leq d(c(0), c(t)) + d(c(t), \tilde{c}(t)) \leq tL(c) + td(c(1), \tilde{c}(1)), \end{aligned}$$

i.e., $L(\tilde{c}) \leq L(c) + d(y, \bar{y})$ and similarly $L(\bar{c}) \leq L(\tilde{c}) + d(x, \bar{x})$. □

Definition 2.3. Let X be a metric space with a local geodesic bicombing σ . For some fixed $x_0 \in X$, we define

$$\tilde{X}_{x_0} := \{c: [0, 1] \rightarrow X \text{ local geodesic with } c(0) = x_0, \text{ consistent with } \sigma\}.$$

We equip \tilde{X}_{x_0} with the metric $D(c, c') = \sup_{t \in [0, 1]} d(c(t), c'(t))$ and define the map

$$\exp: \tilde{X}_{x_0} \rightarrow X, c \mapsto c(1).$$

If X is complete, this map has the following properties.

Lemma 2.4. *Let X be a complete metric space with a convex local geodesic bicombing σ . Then the following holds:*

- (i) *The map $\exp: \tilde{X}_{x_0} \rightarrow X$ is locally an isometry. Hence σ naturally induces a convex local geodesic bicombing $\tilde{\sigma}$ on \tilde{X}_{x_0} .*
- (ii) *\tilde{X}_{x_0} is contractible.*
- (iii) *For each $\tilde{x} \in \tilde{X}_{x_0}$, there is a unique local geodesic from \tilde{x}_0 to \tilde{x} which is consistent with $\tilde{\sigma}$, where \tilde{x}_0 is the constant path $\tilde{x}_0(t) = x_0$.*
- (iv) *\tilde{X}_{x_0} is complete.*

Proof. By Lemma 2.2, for every $c \in \tilde{X}_{x_0}$, there is some $\epsilon > 0$ such that the map $\exp|_{U(c, \epsilon)}: U(c, \epsilon) \rightarrow U(c(1), \epsilon)$ is an isometry. Hence, σ naturally induces a convex local geodesic bicombing $\tilde{\sigma}$ on \tilde{X}_{x_0} .

Consider the map $r: \tilde{X}_{x_0} \times [0, 1] \rightarrow \tilde{X}_{x_0}, (c, s) \mapsto (r_s(c): t \mapsto c(st))$. This defines a retraction of \tilde{X}_{x_0} to \tilde{x}_0 .

A continuous path $\tilde{c}: [0, 1] \rightarrow \tilde{X}_{x_0}$ is a local geodesic in \tilde{X}_{x_0} which is consistent with $\tilde{\sigma}$ if and only if $\exp \circ \tilde{c}$ is a local geodesic in X which is consistent with σ . Therefore, for any $c \in \tilde{X}_{x_0}$, the map $s \mapsto r_s(c)$ is the unique local geodesic from \tilde{x}_0 to c .

Finally, if $(c_n)_n$ is a Cauchy sequence in \tilde{X} , by completeness of X , for every $t \in [0, 1]$, the sequences $(c_n(t))_n$ converge in X , to $c(t)$ say. Locally, i.e., inside $U(c(t), r_c(t))$, the subsegment $c|_{[t-\epsilon, t+\epsilon]}$ is the limit of the consistent geodesics $(c_n|_{[t-\epsilon, t+\epsilon]})_n$ and hence c is consistent with σ by the convexity of the local geodesic bicombing. \square

The following criterion will ensure that \exp is a covering map.

Lemma 2.5. *Let $p: \tilde{X} \rightarrow X$ be a map of length spaces such that*

- (i) X is connected,
- (ii) p is a local homeomorphism,
- (iii) for all rectifiable curves $\tilde{c}: [0, 1] \rightarrow \tilde{X}$, we have $L(\tilde{c}) \leq L(p \circ \tilde{c})$,
- (iv) X has a convex local geodesic bicombing σ , and
- (v) \tilde{X} is complete.

Then p is a covering map.

Proof. The proof of Proposition I.3.28 in [BH] also works in our setting. In the second step, take $U = U(x, r_x)$ and define the maps $s_{\tilde{x}}: U(x, r_x) \rightarrow \tilde{X}$ by $s_{\tilde{x}}(y) = \tilde{\sigma}_{xy}(1)$, where $\tilde{\sigma}_{xy}$ is the unique lift of σ_{xy} with $\tilde{\sigma}_{xy}(0) = \tilde{x}$. \square

Remark. For a local isometry p , conditions (ii) and (iii) are satisfied.

Corollary 2.6. *Let (X, d) be a complete, connected metric space with a convex local geodesic bicombing σ . Then $\exp: \tilde{X}_{x_0} \rightarrow X$ is a universal covering map.*

Proof. Consider the induced length metrics \bar{d} and \bar{D} on X and \tilde{X}_{x_0} . Since (X, d) locally is a length space, the metrics d and D locally coincide with \bar{d} and \bar{D} , respectively. Hence p still is a local isometry with respect to the length metrics and σ is a convex local geodesic bicombing. Thus Lemma 2.5 applies. \square

Proof of Theorem 1.1. First, we show that, for all $x, y \in X$, there is a unique consistent local geodesic from x to y . Since X is simply-connected, the covering

map $\exp: \tilde{X}_x \rightarrow X$ is a homeomorphism which is a local isometry and by Lemma 2.4, there is a unique consistent local geodesic $\tilde{\sigma}_{xy}$ from x to y .

Next, we prove that $\tilde{\sigma}_{xy}$ is a geodesic. To do so, it is enough to show that, for every curve $\gamma: [0, 1] \rightarrow X$ and every $t \in [0, 1]$, we have $L(\tilde{\sigma}_{\gamma(0)\gamma(t)}) \leq L(\gamma|_{[0,t]})$. Let

$$A := \{s \in [0, 1] : \forall t \in [0, s] \text{ we have } L(\tilde{\sigma}_{\gamma(0)\gamma(t)}) \leq L(\gamma|_{[0,t]})\}.$$

Clearly, A is non-empty and closed. To prove that A is open, consider $s \in A$. For $\delta > 0$ small enough, by Lemma 2.2, we have

$$\begin{aligned} L(\tilde{\sigma}_{\gamma(0)\gamma(s+\delta)}) &\leq L(\tilde{\sigma}_{\gamma(0)\gamma(s)}) + d(\gamma(s), \gamma(s+\delta)) \\ &\leq L(\gamma|_{[0,s]}) + L(\gamma|_{[s,s+\delta]}) = L(\gamma|_{[0,s+\delta]}). \end{aligned}$$

Hence, $A = [0, 1]$ as desired.

Finally, we show that $t \mapsto d(\tilde{\sigma}_{xy}(t), \tilde{\sigma}_{\bar{x}\bar{y}}(t))$ is convex. By Lemma 2.2, there is a sequence $0 = t_1 < \dots < t_n = 1$ and $\epsilon_k > 0$ such that

- the balls $U(\tilde{\sigma}_{x\bar{x}}(t_1), \epsilon_1), \dots, U(\tilde{\sigma}_{x\bar{x}}(t_n), \epsilon_n)$ cover $\tilde{\sigma}_{x\bar{x}}$,
- the balls $U(\tilde{\sigma}_{y\bar{y}}(t_1), \epsilon_1), \dots, U(\tilde{\sigma}_{y\bar{y}}(t_n), \epsilon_n)$ cover $\tilde{\sigma}_{y\bar{y}}$, and
- for all $p, \bar{p} \in U(\tilde{\sigma}_{x\bar{x}}(t_k), \epsilon_k)$ and $q, \bar{q} \in U(\tilde{\sigma}_{y\bar{y}}(t_k), \epsilon_k)$, the map $t \mapsto d(\tilde{\sigma}_{pq}(t), \tilde{\sigma}_{\bar{p}\bar{q}}(t))$ is convex.

Consider now a sequence $0 = s_0 < s_1 < \dots < s_n = 1$ with

$$\begin{aligned} \tilde{\sigma}_{x\bar{x}}(s_k) &\in U(\tilde{\sigma}_{x\bar{x}}(t_k), \epsilon_k) \cap U(\tilde{\sigma}_{x\bar{x}}(t_{k+1}), \epsilon_{k+1}), \\ \tilde{\sigma}_{y\bar{y}}(s_k) &\in U(\tilde{\sigma}_{y\bar{y}}(t_k), \epsilon_k) \cap U(\tilde{\sigma}_{y\bar{y}}(t_{k+1}), \epsilon_{k+1}), \end{aligned}$$

for $k = 1, \dots, n-1$. Then we get

$$\begin{aligned} &d(\tilde{\sigma}_{xy}(t), \tilde{\sigma}_{\bar{x}\bar{y}}(t)) \\ &\leq \sum_{k=1}^n d(\tilde{\sigma}_{\tilde{\sigma}_{x\bar{x}}(s_{k-1})\tilde{\sigma}_{y\bar{y}}(s_{k-1})}(t), \tilde{\sigma}_{\tilde{\sigma}_{x\bar{x}}(s_k)\tilde{\sigma}_{y\bar{y}}(s_k)}(t)) \\ &\leq (1-t) \left(\sum_{k=1}^n d(\tilde{\sigma}_{x\bar{x}}(s_{k-1}), \tilde{\sigma}_{x\bar{x}}(s_k)) \right) + t \left(\sum_{k=1}^n d(\tilde{\sigma}_{y\bar{y}}(s_{k-1}), \tilde{\sigma}_{y\bar{y}}(s_k)) \right) \\ &= (1-t)d(x, \bar{x}) + td(y, \bar{y}). \end{aligned}$$

Hence, $\tilde{\sigma}$ is a convex geodesic bicombing on X .

If σ is reversible, then $\tilde{\sigma}_{xy}^*(t) := \tilde{\sigma}_{yx}(1-t)$ also defines a convex geodesic bicombing on X which is consistent with σ . Therefore, by uniqueness, $\tilde{\sigma}^*$ and $\tilde{\sigma}$ coincide, i.e. $\tilde{\sigma}$ is reversible. \square

3. Locally injective metric spaces

N. Aronszajn and P. Panitchpakdi [AP] proved that injective metric spaces are exactly the *hyperconvex* metric spaces, namely metric spaces with the property that for every family of closed balls $\{B(x_i, r_i)\}_{i \in I}$ with $d(x_i, x_j) \leq r_i + r_j$, for all $i, j \in I$, we have $\bigcap_{i \in I} B(x_i, r_i) \neq \emptyset$. Note that in hyperconvex metric spaces closed balls are hyperconvex.

Definition 3.1. A metric space X is *locally injective* if, for every $x \in X$, there is some $r_x > 0$ such that $B(x, r_x)$ is injective. If we can take $r_x = r$ for all x , we call X *uniformly locally injective*.

Lemma 3.2. *Let X be a metric space with the property that every closed ball $B(x, r)$ is injective, then X is itself injective.*

Proof. Let $\{B(x_i, r_i)\}_{i \in I}$ be a family of closed balls with $d(x_i, x_j) \leq r_i + r_j$. Fix some $i_0 \in I$ and set $A_i := B(x_i, r_i) \cap B(x_{i_0}, r_{i_0})$. Since, for r big enough, we have $x_i, x_j \in B(x_{i_0}, r)$, we get that the A_i 's are externally hyperconvex in A_{i_0} and $A_i \cap A_j \neq \emptyset$ for all $i, j \in I$. Hence, it follows

$$\bigcap_{i \in I} B(x_i, r_i) = \bigcap_{i \in I} A_i \neq \emptyset$$

by [Mie, Proposition 1.2]. □

Proposition 3.3. *Let X be a uniformly locally injective metric space with a reversible, convex geodesic bicombing σ . Then X is injective.*

Proof. Consider the following property:

P(R): For every family $\{B(x_i, r_i)\}_{i \in I}$ with $d(x_i, x_j) \leq r_i + r_j$ and $r_i \leq R$, there is some $x \in \bigcap_{i \in I} B(x_i, r_i)$.

Since X is uniformly locally injective, this clearly holds for some $R_0 > 0$. Next, we show $P(R) \Rightarrow P(2R)$ and therefore $P(R)$ holds for any $R \geq 0$.

Let $\{B(x_i, r_i)\}_{i \in I}$ be a family of closed balls with $d(x_i, x_j) \leq r_i + r_j$ and $r_i \leq 2R$. For $i, j \in I$, define $y_{ij} := \sigma_{x_i x_j}(\frac{1}{2})$. By convexity of σ , we have

$$d(y_{ij}, y_{ik}) = d(\sigma_{x_i x_j}(\frac{1}{2}), \sigma_{x_i x_k}(\frac{1}{2})) \leq \frac{1}{2}d(x_j, x_k) \leq \frac{r_j}{2} + \frac{r_k}{2}.$$

Hence, for every $i \in I$, there is some $z_i \in \bigcap_{j \in I} B(y_{ij}, \frac{r_j}{2})$. Now, observe that $d(z_i, z_j) \leq d(z_i, y_{ij}) + d(y_{ij}, z_j) \leq \frac{r_i}{2} + \frac{r_j}{2}$ and therefore, we find

$$x \in \bigcap_{i \in I} B(z_i, \frac{r_i}{2}) \subset \bigcap_{i \in I} B(x_i, r_i).$$

Since all balls with center in $B(x, r)$ and radius larger than $2r$ contain $B(x, r)$, $P(R)$ for $R = 2r$ implies that $B(x, r)$ is injective. Hence, by Lemma 3.2, X is injective. \square

Since compact, locally injective metric spaces are always uniformly locally injective we conclude the following.

Corollary 3.4. *Let X be a compact, locally injective metric space with a reversible, convex geodesic bicombing σ . Then X is injective.*

Corollary 3.5. *Let X be a proper, locally injective metric space with a reversible, convex geodesic bicombing σ . Then X is injective.*

Proof. Let $\{B(x_i, r_i)\}_{i \in I}$ be a family of balls with $d(x_i, x_j) \leq r_i + r_j$. Fix some $i_0 \in I$ and define $I_n = \{i \in I : d(x_i, x_{i_0}) \leq n\}$, for $n \in \mathbb{N}$. Since $B(x_{i_0}, n)$ is compact, by the previous corollary, there is some $y_n \in \bigcap_{i \in I_n} B(x_i, r_i)$. Especially, $(y_n)_n \subset B(x_{i_0}, r_{i_0})$ and hence, there is some converging subsequence $y_{n_k} \rightarrow y \in \bigcap_{i \in I} B(x_i, r_i)$. \square

Remark. In [Lan], U. Lang proves that every injective metric space admits a reversible, conical geodesic bicombing (Proposition 3.8). Observe also that this is the only property of the geodesic bicombing used in the proof of Proposition 3.3. Therefore, we get the following equivalence statement (in the terminology of [Lan]): A metric space is injective if and only if it is uniformly locally injective and admits a reversible, conical geodesic bicombing.

If an injective metric space X is proper, it also admits a (possibly non-consistent) convex geodesic bicombing [DL1, Theorem 1.1] and if X has finite combinatorial dimension in the sense of A. Dress [Dre], this convex geodesic bicombing is consistent, reversible and unique [DL1, Theorem 1.2]. In our terms, this is:

Proposition 3.6. *Every proper, injective metric space with finite combinatorial dimension admits a unique reversible, convex geodesic bicombing.*

Recall that, by the Hopf–Rinow Theorem, any complete, locally compact length space is proper.

Corollary 3.7. *Let X be a locally compact, locally injective metric space with locally finite combinatorial dimension. Then X admits a reversible, convex local geodesic bicombing.*

Proof. For every $x \in X$, there is some $r_x > 0$ such that $B(x, 3r_x)$ is compact, injective and has finite combinatorial dimension. This also holds for $B(x, r_x)$ and therefore, there is a reversible, convex geodesic bicombing σ^x on $B(x, r_x)$.

We will check that for $B(x, r_x)$ and $B(y, r_y)$ with $B(x, r_x) \cap B(y, r_y) \neq \emptyset$ the two geodesic bicomblings σ^x, σ^y coincide on the intersection. Assume without loss of generality that $r_x \geq r_y$ and hence $B(x, r_x), B(y, r_y) \subset B(x, 3r_x)$. Then the convex geodesic bicombing τ on $B(x, 3r_x)$ restricts to both $B(x, r_x)$ and $B(y, r_y)$ since, for $p, q \in B(z, r_z)$, we have $d(z, \tau_{pq}(t)) \leq (1 - t)d(z, p) + td(z, q) \leq r_z$. Hence, by uniqueness, the geodesic bicomblings σ^x, σ^y are both restrictions of τ and thus coincide on $B(x, r_x) \cap B(y, r_y)$.

Therefore σ , defined by $\sigma|_{B(x, r_x) \times B(x, r_x)} := \sigma^x|_{B(x, r_x) \times B(x, r_x)}$, is a reversible, convex local geodesic bicombing on X . □

Proof of Theorem 1.2. Let X be a complete, locally compact, simply-connected, locally injective length space with locally finite combinatorial dimension. By Corollary 3.7, X has a reversible, convex local geodesic bicombing, which induces a reversible, convex geodesic bicombing by Theorem 1.1. Hence, we can apply Corollary 3.5 and deduce that X is injective. □

4. Absolute 1-Lipschitz Neighborhood Retracts

Absolute Lipschitz uniform neighborhood retracts appear for instance in the study of approximations of Lipschitz maps, see [HJ, Section 7]. The question arises, how much absolute Lipschitz uniform neighborhood retracts differ from being an absolute Lipschitz retract. Theorem 1.3 will give a first answer in the case of absolute 1-Lipschitz retracts.

A metric space X is an *absolute 1-Lipschitz neighborhood retract* if, for every metric space Y with $X \subset Y$, there is some neighborhood U of X in Y and a 1-Lipschitz retraction $\rho: U \rightarrow X$. Furthermore, if we can take $U = U(X, \epsilon)$ for some $\epsilon > 0$, we call X an *absolute 1-Lipschitz uniform neighborhood retract*. In this case, ϵ can be chosen independent of Y ; see [HJ, Proposition 7.78]. Finally, if there is always a 1-Lipschitz retraction $r: Y \rightarrow X$, then X is an *absolute 1-Lipschitz retract*. This is equivalent to X being an injective metric space [AP, Theorem 8].

Lemma 4.1. *Let X be an absolute 1-Lipschitz (uniform) neighborhood retract. Then X is (uniformly) locally injective.*

Proof. Consider $X \subset l_\infty(X)$. Since X is an absolute 1-Lipschitz neighborhood retract, there is some neighborhood U of X and a 1-Lipschitz retraction $\rho: U \rightarrow X$. For $x \in X$, there is some $r_x > 0$ such that $B(x, r_x) \subset U$. Let now $\{B(x_i, r_i)\}_{i \in I}$ be a family of closed balls with $x_i \in B(x, r_x) \cap X$ and $d(x_i, x_j) \leq r_i + r_j$. Then, since $l_\infty(X)$ is injective, there is some $y \in B(x, r_x) \cap \bigcap_{i \in I} B(x_i, r_i) \subset U$. Hence, we have $\rho(y) \in B(x, r_x) \cap \bigcap_{i \in I} B(x_i, r_i) \cap X$ and therefore $B(x, r_x) \cap X$ is injective.

If X is an absolute 1-Lipschitz uniform neighborhood retract, we have $U = U(X, \epsilon)$ for some $\epsilon > 0$ and therefore, we can choose $r_x = \frac{\epsilon}{2}$ for all $x \in X$. \square

The converse is not true, as the following example shows.

Example 4.2. Consider the unit sphere S^1 endowed with the inner metric. Since, for every $x \in S^1$ and $\epsilon \in (0, \frac{\pi}{2}]$, the ball $B(x, \epsilon)$ is isometric to the interval $[-\epsilon, \epsilon]$, the unit sphere S^1 is uniformly locally injective.

But S^1 is not an absolute 1-Lipschitz neighborhood retract. Fix some inclusion $S^1 \subset l_\infty(S^1)$. We choose three points $x, y, z \in S^1$ with $r := d(x, y) = d(x, z) = d(y, z) = \frac{2\pi}{3}$. Let U be a neighborhood of S^1 in $l_\infty(S^1)$. As U is open, there is some $\epsilon \in (0, \frac{r}{2})$ such that $B(x, \epsilon) \subset U$. By hyperconvexity of $l_\infty(S^1)$, there is some

$$p \in B(x, \epsilon) \cap B(y, r - \epsilon) \cap B(z, r - \epsilon) \subset U.$$

But since

$$B(x, \epsilon) \cap B(y, r - \epsilon) \cap B(z, r - \epsilon) \cap S^1 = \emptyset,$$

there is no 1-Lipschitz retraction $\rho: S^1 \cup \{p\} \rightarrow S^1$.

In fact, the notion of an absolute 1-Lipschitz uniform neighborhood retract is quite restrictive.

Lemma 4.3. *Let X be an absolute 1-Lipschitz uniform neighborhood retract. Then X is*

- (i) *complete,*
- (ii) *geodesic, and*
- (iii) *simply-connected.*

Proof. Fix some inclusion $X \subset l_\infty(X)$ and $r := \frac{\epsilon}{2} > 0$ such that there is a 1-Lipschitz retraction $\rho: U(X, \epsilon) \rightarrow X$.

First, if $(x_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in X , it converges to some $x \in U(X, \epsilon)$. It follows that $x = \rho(x) \in X$.

Next, assume that there is a geodesic in X between points at distance less than d . By Lemma 4.1, this is clearly true for $d = r$. Consider two points $x, y \in X$ with $d(x, y) \leq d + r$. Now, since $l_\infty(X)$ is geodesic, there is some $z \in l_\infty(X)$ with $d(x, y) = d(x, z) + d(z, y)$, $d(x, z) \leq r$ and $d(z, y) \leq d$. But then, we have $\rho(z) \in X$ with $d(x, y) = d(x, \rho(z)) + d(\rho(z), y)$ and, by our hypothesis, there are geodesics from x to $\rho(z)$ and from $\rho(z)$ to y which combine to a geodesic from x to y .

Finally, since X is locally simply-connected, every curve is homotopic to a curve of finite length and hence it is enough to consider loops of finite length. We show that every such loop in X is contractible.

Let γ be a loop in X of length $L(\gamma) = 2\pi R$ with $R > r$. Denote by $S_R^2 := \{x \in \mathbb{R}^3 : |x| = R\}$ the sphere of radius R endowed with the inner metric and let $A := \{x \in S_R^2 : 0 \leq x_3 \leq R \sin(\frac{r}{R})\}$ be the region bounded by the two circles $c := \{x \in S_R^2 : x_3 = 0\}$ and $c' := \{x \in S_R^2 : x_3 = R \sin(\frac{r}{R})\}$. Let $f : c \rightarrow X$ be a parametrization of γ by arclength and let $\bar{f} : A \rightarrow l_\infty(X)$ be a 1-Lipschitz extension. Then $\gamma' := \rho \circ \bar{f}(c')$ is a loop of length $L(\gamma') \leq L(c') = 2\pi R'$ with $R' := R \cos(\frac{r}{R})$, which is homotopic to γ . Since $\cos(\frac{r}{R'}) \leq \cos(\frac{r}{R})$, we find inductively a loop γ_n with $L(\gamma_n) \leq 2\pi R \cos(\frac{r}{R})^n$, which is homotopic to γ .

If $L(\gamma) = 2\pi R$ with $R \leq r$, we can use the same argument with A replaced by the upper hemisphere of radius R to show that γ is contractible. \square

We conclude that an absolute 1-Lipschitz uniform neighborhood retract is a complete, simply-connected, locally injective length space and therefore Theorem 1.3 follows directly from Theorem 1.2.

Acknowledgments. I would like to thank Prof. Dr. Urs Lang for helpful remarks on this work and Dr. Maël Pavón for inspiring discussions. I am also grateful for Giuliano Basso's comments related to Theorem 1.1. The author was supported by the Swiss National Science Foundation.

References

- [AB] S. ALEXANDER and R. BISHOP, The Hadamard–Cartan theorem in locally convex metric spaces. *Enseign. Math.* (2) **36** (1990), 309–320. Zbl 0718.53055 MR1096422
- [ABr] J.M. ALONSO and M.R. BRIDSON, Semihyperbolic groups. *Proc. London Math. Soc.* (3) **70** (1995), 56–114. Zbl 0823.20035 MR1300841
- [AP] N. ARONSZAJN and P. PANITCHPAKDI, Extension of uniformly continuous transformations and hyperconvex metric spaces. *Pacific J. Math.* **6** (1956), 405–439. Zbl 0074.17802 MR 0084762

- [Bal] W. BALLMANN, Singular spaces of nonpositive curvature. Sur les groupes hyperboliques d'après Mikhael Gromov (Bern, 1988), 189–201, Progr. Math., 83, Birkhäuser Boston, Boston, MA, 1990. MR1086658
- [Bas] G. BASSO, Fixed point theorems for metric spaces with a conical geodesic bicombing. *Ergodic Theory Dynam. Systems* (2017), Doi10.1017/etds.2016.106.
- [BH] M. BRIDSON and A. HAEFLIGER, *Metric Spaces of Non-Positive Curvature*. Springer-Verlag, 1999. Zbl0988.53001 MR1744486
- [BM] G. BASSO and B. MIESCH, Conical geodesic bicomblings on subsets of normed vector spaces. arXiv:1604.04163, Aug 2016. To appear in *Adv. Geom.*
- [Des] D. DESCOMBES, Asymptotic rank of spaces with bicomblings. *Math. Z.* **284** (2016), 947–960. Zbl1360.53078 MR3563261
- [DL1] D. DESCOMBES and U. LANG, Convex geodesic bicomblings and hyperbolicity. *Geom. Dedicata* **177** (2015), 367–384. Zbl1343.53036 MR3370039
- [DL2] — Flats in spaces with convex geodesic bicomblings. *Anal. Geom. Metr. Spaces* **4** (2016), 68–84. Zbl1341.53070 MR3483604
- [Dre] A. W. M. DRESS, Trees, tight extensions of metric spaces, and the cohomological dimension of certain groups: A note on combinatorial properties of metric spaces. *Adv. in Math.* **53** (1984), 321–402. Zbl0562.54041 MR0753872
- [EK] R. ESPÍNOLA and M. A. KHAMSI, Introduction to hyperconvex spaces. *Handbook of Metric Fixed Point Theory*, 391–435, Kluwer Acad. Publ., Dordrecht, 2001. Zbl1029.47002 MR1904284
- [HJ] P. HÁJEK and M. JOHANIS, *Smooth Analysis in Banach Spaces*. De Gruyter Series in Nonlinear Analysis and Applications, 19. De Gruyter, Berlin, 2014. Zbl1327.46002 MR3244144
- [Isb] J. R. ISBELL, Six theorems about injective metric spaces. *Comment. Math. Helv.* **39** (1964), 65–76. Zbl0151.30205 MR0182949
- [Ito] S. ITOH, Some fixed-point theorems in metric spaces. *Fund. Math.* **102** (1979), 109–117. Zbl0412.54054 MR0525934
- [KL] U. KOHLENBACH and L. LEUȘTEAN, Asymptotically nonexpansive mappings in uniformly convex hyperbolic spaces. *J. Eur. Math. Soc. (JEMS)* **12** (2010), 71–92. Zbl1184.03057 MR2578604
- [Kle] B. KLEINER, The local structure of length spaces with curvature bounded above. *Math. Z.* **231** (1999), 409–456. Zbl0940.53024 MR1704987
- [Koh] U. KOHLENBACH, Some logical metatheorems with applications in functional analysis. *Trans. Amer. Math. Soc.* **357** (2005), 89–128. Zbl1079.03046 MR2098088
- [Lan] U. LANG, Injective hulls of certain discrete metric spaces and groups. *J. Topol. Anal.* **5** (2013), 297–331. Zbl1292.20046 MR3096307
- [Mie] B. MIESCH, Gluing hyperconvex metric spaces. *Anal. Geom. Metr. Spaces* **3** (2015), 102–110. Zbl1321.54053 MR3349339
- [Sin] R. SINE, On linear contraction semigroups in sup norm spaces, *Nonlinear Anal.* **3** (1979), 885–890.

- [Soa] P. M. SOARDI, Existence of fixed points of nonexpansive mappings in certain Banach lattices. *Proc. Amer. Math. Soc.* **73** (1979), 25–29. Zbl 0371.47048 MR 0512051
- [Tak] W. TAKAHASHI, A convexity in metric space and nonexpansive mappings. I. *Kōdai Math. Sem. Rep.* **22** (1970), 142–149. Zbl 0268.54048 MR 0267565

(Reçu le 23 novembre 2016)

Benjamin MIESCH, Department of Mathematics, ETH Zürich,
Raemistrasse 101, 8092 Zürich, Switzerland

e-mail: benjamin.miesch@math.ethz.ch

