

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 63 (2017)
Heft: 3-4

Artikel: The Horn inequalities from a geometric point of view
Autor: Berline, Nicole / Vergne, Michèle / Walter, Michael
DOI: <https://doi.org/10.5169/seals-787391>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The Horn inequalities from a geometric point of view

Nicole BERLINE, Michèle VERGNE and Michael WALTER

Abstract. We give an exposition of the Horn inequalities and their triple role characterizing tensor product invariants, eigenvalues of sums of Hermitian matrices, and intersections of Schubert varieties. We follow Belkale’s geometric method, but assume only basic representation theory and algebraic geometry, aiming for self-contained, concrete proofs. In particular, we do not assume the Littlewood–Richardson rule nor an a priori relation between intersections of Schubert cells and tensor product invariants. Our motivation is largely pedagogical, but the desire for concrete approaches is also motivated by current research in computational complexity theory and effective algorithms.

Mathematics Subject Classification (2010). Primary: 22E46; Secondary: 14L24, 53D20, 15A42.

Keywords. Horn inequalities, Littlewood–Richardson coefficients, saturation property, Fulton conjecture, Kirwan cone.

Contents

1	Introduction	404
2	A panorama of invariants, eigenvalues, and intersections	406
3	Subspaces, flags, positions	415
3.1	Schubert positions	416
3.2	Induced flags and positions	420
3.3	The flag variety	425
4	Intersections and Horn inequalities	429
4.1	Coordinates	429
4.2	Intersections and dominance	430
4.3	Slopes and Horn inequalities	433
5	Sufficiency of Horn inequalities	439
5.1	Tangent maps	439
5.2	Kernel dimension and position	442

5.3	The kernel recurrence	448
6	Invariants and Horn inequalities	450
6.1	Borel–Weil construction	450
6.2	Invariants from intersecting tuples	451
6.3	Kirwan cone and saturation	454
6.4	Invariants and intersection theory	455
7	Proof of Fulton’s conjecture	457
7.1	Nonzero invariants and intersections	457
7.2	Sherman’s refined lemma	461
A	Horn triples in low dimensions	464
B	Kirwan cones in low dimensions	466
	References	468

1. Introduction

The possible eigenvalues of Hermitian matrices X_1, \dots, X_s such that $X_1 + \dots + X_s = 0$ form a convex polytope. They can thus be characterized by a finite set of linear inequalities, most famously so by the inductive system of linear inequalities conjectured by Horn [Hor]. The very same inequalities give necessary and sufficient conditions on highest weights $\lambda_1, \dots, \lambda_s$ such that the tensor product of the corresponding irreducible $\mathrm{GL}(r)$ -representations $L(\lambda_1), \dots, L(\lambda_s)$ contains a nonzero invariant vector, i.e., $c(\vec{\lambda}) := \dim(L(\lambda_1) \otimes \dots \otimes L(\lambda_s))^{\mathrm{GL}(r)} > 0$. For $s = 3$, the multiplicities $c(\vec{\lambda})$ can be identified with the *Littlewood-Richardson coefficients*. Since the Horn inequalities are linear, $c(\vec{\lambda}) > 0$ if and only if $c(N\vec{\lambda}) > 0$ for any integer $N > 0$. This is the celebrated *saturation property* of $\mathrm{GL}(r)$, first established combinatorially by Knutson and Tao [KT] building on work by Klyachko [Kly]. Some years after, Belkale has given an alternative proof of the Horn inequalities and the saturation property [Bel3]. His main insight is to ‘geometrize’ the classical relationship between the invariant theory of $\mathrm{GL}(r)$ and the intersection theory of Schubert varieties of the Grassmannian. In particular, by a careful study of the tangent space of intersections, he shows how to obtain a geometric basis of invariants.

The aim of this text is to give a self-contained exposition of the Horn inequalities, assuming only linear algebra and some basic representation theory and algebraic geometry, similar in spirit to the approach taken in [VW]. We also discuss a proof of Fulton’s conjecture which asserts that $c(\vec{\lambda}) = 1$ if and only if $c(N\vec{\lambda}) = 1$ for any integer $N \geq 1$. We follow Belkale’s geometric method [Bel2, Bel3, Bel4], as recently refined by Sherman [She], and do not claim any originality. Instead, we hope that our text might be useful by

providing a more accessible introduction to these topics, since we tried to give simple and concrete proofs of all results. In particular, we do not use the Littlewood-Richardson rule for determining $c(\vec{\lambda})$, and we do not discuss the relation of a basis of invariants to the integral points of the hive polytope [KT]. Instead, we describe a basis of invariants that can be identified with the Howe-Tan-Willenbring basis, which is constructed using determinants associated to Littlewood-Richardson tableaux, as we explained in [VW]. We will come back to this subject in the future. We note that Derksen and Weyman's work [DW] can be understood as a variant of the geometric approach in the context of quivers. For alternative accounts we refer to the work by Knutson and Tao [KT] and Woodward [KTW], Ressayre [Res1, Res3] and to the expositions by Fulton and Knutson [Ful2, Knu].

The desire for concrete approaches to questions of representation theory and algebraic geometry is also motivated by recent research in computational complexity and the interest in efficient algorithms. Indeed, the saturation property implies that deciding the nonvanishing of a Littlewood-Richardson coefficient can be decided in polynomial time [MNS]. In contrast, the analogous problem for the Kronecker coefficients, which are not saturated, is NP-hard, but believed to simplify in the asymptotic limit [IMW, BCM]. We refer to [Mul, BLMW] for further detail.

These notes are organized as follows: In Section 2, we start by motivating the triple role of the Horn inequalities characterizing invariants, eigenvalues, and intersections. Then, in Section 3, we collect some useful facts about positions and flags. This is used in Sections 4 and 5 to establish Belkale's theorem characterizing intersecting Schubert varieties in terms of Horn's inequalities. In Section 6, we explain how to construct a geometric basis of invariants from intersecting Schubert varieties. This establishes the Horn inequalities for the Littlewood-Richardson coefficients, and thereby the saturation property, as well as for the eigenvalues of Hermitian matrices that sum to zero. In Section 7, we sketch how Fulton's conjecture can be proved geometrically by similar techniques. Lastly, in the appendix, we have collected the Horn inequalities for three tensor factors and low dimensions.

Notation. We write $[n] := \{1, \dots, n\}$ for any positive integer n . For any group G and representation M , we write M^G for the linear subspace of G -invariant vectors. For any subgroup $H \subseteq G$, we denote by $G/H = \{gH\}$ the right coset space. If F is an H -space, we denote by $G \times_H F$ the quotient of $G \times F$ by the equivalence relation $(g, f) \sim (gh^{-1}, hf)$ for $g \in G$, $f \in F$, $h \in H$. Note that $G \times_H F$ is a G -space fibered over G/H , with fiber F . If F is a subspace of a G -space X , then $G \times_H F$ is identified by the G -equivariant map

$[g, f] \mapsto (gH, gf)$ with the subspace of $G/H \times X$ (equipped with the diagonal G -action) consisting of the (gH, x) such that $g^{-1}x \in F$.

2. A panorama of invariants, eigenvalues, and intersections

In this section we give a panoramic overview of the relationship between invariants, eigenvalues, and intersections. Our focus is on explaining the intuition, connections, and main results. To keep the discussion streamlined, more difficult proofs are postponed to later sections (in which case we use the numbering of the later section, so that the proofs can easily be found). The rest of this article, from Section 3 onwards, is concerned with developing the necessary mathematical theory and giving these proofs.

We start by recalling the basic representation theory of the general linear group $GL(r) := GL(r, \mathbb{C})$. Consider \mathbb{C}^r with the ordered standard basis $e(1), \dots, e(r)$ and standard Hermitian inner product. Let $H(r)$ denote the subgroup of invertible matrices $t \in GL(r)$ that are diagonal in the standard basis, i.e., $t e(i) = t(i) \cdot e(i)$ with all $t(i) \neq 0$. We write $t = (t(1), \dots, t(r))$ and thereby identify $H(r) \cong (\mathbb{C}^*)^r$. To any sequence of integers $\mu = (\mu(1), \dots, \mu(r))$, we can associate a character of $H(r)$ by $t \mapsto t^\mu := t(1)^{\mu(1)} \dots t(r)^{\mu(r)}$. We say that μ is a weight and call $\Lambda(r) = \mathbb{Z}^r$ the weight lattice. A weight is dominant if $\mu(1) \geq \dots \geq \mu(r)$, and the set of all dominant weights form a semigroup, denoted by $\Lambda_+(r)$. We later also consider antidominant weights ω , which satisfy $\omega(1) \leq \dots \leq \omega(r)$.

For any dominant weight $\lambda \in \Lambda_+(r)$, there is a unique irreducible representation $L(\lambda)$ of $GL(r)$ with highest weight λ . That is, if $B(r)$ denotes the group of upper-triangular invertible matrices (the standard Borel subgroup of $GL(r)$) and $N(r) \subseteq B(r)$ the subgroup of upper-triangular matrices with all ones on the diagonal (i.e., the corresponding unipotent), then $L(\lambda)^{N(r)} = \mathbb{C}v_\lambda$ is a one-dimensional eigenspace of $B(r)$ of $H(r)$ -weight λ . We say that v_λ is a highest weight vector of $L(\lambda)$. In Section 6.1 we describe a concrete construction of $L(\lambda)$ due to Borel and Weil. Now let $U(r)$ denote the group of unitary matrices, which is a maximally compact subgroup of $GL(r)$. We can choose an $U(r)$ -invariant Hermitian inner product $\langle \cdot, \cdot \rangle$ (by convention complex linear in the second argument) on each $L(\lambda)$ so that the representation $L(\lambda)$ restricts to an irreducible unitary representation of $U(r)$. Any two such representations of $U(r)$ are pairwise inequivalent, and, by Weyl's trick, any irreducible unitary representation can be obtained in this way. Let us now decompose their Lie algebras as $\mathfrak{gl}(r) = \mathfrak{u}(r) \oplus i\mathfrak{u}(r)$, where $i = \sqrt{-1}$, and likewise $\mathfrak{h}(r) = \mathfrak{t}(r) \oplus i\mathfrak{t}(r)$, where we write $\mathfrak{t}(r)$ for the Lie algebra of $T(r)$, the group of diagonal unitary

matrices, and similarly for the other Lie groups. Here, $iu(r)$ denotes the space of Hermitian matrices and $it(r)$ the subspace of diagonal matrices with real entries. We freely identify vectors in \mathbb{R}^r with the corresponding diagonal matrices in $it(r)$ and denote by (\cdot, \cdot) the usual inner product of $it(r) \cong \mathbb{R}^r$. For a subset $J \subseteq [r]$, we write T_J for the vector (diagonal matrix) in $it(r)$ that has ones in position J , and otherwise zero.

Now let \mathcal{O}_λ denote the set of Hermitian matrices with eigenvalues $\lambda(1) \geq \dots \geq \lambda(r)$. By the spectral theorem, \mathcal{O}_λ is a $U(r)$ -orbit with respect to the adjoint action, $u \cdot X := uXu^*$, and so $\mathcal{O}_\lambda = U(r) \cdot \lambda$, where we identify λ with the diagonal matrix with entries $\lambda(1) \geq \dots \geq \lambda(r)$. On the other hand, recall that any invertible matrix $g \in GL(r)$ can be written as a product $g = ub$, where $u \in U(r)$ is unitary and $b \in B(r)$ upper-triangular. Since v_λ is an eigenvector of $B(r)$, it follows that, in projective space $\mathbb{P}(L(\lambda))$, the orbits of $[v_\lambda]$ for $GL(r)$ and $U(r)$ are the same! Moreover, it is not hard to see that the $U(r)$ -stabilizers of λ and of $[v_\lambda]$ agree, so we obtain a $U(r)$ -equivariant diffeomorphism

$$(2.1) \quad \mathcal{O}_\lambda \rightarrow U(r) \cdot [v_\lambda] = GL(r) \cdot [v_\lambda] \subseteq \mathbb{P}(L(\lambda)), \quad u \cdot \lambda \mapsto u \cdot [v_\lambda] = [u \cdot v_\lambda]$$

which also allows us to think of the adjoint orbit \mathcal{O}_λ as a complex projective $GL(r)$ -variety. An important observation is that

$$(2.2) \quad \text{tr}((u \cdot \lambda)A) = \frac{\langle u \cdot v_\lambda, \rho_\lambda(A)(u \cdot v_\lambda) \rangle}{\|v_\lambda\|^2}$$

for all complex $r \times r$ -matrices A , i.e., elements of the Lie algebra $\mathfrak{gl}(r)$ of $GL(r)$; ρ_λ denotes the Lie algebra representation on $L(\lambda)$. To see that (2.2) holds true, we may assume that $\|v_\lambda\| = 1$ as well as that $u = 1$, the latter by $U(r)$ -equivariance. Now $\text{tr}(A\lambda) = \langle v_\lambda, \rho_\lambda(A)v_\lambda \rangle$ is easily verified by decomposing $A = L + H + R$ with L strictly lower triangular, R strictly upper triangular, and $H \in \mathfrak{h}(r)$ diagonal and comparing term by term. These observations lead to the following fundamental connection between the eigenvalues of Hermitian matrices and the invariant theory of the general linear group:

Proposition 2.3 (Kempf-Ness, [KN]). *Let $\lambda_1, \dots, \lambda_s$ be dominant weights for $GL(r)$ such that $(L(\lambda_1) \otimes \dots \otimes L(\lambda_s))^{GL(r)} \neq \{0\}$. Then there exist Hermitian matrices $X_k \in \mathcal{O}_{\lambda_k}$ such that $\sum_{k=1}^s X_k = 0$.*

Proof. Let $0 \neq w \in (L(\lambda_1) \otimes \dots \otimes L(\lambda_s))^{GL(r)}$ be a nonzero invariant vector. Then, $P(v) := \langle w, v \rangle$ is a nonzero linear function on $L(\lambda_1) \otimes \dots \otimes L(\lambda_s)$ that is invariant under the diagonal action of $GL(r)$; indeed, $\langle w, g \cdot v \rangle = \langle g^* \cdot w, v \rangle = \langle w, v \rangle$. Since the $L(\lambda_k)$ are irreducible, they are spanned by the orbits $U(r)v_{\lambda_k}$. Thus we can find $u_1, \dots, u_s \in U(r)$ such that $P(v) \neq 0$ for $v = (u_1 \cdot v_{\lambda_1}) \otimes \dots \otimes (u_s \cdot v_{\lambda_s})$.

Consider the class $[v]$ of v in the corresponding projective space $\mathbb{P}(L(\lambda_1) \otimes \cdots \otimes L(\lambda_s))$. The orbit of $[v]$ under the diagonal $GL(r)$ -action is contained in the $GL(r)^s$ -orbit, which is the closed set $[U(r) \cdot v_{\lambda_1} \otimes \cdots \otimes U(r) \cdot v_{\lambda_s}]$ according to the discussion preceding (2.1). It follows that $GL(r) \cdot v$ and its closure, $\overline{GL(r) \cdot v}$ (say, in the Euclidean topology), are contained in the closed set $\{\kappa(u'_1 \cdot v_{\lambda_1}) \otimes \cdots \otimes (u'_s \cdot v_{\lambda_s})\}$ for $\kappa \in \mathbb{C}$ and $u'_1, \dots, u'_s \in U(r)$.

Since P is $GL(r)$ -invariant, $P(v') = P(v) \neq 0$ for any vector v' in the diagonal $GL(r)$ -orbit of v . By continuity, this is also true in the orbits' closure, $\overline{GL(r) \cdot v}$. On the other hand, $P(0) = 0$. It follows that $0 \notin \overline{GL(r) \cdot v}$, i.e., the origin does not belong to the orbit closure. Consider then a nonzero vector v' of minimal norm in $\overline{GL(r) \cdot v}$. By the discussion in the preceding paragraph, this vector is of the form $v' = \kappa(u'_1 \cdot v_{\lambda_1}) \otimes \cdots \otimes (u'_s \cdot v_{\lambda_s})$ for some $0 \neq \kappa \in \mathbb{C}$ and $u'_1, \dots, u'_s \in U(r)$. By rescaling v we may moreover assume that $\kappa = 1$, so that v' is a unit vector.

The vector v' is by construction a vector of minimal norm in its own $GL(r)$ -orbit. It follows that, for any Hermitian matrix A ,

$$\begin{aligned} 0 &= \frac{1}{2} \partial_{t=0} \| (e^{At} \otimes \cdots \otimes e^{At}) \cdot v' \|^2 \\ &= \langle v', (\rho_{\lambda_1}(A) \otimes I \otimes \cdots \otimes I + \cdots + I \otimes \cdots \otimes I \otimes \rho_{\lambda_s}(A)) v' \rangle \\ &= \sum_{k=1}^s \langle u'_k \cdot v_{\lambda_k}, \rho_{\lambda_k}(A)(u'_k \cdot v_{\lambda_k}) \rangle = \sum_{k=1}^s \text{tr}(A(u'_k \cdot \lambda_k)) = \sum_{k=1}^s \text{tr}(AX_k), \end{aligned}$$

where we have used Eq. (2.2) and set $X_k := u'_k \cdot \lambda_k$ for $k \in [s]$. This implies at once that $\sum_{k=1}^s X_k = 0$. □

The adjoint orbits $\mathcal{O}_\lambda = U(r) \cdot \lambda$ (but not the map (2.1)) can be defined not only for dominant weights λ but in fact for arbitrary Hermitian matrices. Conversely, any Hermitian matrix is conjugate to a unique element $\xi \in i\mathfrak{t}(r)$ such that $\xi(1) \geq \cdots \geq \xi(r)$. The set of all such ξ is a convex cone, known as the positive Weyl chamber $C_+(r)$, and it contains the semigroup of dominant weights. Throughout this text, we only ever write $\mathcal{O}_\xi = U(r) \cdot \xi$ for ξ that are in the positive Weyl chamber. For example, if $\xi \in C_+(r)$ then $-\xi \in \mathcal{O}_{\xi^*}$, where $\xi^* = (-\xi(r), \dots, -\xi(1)) \in C_+(r)$. If λ is a dominant weight then $\lambda^* = (-\lambda(d), \dots, -\lambda(1))$ is the highest weight of the dual representation of $L(\lambda)$, i.e., $L(\lambda^*) \cong L(\lambda)^*$.

Remark. Using the inner product $(A, B) := \text{tr}(AB)$ on Hermitian matrices we may also think of λ as an element in $i\mathfrak{t}(r)^*$ and of \mathcal{O}_λ as a *coadjoint* orbit in $i\mathfrak{u}(r)^*$. From the latter point of view, the map $(X_1, \dots, X_s) \mapsto \sum_{k=1}^s X_k$ is the moment map for the diagonal $U(r)$ -action on the product of Hamiltonian

manifolds \mathcal{O}_{λ_k} , $k \in [s]$. Proposition 2.3 thus relates the existence of nonzero invariants to the statement that the zero set of the corresponding moment map is nonempty. This is a general fact of Mumford's geometric invariant theory.

Definition 2.4. The *Kirwan cone* $\text{Kirwan}(r, s)$ is defined as the set of $\vec{\xi} = (\xi_1, \dots, \xi_s) \in C_+(r)^s$ such that there exist $X_k \in \mathcal{O}_{\xi_k}$ with $\sum_{k=1}^s X_k = 0$.

Using this language, Proposition 2.3 asserts that if the generalized Littlewood-Richardson coefficient $c(\vec{\lambda}) := \dim(L(\lambda_1) \otimes \dots \otimes L(\lambda_s))^{\text{GL}(r)} > 0$ is nonzero then $\vec{\lambda}$ is a point in the Kirwan cone $\text{Kirwan}(r, s)$.

Remark. We will see in Section 6 that, conversely, if $\vec{\lambda} \in \text{Kirwan}(r, s)$, then $c(\vec{\lambda}) > 0$ (by constructing an explicit nonzero invariant). As a consequence, it will follow that $c(\vec{\lambda}) > 0$ if and only if $c(N\vec{\lambda}) > 0$ for some integer $N > 0$. This is the remarkable saturation property of the Littlewood-Richardson coefficients. In fact, we will show that the Horn inequalities give a complete set of conditions for nonvanishing $c(\vec{\lambda})$ as well as for $\vec{\xi} \in \text{Kirwan}(r, s)$, which in particular establishes that $\text{Kirwan}(r, s)$ is indeed a convex polyhedral cone. We will come back to these points at the end of this section.

If there exist permutations w_k such that $\sum_{k=1}^s w_k \cdot \xi_k = 0$ then $\vec{\xi} \in \text{Kirwan}(r, s)$ (choose each X_k as the diagonal matrix $w_k \cdot \xi_k$). This suffices to characterize the Kirwan cone for $s \leq 2$:

Example. For $s = 1$, it is clear that $\text{Kirwan}(r, 1) = \{0\}$. When $s = 2$, then $\text{Kirwan}(r, 2) = \{(\xi, \xi^*)\}$. Indeed, if $X_1 \in \mathcal{O}_{\xi_1}$ and $X_2 \in \mathcal{O}_{\xi_2}$ with $X_1 + X_2 = 0$, then $X_2 = -X_1 \in \mathcal{O}_{\xi_1^*}$.

In general, however, it is quite delicate to determine if a given $\vec{\xi} \in C_+(r)^s$ is in $\text{Kirwan}(r, s)$ or not. Clearly, one necessary condition is that $\sum_{k=1}^s |\xi_k| = 0$, where we have defined $|\mu| := \sum_{j=1}^r \mu(j)$ for an arbitrary $\mu \in \mathfrak{h}(r)$. This follows by taking the trace of the equation $\sum_{k=1}^s X_k = 0$. In fact, it is clear that by adding or subtracting appropriate multiples of the identity matrix we can always reduce to the case where each $|\xi_k| = 0$.

Example. Let $X_k \in \mathcal{O}_{\xi_k}$ such that $\sum_{k=1}^s X_k = 0$. For each k , let v_k denote a unit eigenvector of X_k with eigenvalue $\xi_k(1)$. Then we have

$$(2.5) \quad 0 = \langle v_k, (\sum_{l=1}^s X_l)v_k \rangle = \xi_k(1) + \sum_{l \neq k} \langle v_k, X_l v_k \rangle \geq \xi_k(1) + \sum_{l \neq k} \xi_l(r)$$

since $\xi_l(r) = \min_{\|v\|=1} \langle v, X_l v \rangle$ by the variational principle for the minimal eigenvalue of a Hermitian matrix X_l . These inequalities, together with $\sum_{k=1}^s |\xi_k| = 0$, characterize the Kirwan cone for $r = 2$, as can be verified by brute force.

There is also a pleasant geometric way of understanding these inequalities in the case $r = 2$. As discussed above, we may assume that the X_k are traceless, i.e., that $\xi_k = (j_k, -j_k)$ for some $j_k \geq 0$. Recall that the traceless Hermitian matrices form a three-dimensional real vector space, spanned by the Pauli matrices. Thus each X_k identifies with a vector $x_k \in \mathbb{R}^3$, and the condition that $X_k \in \mathcal{O}_{\xi_k}$ translates into $\|x_k\| = j_k$. Thus we seek to characterize necessary and sufficient conditions on the lengths j_k of vectors x_k that sum to zero, $\sum_{k=1}^s x_k = 0$. By the triangle inequality, $j_k = \|x_k\| \leq \sum_{l \neq k} \|x_l\| = \sum_{l \neq k} j_l$, which is equivalent to the above. It is instructive to observe that $j_k \leq \sum_{l \neq k} j_l$ is precisely the Clebsch-Gordan rule for $\mathrm{SL}(2)$ when the j_k are half-integers.

The proof of Eq. (2.5), which was valid for any s and r , suggests that a more general variational principle for eigenvalues might be useful to produce linear inequalities for the Kirwan cones.

Definition 2.6. A (*complete*) flag F on a vector space V , $\dim V = r$, is a chain of subspaces

$$\{0\} = F(0) \subset F(1) \subset \cdots \subset F(j) \subset F(j+1) \subset \cdots \subset F(r) = V,$$

such that $\dim F(j) = j$ for all $j = 0, \dots, r$. Any ordered basis $f = (f(1), \dots, f(r))$ of V determines a flag by $F(j) = \mathrm{span}\{f(1), \dots, f(j)\}$. We say that f is *adapted* to F .

Now let $X \in \mathcal{O}_{\xi}$ be a Hermitian matrix with eigenvalues $\xi(1) \geq \cdots \geq \xi(r)$. Let $(f_X(1), \dots, f_X(r))$ denote an orthonormal eigenbasis, ordered correspondingly, and denote by F_X the corresponding *eigenflag* of X , defined as above. Note that F_X is uniquely defined if the eigenvalues $\xi(j)$ are all distinct. We can quantify the position of a subspace with respect to a flag in the following way:

Definition 2.7. The *Schubert position* of an d -dimensional subspace $S \subseteq V$ with respect to a flag F on V is the strictly increasing sequence J of integers defined by

$$J(b) := \min \{j \in [r], \dim F(j) \cap S = b\}$$

for $b \in [d]$. We write $\mathrm{Pos}(S, F) = J$ and freely identify J with the subset $\{J(1) < \cdots < J(d)\}$ of $[r]$. In particular, $\mathrm{Pos}(S, F) = \emptyset$ for $S = \{0\}$ the zero-dimensional subspace.

The upshot of these definitions is the following variational principle:

Lemma 2.8. *Let $\xi \in C_+(r)$, $X \in \mathcal{O}_\xi$ with eigenflag F_X , and $J \subseteq [r]$ a subset of cardinality d . Then,*

$$\min_{S: \text{Pos}(S, F_X) = J} \text{tr}(P_S X) = \sum_{j \in J} \xi(j) = (T_J, \xi),$$

where P_S denotes the orthogonal projector onto an d -dimensional subspace $S \subseteq \mathbb{C}^r$.

Proof. Recall that $F_X(j) = \text{span}\{f_X(1), \dots, f_X(j)\}$, where $(f_X(1), \dots, f_X(r))$ is an orthonormal eigenbasis of X , ordered according to $\xi(1) \geq \dots \geq \xi(r)$. Given a subspace S with $\text{Pos}(S, F_X) = J$, we can find an ordered orthonormal basis $(s(1), \dots, s(d))$ of S where each $s(a) \in F_X(J(a))$. Therefore,

$$\text{tr}(P_S X) = \sum_{a=1}^d \langle s(a), X s(a) \rangle \geq \sum_{a=1}^d \xi(J(a)) = \sum_{j \in J} \xi(j).$$

The inequality holds term by term, as the Hermitian matrix obtained by restricting X to the subspace $F_X(J(a))$ has smallest eigenvalue $\xi(J(a))$. Since $\text{tr}(P_S X) = \sum_{j \in J} \xi(j)$ for $S = \text{span}\{f_X(j) : j \in J\}$, this establishes the lemma. \square

Recall that the Grassmannian $\text{Gr}(d, V)$ is the space of d -dimensional subspaces of V . We may partition $\text{Gr}(d, V)$ according to the Schubert position with respect to a fixed flag:

Definition 2.9. Let F be a flag on V , $\dim V = r$, and $J \subseteq [r]$ a subset of cardinality d . The *Schubert cell* is

$$\Omega_J^0(F) = \{S \subseteq V : \dim S = d, \text{Pos}(S, F) = J\}.$$

The *Schubert variety* $\Omega_J(F)$ is defined as the closure of $\Omega_J^0(F)$ in the Grassmannian $\text{Gr}(d, V)$.

The closures in the Euclidean and Zariski topology coincide; the $\Omega_J(F)$ are indeed algebraic varieties. Using these definitions, Lemma 2.8 asserts that $\min_{S \in \Omega_J^0(F_X)} \text{tr}(P_S X) = \sum_{j \in J} \xi(j)$ for any $X \in \mathcal{O}_\xi$. Since the orthogonal projector P_S is a continuous function of $S \in \text{Gr}(d, V)$ (in fact, the Grassmannian is homeomorphic to the space of orthogonal projectors of rank d), it follows at once that

$$(2.10) \quad \min_{S \in \Omega_J(F_X)} \text{tr}(P_S X) = \sum_{j \in J} \xi(j) = (T_J, \xi).$$

As a consequence, intersections of Schubert varieties imply linear inequalities of eigenvalues of matrices summing to zero:

Lemma 2.11. *Let $X_k \in \mathcal{O}_{\xi_k}$ be Hermitian matrices with $\sum_{k=1}^s X_k = 0$. If $J_1, \dots, J_s \subseteq [r]$ are subsets of cardinality d such that $\bigcap_{k=1}^s \Omega_{J_k}(F_{X_k}) \neq \emptyset$, then $\sum_{k=1}^s (T_{J_k}, \xi_k) \leq 0$.*

Proof. Let $S \in \bigcap_{k=1}^s \Omega_{J_k}(F_{X_k})$. Then, $0 = \sum_{k=1}^s \text{tr}(P_S X_k) \geq \sum_{k=1}^s (T_{J_k}, \xi_k)$ by (2.10). \square

Remarkably, we will find that it suffices to consider only those J_1, \dots, J_s such that $\bigcap_{k=1}^s \Omega_{J_k}(F_k) \neq \emptyset$ for all flags F_1, \dots, F_s . We record the corresponding eigenvalue inequalities, together with the trace condition, in Corollary 2.13 below. Following [Bel3], we denote s -tuples by calligraphic letters, e.g., $\mathcal{J} = (J_1, \dots, J_s)$, $\mathcal{F} = (F_1, \dots, F_s)$, etc. In the case of Greek letters we continue to write $\vec{\lambda} = (\lambda_1, \dots, \lambda_s)$, etc., as above.

Definition 2.12. We denote by $\text{Subsets}(d, r, s)$ the set of s -tuples \mathcal{J} , where each J_k is a subset of $[r]$ of cardinality d . Given such a \mathcal{J} , let \mathcal{F} be an s -tuple of flags on V , with $\dim V = r$. Then we define

$$\Omega_{\mathcal{J}}^0(\mathcal{F}) := \bigcap_{k=1}^s \Omega_{J_k}^0(F_k), \quad \Omega_{\mathcal{J}}(\mathcal{F}) := \bigcap_{k=1}^s \Omega_{J_k}(F_k).$$

We shall say that \mathcal{J} is *intersecting* if $\Omega_{\mathcal{J}}(\mathcal{F}) \neq \emptyset$ for every s -tuple of flags \mathcal{F} , and we denote the set of such \mathcal{J} by $\text{Intersecting}(d, r, s) \subseteq \text{Subsets}(d, r, s)$.

Corollary 2.13 (Klyachko, [Kly]). *If $\vec{\xi} \in \text{Kirwan}(r, s)$ then $\sum_{k=1}^s |\xi_k| = 0$, and for any $0 < d < r$ and any s -tuple $\mathcal{J} \in \text{Intersecting}(d, r, s)$ we have that $\sum_{k=1}^s (T_{J_k}, \xi_k) \leq 0$.*

Example. If $J = \{1, \dots, d\} \subseteq [r]$ then $\Omega_J^0(F) = \{F(d)\}$ is a single point. On the other end, if $J = \{r - d + 1, \dots, r\}$ then $\Omega_J^0(F)$ is dense in $\text{Gr}(r, V)$, so that $\Omega_J(F) = \text{Gr}(r, V)$. It follows that $\mathcal{J} = (J_1, \{r - d + 1, \dots, r\}, \dots, \{r - d + 1, \dots, r\}) \in \text{Intersecting}(d, r, s)$ is intersecting for any J_1 (and likewise for permutations of the s factors).

For $d = 1$, this means that $\Omega_{\{r\}}(F) = \mathbb{P}(V)$, so that (2.10) reduces to the variational principle for the minimal eigenvalue, $\xi(r) = \min_{\|v\|=1} \langle v, Xv \rangle$, which we used to derive (2.5) above. Indeed, since $(\{a\}, \{r\}, \dots, \{r\})$ is intersecting for any a , we find that (2.5) is but a special case of Corollary 2.13.

In order to understand the linear inequalities in Corollary 2.13, we need to understand the sets of intersecting tuples. In the remainder of this section we thus motivate Belkale's inductive system of conditions for an s -tuple to be intersecting. For reasons that will become clear shortly, we slightly change notation: E will be

a complete flag on some n -dimensional vector space W , I will be a subset of $[n]$ of cardinality r , and hence $\Omega_I^0(E)$ will be a Schubert cell in the Grassmannian $\text{Gr}(r, W)$. We will describe $\text{Gr}(r, W)$ and $\Omega_I^0(E)$ in detail in Section 3. For now, we note that the dimension of $\text{Gr}(r, W)$ is $r(n - r)$. In fact, $\text{Gr}(r, W)$ is covered by affine charts isomorphic to $\mathbb{C}^{r(n-r)}$. The dimension of a Schubert cell and the corresponding Schubert variety (its Zariski closure) is given by

$$(3.1.8) \quad \dim \Omega_I^0(E) = \dim \Omega_I(E) = \sum_{a=1}^r (I(a) - a) =: \dim I.$$

Indeed, $\Omega_I^0(E)$ is contained in an affine chart $\mathbb{C}^{r(n-r)}$ and is isomorphic to a vector subspace of dimension $\dim I$. So locally $\Omega_I^0(E)$ is defined by $r(n - r) - \dim I$ equations. This is easy to see and we give a proof in Section 3.

Definition 2.14. Let $\mathcal{I} \in \text{Subsets}(r, n, s)$. The *expected dimension* associated with \mathcal{I} is

$$\text{edim } \mathcal{I} := r(n - r) - \sum_{k=1}^s (r(n - r) - \dim I_k).$$

This definition is natural in terms of intersections, as the following lemma shows:

Lemma 2.15. Let \mathcal{E} be an s -tuple of flags on W , $\dim W = n$, and $\mathcal{I} \in \text{Subsets}(r, n, s)$. If $\Omega_{\mathcal{I}}^0(\mathcal{E}) \neq \emptyset$ then its irreducible components (in the sense of algebraic geometry) are all of dimension at least $\text{edim } \mathcal{I}$.

Proof. Each Schubert cell $\Omega_{I_k}^0(E_k)$ is locally defined by $r(n - r) - \dim I_k$ equations. It follows that any irreducible component $\mathcal{Z} \subseteq \Omega_{\mathcal{I}}^0(\mathcal{E}) = \bigcap_{k=1}^s \Omega_{I_k}^0(E_k)$ is locally defined by $\sum_{k=1}^s (r(n - r) - \dim I_k)$ equations. These equations, however, are not necessarily independent. Thus the codimension of \mathcal{Z} is at most that number, and we conclude that $\dim \mathcal{Z} \geq \text{edim } \mathcal{I}$. \square

Belkale’s first observation is that the expected dimension of an intersecting tuple $\mathcal{I} \in \text{Intersecting}(r, n, s)$ is necessarily nonnegative,

$$(4.2.7) \quad \text{edim } \mathcal{I} = r(n - r) - \sum_{k=1}^s (r(n - r) - \dim I_k) \geq 0.$$

This inequality, as well as some others, will be proved in detail in Section 4. For now, we remark that the condition is rather natural from the perspective of Kleiman’s moving lemma. Given $\mathcal{I} \in \text{Intersecting}(r, n, s)$, it not only implies that the intersection of the Schubert cells, $\Omega_{\mathcal{I}}^0(\mathcal{E}) = \bigcap_{k=1}^s \Omega_{I_k}^0(E_k) \neq \emptyset$, is nonempty

for generic flags, but in fact transverse, so that the dimensions of its irreducible components are exactly equal to the expected dimension; hence, $\text{edim } \mathcal{I} \geq 0$.

We now show that (4.2.7) gives rise to an inductive system of conditions. Given a flag E on W and a subspace $V \subseteq W$, we denote by E^V the flag obtained from the distinct subspaces in the sequence $E(i) \cap V$, $i = 0, \dots, n$. Given subsets $I \subseteq [n]$ of cardinality r and $J \subseteq [r]$ of cardinality d , we also define their *composition* IJ as the subset $IJ = \{I(J(1)) < \dots < I(J(d))\} \subseteq [n]$. (For s -tuples \mathcal{I} and \mathcal{J} we define $\mathcal{I}\mathcal{J}$ componentwise.) Then we have the following ‘chain rule’ for positions: If $S \subseteq V \subseteq W$ are subspaces and E is a flag on W then

$$(3.2.9) \quad \text{Pos}(S, E) = \text{Pos}(V, E) \text{Pos}(S, E^V).$$

We also have the following description of Schubert varieties in terms of Schubert cells:

$$(3.1.6) \quad \Omega_I(E) = \bigcup_{I' \leq I} \Omega_{I'}^0(E),$$

where the union is over all subsets $I' \subseteq [n]$ of cardinality r such that $I'(a) \leq I(a)$ for $a \in [r]$. Both statements are not hard to see; we will give careful proofs in Section 3 below. We thus obtain a corresponding chain rule for intersecting tuples:

Lemma 2.16. *If $\mathcal{I} \in \text{Intersecting}(r, n, s)$ and $\mathcal{J} \in \text{Intersecting}(d, r, s)$, then we have $\mathcal{I}\mathcal{J} \in \text{Intersecting}(d, n, s)$.*

Proof. Let \mathcal{E} be an s -tuple of flags on $W = \mathbb{C}^n$. Since \mathcal{I} is intersecting, there exists $V \in \Omega_{\mathcal{I}}(\mathcal{E})$. Let \mathcal{E}^V denote the s -tuple of induced flags on V . Likewise, since \mathcal{J} is intersecting, we can find $S \in \Omega_{\mathcal{J}}(\mathcal{E}^V)$. In particular, $\text{Pos}(V, E_k)(a) \leq I_k(a)$ for $a \in [r]$ and $\text{Pos}(S, E_k^V) \leq J_k(b)$ for $b \in [d]$ by (3.1.6). Thus (3.2.9) shows that $\text{Pos}(S, E_k)(b) = \text{Pos}(V, E_k)(\text{Pos}(S, E_k^V)(b)) \leq \text{Pos}(V, E_k)(J_k(b)) \leq I_k(J_k(b))$. Using (3.1.6) one last time, we conclude that $S \in \Omega_{\mathcal{I}\mathcal{J}}(\mathcal{E})$. □

As an immediate consequence of Inequality (4.2.7) and Lemma 2.16 we obtain the following set of necessary conditions for an s -tuple \mathcal{I} to be intersecting:

Corollary 2.17. *If $\mathcal{I} \in \text{Intersecting}(r, n, s)$ then for any $0 < d < r$ and any s -tuple $\mathcal{J} \in \text{Intersecting}(d, r, s)$ we have that $\text{edim } \mathcal{I}\mathcal{J} \geq 0$.*

Belkale’s theorem asserts that these conditions are also sufficient. In fact, it suffices to restrict to intersecting \mathcal{J} with $\text{edim } \mathcal{J} = 0$:

Definition 2.18. Let $\text{Horn}(r, n, s)$ denote the set of s -tuples $\mathcal{I} \in \text{Subsets}(r, n, s)$ defined by the conditions that $\text{edim } \mathcal{I} \geq 0$ and, if $r > 1$, that

$$\text{edim } \mathcal{I} \mathcal{J} \geq 0$$

for all $\mathcal{J} \in \text{Horn}(d, r, s)$ with $0 < d < r$ and $\text{edim } \mathcal{J} = 0$.

Theorem 5.3.4 (Belkale, [Bel3]). *For $r \in [n]$ and $s \geq 2$, $\text{Intersecting}(r, n, s) = \text{Horn}(r, n, s)$.*

We will prove Theorem 5.3.4 in Section 5. The inequalities defining $\text{Horn}(r, n, s)$ are in fact tightly related to those constraining the Kirwan cone $\text{Kirwan}(r, s)$ and the existence of nonzero invariant vectors. To any s -tuple of dominant weights $\vec{\lambda}$ for $\text{GL}(r)$ such that $\sum_{k=1}^s |\lambda_k| = 0$, we will associate an s -tuple $\mathcal{I} \in \text{Subsets}(r, n, s)$ for some $[n]$ such that $\text{edim } \mathcal{I} = 0$. Furthermore, if $\vec{\lambda}$ satisfies the inequalities in Corollary 2.13 then $\mathcal{I} \in \text{Horn}(r, n, s)$. In Section 6 we will explain this more carefully and show how Belkale’s considerations allow us to construct a corresponding nonzero $\text{GL}(r)$ -invariant in $L(\lambda_1) \otimes \cdots \otimes L(\lambda_s)$. By Proposition 2.3, we will thus obtain at once a characterization of the Kirwan cone as well as of the existence of nonzero invariants in terms of Horn’s inequalities:

Corollary 6.3.3 (Knutson–Tao, [KT]). (a) ***Horn inequalities:** The Kirwan cone $\text{Kirwan}(r, s)$ is the convex polyhedral cone of $\vec{\xi} \in C_+(r)^s$ such that $\sum_{k=1}^s |\xi_k| = 0$, and for any $0 < d < r$ and any s -tuple $\mathcal{J} \in \text{Horn}(d, r, s)$ with $\text{edim } \mathcal{J} = 0$ we have that $\sum_{k=1}^s (T_{\mathcal{J}_k}, \xi_k) \leq 0$.*

(b) ***Saturation property:** For a dominant weight $\vec{\lambda} \in \Lambda_+(r)^s$, the space of invariants $(L(\lambda_1) \otimes \cdots \otimes L(\lambda_s))^{\text{GL}(r)}$ is nonzero if and only if $\vec{\lambda} \in \text{Kirwan}(r, s)$.*

In particular, $c(\vec{\lambda}) := \dim(L(\lambda_1) \otimes \cdots \otimes L(\lambda_s))^{\text{GL}(r)} > 0$ if and only if $c(N\vec{\lambda}) > 0$ for some integer $N > 0$.

The proof of Corollary 6.3.3 will be given in Section 6. In Appendices A and B, we list the Horn triples as well as the Horn inequalities for the Kirwan cones up to $r = 4$.

3. Subspaces, flags, positions

In this section, we study the geometry of subspaces and flags in more detail and supply proofs of some linear algebra facts used previously in Section 2.

3.1. Schubert positions. We start with some remarks on the Grassmannian $\text{Gr}(r, W)$, which is an irreducible algebraic variety on which the general linear group $\text{GL}(W)$ acts transitively. The stabilizer of a subspace $V \in \text{Gr}(r, W)$ is equal to the parabolic subgroup $P(V, W) = \{\gamma \in \text{GL}(W) : \gamma V \subseteq V\}$, with Lie algebra $\mathfrak{p}(V, W) = \{x \in \mathfrak{gl}(W) : xV \subseteq V\}$. Thus we obtain that

$$\text{Gr}(r, W) = \text{GL}(W) \cdot V \cong \text{GL}(W)/P(V, W),$$

and we can identify the tangent space at V with

$$T_V \text{Gr}(r, W) = \mathfrak{gl}(W) \cdot V \cong \mathfrak{gl}(W)/\mathfrak{p}(V, W) \cong \text{Hom}(V, W/V).$$

If we choose a complement Q of V in W then

$$(3.1.1) \quad \text{Hom}(V, Q) \rightarrow \text{Gr}(r, W), \quad \phi \mapsto (\text{id} + \phi)(V)$$

parametrizes a neighborhood of V . This gives a system of affine charts in $\text{Gr}(r, W)$ isomorphic to $\mathbb{C}^{r(n-r)}$. In particular, $\dim \text{Gr}(r, W) = r(n-r)$, a fact we use repeatedly in this article.

We now consider Schubert positions and the associated Schubert cells and varieties in more detail (Definitions 2.7 and 2.9) For all $\gamma \in \text{GL}(W)$, we have the following equivariance property:

$$(3.1.2) \quad \text{Pos}(\gamma^{-1}V, E) = \text{Pos}(V, \gamma E),$$

which in particular implies that

$$(3.1.3) \quad \gamma \Omega_I^0(E) = \Omega_I^0(\gamma E).$$

Thus $\Omega_I^0(E)$ is preserved by the Borel subgroup $B(E) = \{\gamma \in \text{GL}(W) : \gamma E(i) \subseteq E(i) \ (\forall i)\}$, which is the stabilizer of the flag E . We will see momentarily that $\Omega_I^0(E)$ is in fact a single $B(E)$ -orbit. We first state the following basic lemma, which shows that adapted bases (Definition 2.6) provide a convenient way of computing Schubert positions:

Lemma 3.1.4. *Let E be a flag on W , $\dim W = n$, $V \subseteq W$ an r -dimensional subspace, and $I \subseteq [n]$ a subset of cardinality r , with complement I^c . The following are equivalent:*

- (i) $\text{Pos}(V, E) = I$.
- (ii) *For any ordered basis $(f(1), \dots, f(n))$ adapted to E , there exists a (unique) basis $(v(1), \dots, v(r))$ of V of the form*

$$v(a) \in f(I(a)) + \text{span} \{f(i) : i \in I^c, i < I(a)\}.$$

(iii) *There exists an ordered basis $(f(1), \dots, f(n))$ adapted to E such that $\{f(I(1)), \dots, f(I(r))\}$ is a basis of V .*

The proof of Lemma 3.1.4 is left as an exercise to the reader. Clearly, $B(E)$ acts transitively on the set of ordered bases adapted to E . Thus, Lemma 3.1.4, (iii) shows that $\Omega_I^0(E)$ is a single $B(E)$ -orbit. That is, just like Grassmannian itself, each Schubert cell is a homogeneous space. In particular, $\Omega_I^0(E)$ and its closure $\Omega_I(E)$ (Definition 2.9) are both irreducible algebraic varieties.

Example. Consider the flag E on $W = \mathbb{C}^4$ with adapted basis $(f(1), \dots, f(4))$, where $f(1) = e(1) + e(2) + e(3)$, $f(2) = e(2) + e(3)$, $f(3) = e(3) + e(4)$, $f(4) = e(4)$. If $V = \text{span}\{e(1), e(2)\}$ then $\text{Pos}(V, E) = \{2, 4\}$, while $\text{Pos}(V, E_0) = \{1, 2\}$ for the standard flag E_0 with adapted basis $(e(1), e(2), e(3), e(4))$.

Note that the basis $(v(1), v(2))$ of V given by $v(1) = f(2) - f(1) = e(1)$ and $v(2) = f(4) - f(3) + f(1) = e(1) + e(2)$ satisfies the conditions in Lemma 3.1.4, (ii). It follows that $(f(1), v(1), f(3), v(2))$ is an adapted basis of E that satisfies the conditions in (iii).

The following lemma characterizes each Schubert variety explicitly as a union of Schubert cells:

Lemma 3.1.5. *Let E be a flag on W , $\dim W = n$, and $I \subseteq [n]$ a subset of cardinality r . Then,*

$$(3.1.6) \quad \Omega_I(E) = \bigcup_{I' \leq I} \Omega_{I'}^0(E),$$

where the union is over all subsets $I' \subseteq [n]$ of cardinality r such that $I'(a) \leq I(a)$ for $a \in [r]$.

Proof. Recall that $\Omega_I(E)$ can be defined as the Euclidean closure of $\Omega_I^0(E)$. Thus let (V_k) denote a convergent sequence of subspaces in $\Omega_I^0(E)$ with limit some $V \in \text{Gr}(r, W)$. Then $\dim E(I(a)) \cap V \geq \dim E(I(a)) \cap V_k$ for sufficiently large k , since intersections can only become larger in the limit, but $\dim E(I(a)) \cap V_k = a$ for all k . It follows that $\text{Pos}(V, E)(a) \leq I(a)$.

Conversely, suppose that $V' \in \Omega_{I'}^0(E)$, where $I'(a) \leq I(a)$ for all a . Let a' denote the minimal integer such that $I'(a) = I(a)$ for $a = a' + 1, \dots, r$. We will show that $V' \in \Omega_I(E)$ by induction on a' . If $a' = 0$ then $I' = I$ and there is nothing to show. Otherwise, let $(f'(1), \dots, f'(n))$ denote an adapted basis for E such that $v'(a) = f'(I'(a))$ is a basis of V' (as in (iii) of Lemma 3.1.4). For each $\varepsilon > 0$, consider the subspace V_ε with basis vectors $v_\varepsilon(a) = v'(a)$ for all $a \neq a'$ together with $v_\varepsilon(a') := v'(a') + \varepsilon f'(I(a'))$. Then the space V_ε is of dimension

r and in position $\{I'(1), \dots, I'(a' - 1), I(a'), \dots, I(r)\}$ with respect to E . By the induction hypothesis, $V_\varepsilon \in \Omega_I(E)$ for any $\varepsilon > 0$, and thus $V' \in \Omega_I(E)$ as $V_\varepsilon \rightarrow V'$ for $\varepsilon \rightarrow 0$. \square

We now compute the dimensions of Schubert cells and varieties. This is straightforward from Lemma 3.1.4, however it will be useful to make a slight detour and introduce some notation. This will allow us to show that we can exactly parametrize $\Omega_I^0(E)$ by a unipotent subgroup of $B(E)$, which in particular shows that it is an affine space.

Choose an ordered basis $(f(1), \dots, f(n))$ that is adapted to E . Then $V := \text{span}\{f(i) : i \in I\} \in \Omega_I^0(E)$. By Lemma 3.1.4, (ii) any $V \in \Omega_I^0(E)$ is of this form. Now define

$$\begin{aligned} \text{Hom}_E(V, W/V) &:= \{\phi \in \text{Hom}(V, W/V) : \phi(E(i) \cap V) \subseteq (E(i) + V)/V \text{ for } i \in [n]\} \\ &= \left\{ \phi \in \text{Hom}(V, W/V) : \phi\left(f(I(a))\right) \right. \\ &\quad \left. \subseteq \text{span}\{f(I^c(b)) + V : b \in [I(a) - a]\} \text{ for } a \in [r] \right\} \end{aligned}$$

where the $f(j) + V$ for $j \in I^c$ form a basis of W/V . In particular, $\text{Hom}_E(V, W/V)$ is of dimension $\sum_{a=1}^r (I(a) - a)$. Using this basis, we can identify W/V with $Q := \text{span}\{f(j) : j \in I^c\}$. Then $W = V \oplus Q$ and we can identify $\text{Hom}_E(V, W/V)$ with

$$\begin{aligned} H_E(V, Q) &:= \left\{ \phi \in \text{Hom}(V, Q) : \phi\left(f(I(a))\right) \right. \\ &\quad \left. \subseteq \text{span}\{f(I^c(b)) : b \in [I(a) - a]\} \text{ for } a \in [r] \right\}. \end{aligned}$$

Lemma 3.1.4, (ii) shows that for any $\phi \in H_E(V, Q)$, we obtain a distinct subspace $(\text{id} + \phi)(V)$ in $\Omega_I^0(E)$, and that all subspaces in $\Omega_I^0(E)$ can be obtained in this way. Thus, $\Omega_I^0(E)$ is contained in the affine chart $\text{Hom}(V, Q)$ of the Grassmannian described in (3.1.1) and isomorphic to the linear subspace $H_E(V, Q)$ of dimension $\dim I$. We define a corresponding unipotent subgroup,

$$U_E(V, Q) := \{u_\phi = \text{id} + \phi = \begin{pmatrix} \text{id}_V & 0 \\ \phi & \text{id}_Q \end{pmatrix} \in \text{GL}(W) : \phi \in H_E(V, Q)\}.$$

Thus we obtain the following lemma:

Lemma 3.1.7. *Let E be a flag on W , $\dim W = n$, $I \subseteq [n]$ a subset of cardinality r , $V \in \Omega_I^0(E)$, and Q as above. Then we can parametrize $H_E(V, Q) \cong U_E(V, Q) \cong \Omega_I^0(E) = U_E(V, Q)V$, hence $H_E(V, Q) \cong T_V \Omega_I^0(E)$ and*

$$(3.1.8) \quad \dim \Omega_I^0(E) = \dim \Omega_I(E) = \dim H_E(V, Q) = \sum_{a=1}^r (I(a) - a) =: \dim I.$$

It will be useful to rephrase the above to obtain a parametrization of $\Omega_I^0(E)$ in terms of the fixed subspaces

$$(3.1.9) \quad \begin{aligned} V_0 &:= \text{span}\{f(1), \dots, f(r)\} = E(r), \\ Q_0 &:= \text{span}\{\bar{f}(1), \dots, \bar{f}(n-r)\}, \end{aligned}$$

where the $\bar{f}(i) := f(r+i)$ for $i \in [n-r]$ form a basis of Q_0 . Then $W = V_0 \oplus Q_0$.

Definition 3.1.10. Let $I \subseteq [n]$ be a subset of cardinality r . The *shuffle permutation* $\sigma_I \in S_n$ is defined by

$$\sigma_I(a) = \begin{cases} I(a) & \text{for } a = 1, \dots, r, \\ I^c(a-r) & \text{for } a = r+1, \dots, n. \end{cases}$$

and $w_I \in \text{GL}(W)$ is the corresponding permutation operator with respect to the adapted basis $(f(1), \dots, f(n))$, defined as $w_I f(i) := f(\sigma_I^{-1}(i))$ for $i \in [n]$.

Then $V_0 = w_I V$, where $V = \text{span}\{f(i) : i \in I\} \in \Omega_I^0(E)$ as before, and so

$$V_0 \in w_I \Omega_I^0(E) = \Omega_I^0(w_I E)$$

using (3.1.3). The translated Schubert cell can be parametrized by

$$\begin{aligned} &H_{w_I E}(V_0, Q_0) \\ &= \left\{ \phi \in \text{Hom}(V_0, Q_0) : \phi(f(a)) \subseteq \text{span}\{\bar{f}(1), \dots, \bar{f}(I(a)-a)\} \text{ for } a \in [r] \right\}, \end{aligned}$$

where we identify $Q_0 \cong W/V_0$. We thus obtain the following consequence of Lemma 3.1.7:

Corollary 3.1.11. *Let E be a flag on W , $\dim W = n$, $I \subseteq [n]$ of cardinality r , and $V \in \Omega_I^0(E)$. Moreover, define w_I as above for an adapted basis. Then,*

$$\Omega_I^0(E) = w_I^{-1} \Omega_I^0(w_I E) = w_I^{-1} U_{w_I E}(V_0, Q_0) V_0.$$

Example ($r = 3, n = 4$). Let $I = \{1, 3, 4\}$ and E_0 the standard flag on $W = \mathbb{C}^4$, with its adapted basis $(e(1), \dots, e(4))$. Then $\sigma_I = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 3 & 4 & 2 \end{pmatrix}$,

$$w_I^{-1} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

and $V = w_I^{-1}V_0 = \text{span}\{e(1), e(3), e(4)\}$ is indeed in position I with respect to E_0 , in agreement with the preceding discussion. Moreover,

$$H_{w_I E_0}(V_0, Q_0) = \{(0 \quad * \quad *)\} \subseteq \text{Hom}(\mathbb{C}^3, \mathbb{C}^1),$$

$$U_{w_I E_0}(V_0, Q_0) = \left\{ \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & * & * & 1 \end{pmatrix} \right\} \subseteq \text{GL}(4),$$

and so Corollary 3.1.11 asserts that

$$\Omega_I^0(E_0) = w_I^{-1}U_{w_I E_0}(V_0, Q_0) \text{span}\{e(1), e(2), e(3)\} = \text{span} \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ * \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ * \\ 0 \\ 1 \end{pmatrix} \right\},$$

which agrees with Lemma 3.1.4.

3.2. Induced flags and positions. The space $\text{Hom}_E(V, W/V)$ can be understood more conceptually as the space of homomorphisms that respect the filtrations $E(i) \cap V$ and $(E(i) + V)/V$ induced by the flag E . Here we have used the following concept:

Definition 3.2.1. A (complete) filtration F on a vector space V is a chain of subspaces

$$\{0\} = F(0) \subseteq F(1) \subseteq \dots \subseteq F(i) \subseteq F(i+1) \subseteq \dots \subseteq F(l) = V,$$

such that the dimensions increase by no more than one, i.e., $\dim F(i+1) \leq \dim F(i) + 1$ for all $i = 0, \dots, l-1$. Thus distinct subspaces in a filtration determine a flag.

Given a flag E on W and a subspace $V \subseteq W$, we thus obtain an induced flag E^V on V from the distinct subspaces in the sequence $E(i) \cap V$, $i = 0, \dots, n$. We

may also induce a flag $E_{W/V}$ on the quotient W/V from the distinct subspaces in the sequence $(E(i) + V)/V$. These flags can be readily computed from the Schubert position of V :

Lemma 3.2.2. *Let E be a flag on W , $\dim W = n$, and $V \subseteq W$ an r -dimensional subspace in position $I = \text{Pos}(V, E)$. Then the induced flags E^V on V and $E_{W/V}$ on W/V are given by*

$$E^V(a) = E(I(a)) \cap V,$$

$$E_{W/V}(b) = (E(I^c(b)) + V)/V$$

for $a \in [r]$ and $b \in [n - r]$, where I^c denotes the complement of I in $[n]$.

Proof. Using an adapted basis as in Lemma 3.1.4, (iii), it is easy to see that $\dim E(i) \cap V = |[i] \cap I|$ and therefore that $\dim(E(i) + V)/V = |[i] \cap I^c|$. Now observe that $|[i] \cap I| = a$ if and only if $I(a) \leq i < I(a + 1)$, while $|[i] \cap I^c| = b$ if and only if $I^c(b) \leq i < I^c(b + 1)$. Thus we obtain the two assertions. \square

We can use the preceding result to describe $\text{Hom}_E(V, W/V)$ in terms of flags rather than filtrations and without any reference to the ambient space W .

Definition 3.2.3. Let V and Q be vector spaces of dimension r and $n - r$, respectively, $I \subseteq [n]$ a subset of cardinality r , F a flag on V and G a flag on Q . We define

$$H_I(F, G) := \{\phi \in \text{Hom}(V, Q) : \phi(F(a)) \subseteq G(I(a) - a)\},$$

which we note is well-defined by

$$(3.2.4) \quad 0 \leq I(a) - a \leq I(a + 1) - (a + 1) \leq n - r \quad (a = 1, \dots, r - 1).$$

It now easily follows from Lemmas 3.1.7 and 3.2.2 that

$$(3.2.5) \quad T_V \Omega_I^0(E) \cong \text{Hom}_E(V, W/V) = H_I(E^V, E_{W/V}).$$

As a consequence:

$$(3.2.6) \quad H_{w_I E}(V_0, Q_0) = H_I((w_I E)^{V_0}, (w_I E)_{Q_0}) = H_I(E^{V_0}, E_{Q_0})$$

We record the following equivariance property:

Lemma 3.2.7. *Let F be a flag on V and G a flag on Q . If $\phi \in H_I(F, G)$, $a \in GL(V)$ and $d \in GL(Q)$, then $d\phi a^{-1} \in H_I(aF, dG)$. In particular, $H_I(F, G)$ is stable under right multiplication by the Borel subgroup $B(F)$ and left multiplication by the Borel subgroup $B(G)$.*

We now compute the position of subspaces and subquotients with respect to induced flags. Given subsets $I \subseteq [n]$ of cardinality r and $J \subseteq [r]$ of cardinality d , we recall that we had defined their *composition* IJ in Section 2 as the subset

$$IJ = \{I(J(1)) < \cdots < I(J(d))\} \subseteq [n].$$

We also define their *quotient* to be the subset

$$I/J = \{I(J^c(b)) - J^c(b) + b : b \in [r - d]\} \subseteq [n - d],$$

where J^c denotes the complement of J in $[r]$. It follows from (3.2.4) that I/J is indeed a subset of $[n - d]$.

The following lemma establishes the ‘chain rule’ for positions:

Lemma 3.2.8. *Let E be a flag on W , $S \subseteq V \subseteq W$ subspaces, and $I = \text{Pos}(V, E)$, $J = \text{Pos}(S, E^V)$ their relative positions. Then there exists an adapted basis $(f(1), \dots, f(n))$ for E such that $\{f(I(a))\}$ is a basis of V and $\{f(IJ(b))\}$ a basis of S . In particular,*

$$(3.2.9) \quad \text{Pos}(S, E) = IJ = \text{Pos}(V, E) \text{Pos}(S, E^V).$$

Proof. According to Lemma 3.1.4, (iii), there exists an adapted basis $(f(1), \dots, f(n))$ for E such that $(f(I(1)), \dots, f(I(r)))$ is a basis of V , where $r = \dim V$. By Lemma 3.2.2, this ordered basis is in fact adapted to the induced flag E^V . Thus we can apply Lemma 3.1.4, (ii) to E^V and the subspace $S \subseteq V$ to obtain a basis $(v(1), \dots, v(s))$ of S of the form

$$v(b) \in f(IJ(b)) + \text{span} \{f(I(a)) : a \in J^c, a < J(b)\}.$$

It follows that the ordered basis $(f'(1), \dots, f'(n))$ obtained from $(f(1), \dots, f(n))$ by replacing $f(IJ(b))$ with $v(b)$ has all desired properties. We now obtain the chain rule, $\text{Pos}(S, E) = IJ$, as a consequence of Lemma 3.1.4, (iii) applied to f' and $S \subseteq W$. \square

We can visualize the subsets $IJ, IJ^c \subseteq [n]$ and $I/J \subseteq [n - d]$ as follows. Let L denote the string of length n defined by putting the symbol s at the positions in IJ , v at those in $I \setminus IJ = IJ^c$, and w at all other positions. This mirrors the situation in the preceding Lemma 3.2.8, where the adapted basis $(f(1), \dots, f(n))$ can be partitioned into three sets according to membership in S , $V \setminus S$, and $W \setminus V$. Now let L' denote the string of length $n - d$ obtained by deleting all occurrences of the symbol s . Thus the remaining symbols are either v or w , i.e., those that were at locations $(IJ)^c$ in L . We observe that the b -th occurrence of v in L was at location $IJ^c(b)$, where it was preceded by $J^c(b) - b$ occurrences of s . Thus the occurrences of v in L' are given precisely by the quotient position, $(I/J)(b) = IJ^c(b) - (J^c(b) - b)$.

Example. If $n = 6$, $I = \{1, 3, 5, 6\}$ and $J = \{2, 4\}$, then $IJ = \{3, 6\}$ and $L = (v, w, s, w, v, s)$. It follows that $L' = (v, w, w, v)$ and hence the symbols v appear indeed at positions $I/J = \{1, 4\}$.

We thus obtain the following recipe for computing positions of subquotients:

Lemma 3.2.10. *Let E be a flag on W and $S \subseteq V \subseteq W$ subspaces. Then,*

$$\text{Pos}(V/S, E_{W/S}) = \text{Pos}(V, E) / \text{Pos}(S, E^V).$$

Proof. Let $I = \text{Pos}(V, E)$ and $J = \text{Pos}(S, E^V)$. According to Lemma 3.2.8, there exists an adapted basis $(f(1), \dots, f(n))$ of E such that $\{f(I(a))\}$ is a basis of V and $\{f(IJ(b))\}$ a basis of S . This shows not only that $\{f(IJ^c(b))\}$ is a basis of V/S , but also, by Lemma 3.2.2, that $(f((IJ)^c(b)))$ is an adapted basis for $E_{W/S}$. Clearly, $IJ^c \subseteq (IJ)^c$, and the preceding discussion showed that the location of the IJ^c in $(IJ)^c$ is exactly equal to the quotient position I/J . Thus we conclude from Lemma 3.1.4, (iii) that $\text{Pos}(V/S, E_{W/S}) = I/J$. \square

One last consequence of the preceding discussion is the following lemma:

Lemma 3.2.11. *Let E be a flag on W , $\dim W = n$, $S \subseteq V \subseteq W$ subspaces, and $I = \text{Pos}(V, E)$, $J = \text{Pos}(S, E^V)$. Then $F(i) := ((E(i) \cap V) + S)/S$ is a filtration on V/S , and*

$$IJ^c(b) = \min \{i \in [n] : \dim F(i) = b\}$$

for $b = 1, \dots, \dim V/S$.

Proof. As in the preceding proof, we use the adapted basis $(f(1), \dots, f(n))$ from Lemma 3.2.8. Then $\{f(IJ^c(b))\}$ is a basis of V/S and $F(i) = \text{span}\{f(IJ^c(b)) : b \in [q], IJ^c(b) \leq i\}$, and this implies the claim. \square

The following corollary uses Lemma 3.2.11 to compare filtrations for a space that is isomorphic to a subquotient in two different ways, $(S_1 + S_2)/S_2 \cong S_1/(S_1 \cap S_2)$.

Corollary 3.2.12. *Let E be a flag on W , $\dim W = n$, and $S_1, S_2 \subseteq W$ subspaces. Furthermore, let $J = \text{Pos}(S_1, E)$, $K = \text{Pos}(S_1 \cap S_2, E^{S_1})$, $L = \text{Pos}(S_1 + S_2, E)$, and $M = \text{Pos}(S_2, E^{S_1+S_2})$. Then both JK^c and LM^c are subsets of $[n]$ of cardinality $q := \dim S_1/(S_1 \cap S_2) = \dim(S_1 + S_2)/S_2$, and*

$$JK^c(b) \leq LM^c(b)$$

for $b \in [q]$.

Proof. Consider the filtration $F(j) := ((E(j) \cap S_1) + (S_1 \cap S_2))/(S_1 \cap S_2)$ of $S_1/(S_1 \cap S_2)$ and the filtration $F'(j) := ((E(j) \cap (S_1 + S_2)) + S_2)/S_2$ of $(S_1 + S_2)/S_2$. If we identify $S_1/(S_1 \cap S_2) \cong (S_1 + S_2)/S_2$, then $F(j)$ gets identified with the subspace $((E(j) \cap S_1) + S_2)/S_2$ of $F'(j)$. It follows that

$$\begin{aligned} JK^c(b) &= \min \{j \in [n] : \dim F(j) = b\} \geq \min \{j \in [n] : \dim F'(j) = b\} \\ &= LM^c(b), \end{aligned}$$

where we have used Lemma 3.2.11 twice. \square

We now compute the dimension of quotient positions:

Lemma 3.2.13. *Let $I \subseteq [n]$ be a subset of cardinality r and $J \subseteq [r]$ a subset of cardinality d . Then:*

$$\dim I/J = \dim I + \dim J - \dim IJ$$

Proof. Straight from the definition of dimension and quotient position,

$$\begin{aligned} \dim I/J &= \sum_{b=1}^{r-d} I(J^c(b)) - \sum_{b=1}^{r-d} J^c(b) \\ &= \left(\sum_{a=1}^r I(a) - \sum_{b=1}^d I(J(b)) \right) - \left(\sum_{a=1}^r a - \sum_{b=1}^d J(b) \right) \\ &= \sum_{a=1}^r (I(a) - a) + \sum_{b=1}^d (J(b) - b) - \sum_{b=1}^d (I(J(b)) - b) \\ &= \dim I + \dim J - \dim IJ. \end{aligned}$$

\square

Lastly, given subsets $I \subseteq [n]$ of cardinality r and $J \subseteq [r]$ of cardinality d , we define

$$I^J = \{I(J(b)) - J(b) + b : b \in [d]\} \subseteq [n - (r - d)].$$

Clearly, $I^J = I/J^c$, but we prefer to introduce a new notation to avoid confusion, since the role of I^J will be quite different. Indeed, I^J is related to composition, as is indicated by the following lemmas:

Lemma 3.2.14. *Let $I \subseteq [n]$ be a subset of cardinality r , $J \subseteq [r]$ a subset of cardinality d . Then,*

$$\dim I^J K - \dim K = \dim I(JK) - \dim JK$$

for any subset $K \subseteq [d]$. In particular, $\dim I^J = \dim IJ - \dim J$.

Proof. Let m denote the cardinality of K . Then:

$$\begin{aligned} \dim I^J K - \dim K &= \sum_{c=1}^m \left(I^J(K(c)) - K(c) \right) \\ &= \sum_{c=1}^m \left(I \left(J(K(c)) - J(K(c)) \right) \right) = \dim I(JK) - \dim JK. \end{aligned}$$

□

Lemma 3.2.15. *Let $I \subseteq [n]$ be a subset of cardinality r , $\phi \in \text{Hom}(V, Q)$, and F a flag on V . Let $S = \ker \phi$ denote the kernel, $J := \text{Pos}(S, F)$ its position with respect to F , and $\bar{\phi} \in \text{Hom}(V/S, Q)$ the corresponding injection.*

Then $\phi \in H_I(F, G)$ if and only if $\bar{\phi} \in H_{I/J}(F_{V/S}, G)$. In this case, we have for all $\psi \in H_J(F^S, F_{V/S})$ that $\bar{\phi}\psi \in H_{I^J}(F^S, G)$.

Proof. For the first claim, note that if $\phi \in H_I(F, G)$ then

$$\bar{\phi}(F_{V/S}(b)) = \phi(F(J^c(b))) \subseteq G(I(J^c(b)) - J^c(b)) = G((I/J)(b) - b).$$

Conversely, if $\bar{\phi} \in H_{I/J}(F_{V/S}, G)$, then this shows that

$$\phi(F(a)) \subseteq G(I(a) - a)$$

for all $a = J^c(b)$, and hence for all a , since $\phi(F(J^c(b))) = \dots = \phi(F(J^c(b + 1) - 1))$.

For the second, we use $H_J(F^S, F_{V/S}) = \text{Hom}_F(S, V/S)$ (Eq. (3.2.5)) and compute

$$\begin{aligned} \bar{\phi}\psi(F^S(a)) &= \bar{\phi}\psi(F(J(a)) \cap S) \subseteq \bar{\phi}\left(\left(F(J(a)) + S\right)/S\right) \\ &= \phi(F(J(a))) \subseteq G(I(J(a)) - J(a)) = G(I^J(a) - a). \end{aligned}$$

□

3.3. The flag variety. The Schubert cells of the Grassmannian were defined by fixing a flag and classifying subspaces according to their Schubert position. As we will later be interested in intersections of Schubert cells for different flags, it will be useful to also consider variations of the flag for a fixed subspace.

Let $\text{Flag}(W)$ denote the (complete) flag variety, defined as the space of (complete) flags on W . It is a homogeneous space with respect to the transitive $\text{GL}(W)$ -action, so indeed an irreducible variety.

Definition 3.3.1. Let $V \subseteq W$ be a subspace, $\dim V = r$, $\dim W = n$, and $I \subseteq [n]$ a subset of cardinality r . We define

$$\text{Flag}_I^0(V, W) = \{E \in \text{Flag}(W) : \text{Pos}(V, E) = I\},$$

and $\text{Flag}_I(V, W)$ as its closure in $\text{Flag}(W)$ (in either the Euclidean or the Zariski topology).

We have the following equivariance property as a consequence of (3.1.2): For all $\gamma \in \text{GL}(W)$,

$$(3.3.2) \quad \gamma \text{Flag}_I^0(V, W) = \text{Flag}_I^0(\gamma V, W).$$

In particular, $\text{Flag}_I^0(V, W)$ and $\text{Flag}_I(V, W)$ are stable under the action of the parabolic subgroup $P(V, W) = \{\gamma \in \text{GL}(W) : \gamma V \subseteq V\}$, which is the stabilizer of V .

We will now show that $\text{Flag}_I^0(V, W)$ is in fact a single $P(V, W)$ -orbit. This implies that both $\text{Flag}_I^0(V, W)$ and $\text{Flag}_I(V, W)$ are irreducible algebraic varieties.

Definition 3.3.3. Let E be a flag on W , $\dim W = n$, $V_0 = E(r)$, and $I \subseteq [n]$ a subset of cardinality r . We define

$$G_I(V_0, E) := \{\gamma \in \text{GL}(W) : \gamma E \in \text{Flag}_I^0(V_0, W)\},$$

so that $\text{Flag}_I^0(V_0, W) \cong G_I(V_0, E)/B(E)$.

Lemma 3.3.4. Let E be a flag on W , $\dim W = n$, $V_0 = E(r)$, and $I \subseteq [n]$ a subset of cardinality r . Then, $G_I(V_0, E) = P(V_0, W)w_I B(E)$. In particular, $\text{Flag}_I^0(V_0, W) = P(V_0, W)w_I E$.

Proof. Let $\gamma \in \text{GL}(W)$. Then,

$$\gamma \in G_I(V_0, E) \Leftrightarrow V_0 \in \Omega_I^0(\gamma E) = \gamma \Omega_I^0(E) = \gamma B(E)w_I^{-1}V_0 \Leftrightarrow \gamma \in P(V_0, W)w_I B(E),$$

where we have used that $\Omega_I^0(E) = B(E)w_I^{-1}V_0$. □

We now derive a more precise parametrization of $\text{Flag}_I^0(V_0, W)$.

Lemma 3.3.5. Let E be a flag on W , $\dim W = n$, V_0 and Q_0 as in (3.1.9), and $I \subseteq [n]$ a subset of cardinality r . Then we have that $G_I(V_0, E) = P(V_0, W)U_{w_I E}(V_0, Q_0)w_I$.

Proof. Let $\gamma \in \text{GL}(W)$. Then,

$$\begin{aligned} \gamma \in G_I(V_0, E) &\Leftrightarrow V_0 \in \Omega_I^0(\gamma E) = \gamma \Omega_I^0(E) = \gamma w_I^{-1}U_{w_I E}(V_0, Q_0)V_0 \\ &\Leftrightarrow \gamma \in P(V_0, W)U_{w_I E}(V_0, Q_0)w_I, \end{aligned}$$

since $\Omega_I^0(E) = w_I^{-1}U_{w_I E}(V_0, Q_0)V_0$ (Corollary 3.1.11). □

In coordinates, using $W = V_0 \oplus Q_0$ and (3.2.6), we obtain that

$$\begin{aligned} G_I(V_0, E) &= \left\{ \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \phi & 1 \end{pmatrix} : a \in \text{GL}(V_0), d \in \text{GL}(Q_0), \phi \in H_I(E^{V_0}, E_{Q_0}) \right\} w_I \\ &= \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a - bd^{-1}c \in \text{GL}(V_0), d \in \text{GL}(Q_0), d^{-1}c \in H_I(E^{V_0}, E_{Q_0}) \right\} w_I. \end{aligned}$$

In particular, $\dim G_I(V_0, E) = \dim P(V_0, W) + \dim I$. This allows us to compute the dimension of the subvarieties $\text{Flag}_I^0(V, W)$ and to relate their codimension to the codimension of the Schubert cells of the Grassmannian:

Corollary 3.3.6. *Let $V \subseteq W$ be a subspace, $\dim W = n$, $\dim V = r$, and $I \subseteq [n]$ a subset of cardinality r . Then,*

$$(3.3.7) \quad \begin{aligned} \dim \text{Flag}_I^0(V, W) &= \dim \text{Flag}_I(V, W) \\ &= \dim \text{Flag}(V) + \dim \text{Flag}(Q) + \dim I \end{aligned}$$

and

$$(3.3.8) \quad \dim \text{Flag}(W) - \dim \text{Flag}_I^0(V, W) = \dim \text{Gr}(r, W) - \dim I.$$

Proof. Without loss of generality, we may assume that $V = V_0 = E(r)$ for some flag E on W . Then, $\text{Flag}_I^0(V_0, W) \cong G_I(V_0, E)/B(E)$ and hence

$$\begin{aligned} \dim \text{Flag}_I^0(V_0, E) &= \dim P(V_0, W) + \dim I - \dim B(E) \\ &= \dim \text{GL}(W) - \dim \text{Gr}(r, W) + \dim I - \dim B(E) \\ &= \dim \text{Flag}(W) - \dim \text{Gr}(r, W) + \dim I \end{aligned}$$

since $\text{Gr}(r, W) \cong \text{GL}(W)/P(V_0, W)$ and $\text{Flag}(W) = \text{GL}(W)/B(E)$. This establishes (3.3.8). On the other hand, a direct calculation shows that

$$\dim \text{Flag}(W) - \dim \text{Gr}(r, W) = \dim \text{Flag}(V) + \dim \text{Flag}(Q),$$

so we also obtain (3.3.7). □

At last, we study the following set of flags on the target space of a given homomorphism:

Definition 3.3.9. Let V, Q be vector spaces of dimension r and $n - r$, respectively, and $I \subseteq [n]$. Moreover, let F be a flag on V and $\phi \in \text{Hom}(V, Q)$ an *injective* homomorphism. We define

$$\text{Flag}_I^0(F, \phi) := \{G \in \text{Flag}(Q) : \phi \in H_I(F, G)\}$$

where we recall that $H_I(F, G)$ was defined in Definition 3.2.3.

It is clear that $I(a) \geq 2a$ is necessary and sufficient for $\text{Flag}_I^0(F, \phi)$ to be nonempty.

Example ($r=3, n=8$). Let $V_0 \cong \mathbb{C}^3$, with basis $e(1), \dots, e(3)$, and $Q_0 \cong \mathbb{C}^5$, with basis $\bar{e}(1), \dots, \bar{e}(5)$. Take $\phi: V_0 \rightarrow Q_0$ to be the canonical injection and let F_0 denote the standard flag on V_0 . For $I = \{3, 4, 7\}$, $G \in \text{Flag}_I^0(F_0, \phi)$ if and only if

$$\mathbb{C}\bar{e}(1) \subseteq G(2), \quad \mathbb{C}\bar{e}(1) \oplus \mathbb{C}\bar{e}(2) \subseteq G(2), \quad \mathbb{C}\bar{e}(1) \oplus \mathbb{C}\bar{e}(2) \oplus \mathbb{C}\bar{e}(3) \subseteq G(4).$$

For example, the standard flag G_0 on Q_0 is a point in $\text{Flag}_I^0(F_0, \phi)$.

On the other hand, if $I = \{2, 3, 7\}$ then we obtain the condition $\mathbb{C}\bar{e}(1) \oplus \mathbb{C}\bar{e}(2) \subseteq G(1)$ which can never be satisfied. Thus in this case $\text{Flag}_I^0(F_0, \phi) = \emptyset$.

In the following lemma we show that $\text{Flag}_I^0(F, \phi)$ is a smooth variety and compute its dimension.

Lemma 3.3.10. *Let V, Q be vector spaces of dimension r and $n-r$, respectively, and $I \subseteq [n]$ a subset of cardinality r . Moreover, let F be a flag on V and $\phi \in \text{Hom}(V, Q)$ an **injective** homomorphism. If $\text{Flag}_I^0(F, \phi)$ is nonempty, that is, if $I(a) \geq 2a$ for all $a \in [r]$, then it is a smooth irreducible subvariety of $\text{Flag}(Q)$ of dimension*

$$\dim \text{Flag}_I^0(F, \phi) = \dim \text{Flag}(Q_0) + \dim I - r(n - r).$$

Proof. Without loss of generality, we may assume that $V = V_0 \cong \mathbb{C}^r$, $Q = Q_0 \cong \mathbb{C}^{n-r}$, that $F = F_0$ is the standard flag on V_0 and ϕ the canonical injection $\mathbb{C}^r \rightarrow \mathbb{C}^{n-r}$. Then the standard flag G_0 on Q_0 is an element of $\text{Flag}_I^0(F_0, \phi)$. We will show that

$$M_I := \{h \in \text{GL}(Q_0) : h G_0 \in \text{Flag}_I^0(F_0, \phi)\}$$

is a subvariety of $\text{GL}(Q_0)$ and compute its dimension. Note that $h \in M_I$ if and only if $h^{-1}\phi \in H_I(F_0, G_0)$. We now identify V_0 with its image $\phi(V_0)$ and denote by $R_0 \cong \mathbb{C}^{n-2r}$ its standard complement in Q_0 . Thus $Q_0 \cong V_0 \oplus R_0$ and we can think of $h^{-1} \in \text{GL}(Q_0)$ as a block matrix

$$h^{-1} = \begin{pmatrix} A & B \end{pmatrix}$$

where $A \in \text{Hom}(V_0, Q_0)$ and $B \in \text{Hom}(R_0, Q_0)$. The condition $h^{-1}\phi \in H_I(F_0, G_0)$ amounts to demanding that $A \in H_I(F_0, G_0)$, while B is unconstrained. Thus we can identify M_I via $h \mapsto h^{-1}$ with the invertible elements in

$$H_I(F_0, G_0) \times \text{Hom}(R_0, Q_0),$$

which form a nonempty Zariski-open subset, and hence a smooth irreducible subvariety of $\text{GL}(Q_0)$. It follows that $\text{Flag}_I^0(F_0, \phi) = M_I/B(G_0)$ is likewise a smooth irreducible subvariety, and

$$\begin{aligned} \text{Flag}_I^0(F_0, \phi) &= \dim M_I - \dim B(G_0) = \dim I + (n - r)(n - 2r) - \dim B(G_0) \\ &= \dim \text{Flag}(Q_0) + \dim I - (n - r)r, \end{aligned}$$

where we have used Eq. (3.1.8) and that $\text{Flag}(Q_0) \cong \text{GL}(Q_0)/B(G_0)$. □

4. Intersections and Horn inequalities

In this section, we study intersections of Schubert varieties. Recall from Definition 2.12 that given an s -tuple \mathcal{E} of flags on W , $\dim W = n$, and $\mathcal{I} \in \text{Subsets}(r, n, s)$, we had defined

$$\Omega_{\mathcal{I}}^0(\mathcal{E}) = \bigcap_{k=1}^s \Omega_{I_k}^0(E_k) \quad \text{and} \quad \Omega_{\mathcal{I}}(\mathcal{E}) = \bigcap_{k=1}^s \Omega_{I_k}(E_k).$$

We are particularly interested in the intersecting \mathcal{I} , denoted $\mathcal{I} \in \text{Intersecting}(r, n, s)$, for which $\Omega_{\mathcal{I}}(\mathcal{E}) \neq \emptyset$ for every \mathcal{E} .

4.1. Coordinates. Without loss of generality, we may assume that $W = \mathbb{C}^n$, and we shall do so for the remainder of this article. As before, we denote by $(e(1), \dots, e(n))$ the ordered standard basis of \mathbb{C}^n and by E_0 the corresponding *standard flag*. Let $V_0 = E_0(r)$ be the standard r -dimensional subspace, with ordered basis $(e(1), \dots, e(r))$, and Q_0 the subspace with ordered basis $(\bar{e}(1), \dots, \bar{e}(n-r))$, where $\bar{e}(b) := e(r+b)$. Thus $W = V_0 \oplus Q_0$. We denote the corresponding standard flags on V_0 and Q_0 by F_0 and G_0 , respectively. Note that $F_0 = E_0^{V_0}$ and, if we identify $Q_0 \cong W/V_0$, then $G_0 = (E_0)_{W/V_0}$. We further abbreviate the Grassmannian by $\text{Gr}(r, n) := \text{Gr}(r, \mathbb{C}^n)$, the parabolic by $P(r, n) := P(V_0, \mathbb{C}^n)$ and the Borel by $B(n) := B(E_0)$. We write $\text{Flag}(n) := \text{Flag}(W)$ and $\text{Flag}_I^0(r, n) := \text{Flag}_I^0(V_0, W)$ for the set of flags with respect to which V_0 has position I ; $\text{Flag}_I(r, n) := \text{Flag}_I(V_0, W)$ is its closure. We recall from Definition 3.2.3 that

$$H_I(F_0, G_0) = \left\{ \phi \in \text{Hom}(V_0, Q_0) : \phi(e(a)) \subseteq \text{span} \{ \bar{e}(1), \dots, \bar{e}(I(a) - a) \} \right\},$$

and Lemma 3.3.5 reads

$$(4.1.1) \quad G_I(r, n) = P(r, n) \left\{ \begin{pmatrix} \text{id}_{V_0} & 0 \\ \phi & \text{id}_{Q_0} \end{pmatrix} : \phi \in H_I(F_0, G_0) \right\} w_I,$$

where we have introduced $G_I(r, n) := G_I(V_0, E_0)$.

4.2. Intersections and dominance. We start by reformulating the intersecting property in terms of the dominance of certain morphisms of algebraic varieties. This allows us to give a simple proof of Lemma 4.2.6, which states that the expected dimension of an intersecting tuple is necessarily nonnegative.

We caution that while $\Omega_{\mathcal{I}}^0(\mathcal{E}) \subseteq \Omega_{\mathcal{I}}(\mathcal{E})$, the latter is *not* necessarily the closure of the former:

Example. Let $W = \mathbb{C}^2$, $I_1 = \{1\}$, $I_2 = \{2\}$, and $E_1 = E_2$ the same flag on W . Since the Schubert cells $\Omega_{I_k}^0(\mathcal{E})$ partition the projective space $\mathbb{P}(W) = \text{Gr}(1, W)$, $\Omega_{\mathcal{I}}^0(\mathcal{E}) = \emptyset$ is empty, but $\Omega_{\mathcal{I}}(\mathcal{E}) = \{E_1(1)\}$ is a point.

It is also possible that $\Omega_{\mathcal{I}}^0(\mathcal{E})$ or $\Omega_{\mathcal{I}}(\mathcal{E})$ are nonempty for some \mathcal{E} but empty for generic s -tuples \mathcal{E} :

Example. Let $W = \mathbb{C}^2$, $I_1 = I_2 = \{1\}$. Then $\Omega_{\mathcal{I}}^0(\mathcal{E}) = \Omega_{\mathcal{I}}(\mathcal{E}) = E_1(1) \cap E_2(1)$, so the intersection is nonempty if and only if $E_1 = E_2$.

We will later show the existence of a ‘good set’ of sufficiently generic \mathcal{E} such that \mathcal{I} is intersecting if and only if $\Omega_{\mathcal{I}}^0(\mathcal{E}) \neq \emptyset$ for any single ‘good’ \mathcal{E} (Lemma 4.3.1). Here is a more interesting example:

Example 4.2.1. Let $W = \mathbb{C}^6$, $s = 3$, and $\mathcal{I} = (I_1, I_2, I_3)$ where all $I_k = \{2, 4, 6\}$. The triple \mathcal{I} is intersecting. Let

$$f(t) := e_1 + te_2 + \frac{t^2}{2!}e_3 + \frac{t^3}{3!}e_4 + \frac{t^4}{4!}e_5 + \frac{t^5}{5!}e_6$$

and consider the one-parameter family of flags $E(t)$ with adapted basis $(f(t), \frac{d}{dt}f(t), \dots, \frac{d^5}{dt^5}f(t))$. We consider the 3-tuple $\mathcal{E} = (E_1, E_2, E_3)$, where $E_1 := E(0)$ is the standard flag, $E_2 := E(1)$, and $E_3 := E(-1)$. Then the intersection $\Omega_{\mathcal{I}}^0(\mathcal{E})$ consists of precisely two points:

$$V_1 = \text{span} \left\{ e_2 + \sqrt{5}e_1, e_4 - 24\sqrt{5}e_1 - 3\sqrt{5}e_3, e_6 - 24\sqrt{5}e_3 + \sqrt{5}e_5 \right\},$$

$$V_2 = \text{span} \left\{ e_2 - \sqrt{5}e_1, e_4 + 24\sqrt{5}e_1 + 3\sqrt{5}e_3, e_6 + 24\sqrt{5}e_3 - \sqrt{5}e_5 \right\},$$

and coincides with $\Omega_{\mathcal{I}}(\mathcal{E})$.

To study generic intersections of Schubert cells, it is useful to introduce the following maps: Let $\mathcal{I} \in \text{Subsets}(r, n, s)$. We define

$$\omega_{\mathcal{I}}^0: \begin{cases} \text{GL}(n) \times \text{Flag}_{I_1}^0(r, n) \times \dots \times \text{Flag}_{I_s}^0(r, n) \rightarrow \text{Flag}(n)^s \\ (\gamma, E_1, \dots, E_s) \mapsto (\gamma E_1, \dots, \gamma E_s) \end{cases}$$

and its extension

$$(4.2.2) \quad \omega_{\mathcal{I}}: \begin{cases} \mathrm{GL}(n) \times \mathrm{Flag}_{I_1}(r, n) \times \cdots \times \mathrm{Flag}_{I_s}(r, n) \rightarrow \mathrm{Flag}(n)^s \\ (\gamma, E_1, \dots, E_s) \mapsto (\gamma E_1, \dots, \gamma E_s). \end{cases}$$

The following lemma shows that the images of $\omega_{\mathcal{I}}^0$ and $\omega_{\mathcal{I}}$, respectively, characterize the s -tuples \mathcal{E} of flags for which the intersections $\Omega_{\mathcal{I}}^0(\mathcal{E})$ and $\Omega_{\mathcal{I}}(\mathcal{E})$ are nonempty:

Lemma 4.2.3. *Let $\mathcal{I} \in \mathrm{Subsets}(r, n, s)$. Then,*

$$\begin{aligned} \mathrm{im} \omega_{\mathcal{I}}^0 &= \{\mathcal{E} \in \mathrm{Flag}(n)^s : \Omega_{\mathcal{I}}^0(\mathcal{E}) \neq \emptyset\}, \\ \mathrm{im} \omega_{\mathcal{I}} &= \{\mathcal{E} \in \mathrm{Flag}(n)^s : \Omega_{\mathcal{I}}(\mathcal{E}) \neq \emptyset\}. \end{aligned}$$

In particular, $\mathcal{I} \in \mathrm{Intersecting}(r, n, s)$ if and only if $\omega_{\mathcal{I}}$ is surjective.

Proof. If $\mathcal{E} \in \mathrm{im} \omega_{\mathcal{I}}^0$ then there exists $\gamma \in \mathrm{GL}(n)$ such that $E_k \in \gamma \mathrm{Flag}_{I_k}^0(r, n)$ for $k \in [s]$. But

$$\begin{aligned} E_k \in \gamma \mathrm{Flag}_{I_k}^0(r, n) &\Leftrightarrow \gamma^{-1} E_k \in \mathrm{Flag}_{I_k}^0(r, n) \Leftrightarrow V_0 \in \Omega_{I_k}^0(\gamma^{-1} E_k) \\ &\Leftrightarrow \gamma V_0 \in \Omega_{I_k}^0(E_k), \end{aligned}$$

and therefore $\gamma V_0 \in \Omega_{\mathcal{I}}^0(\mathcal{E})$. Conversely, if $V \in \Omega_{\mathcal{I}}^0(\mathcal{E})$, then we write $V = \gamma V_0$ and obtain that $E_k \in \gamma \mathrm{Flag}_{I_k}^0(r, n)$ for all k , and hence that $\mathcal{E} \in \mathrm{im} \omega_{\mathcal{I}}^0$. The result for $\mathrm{im} \omega_{\mathcal{I}}$ is proved in the same way. \square

We now use some basic algebraic geometry (see, e.g., [Per]). Recall that a morphism $f: \mathcal{X} \rightarrow \mathcal{Y}$ of irreducible algebraic varieties is called *dominant* if its image is Zariski dense. In this case, the image contains a nonempty Zariski-open subset \mathcal{Y}_0 such that the dimension of any irreducible component of the fibers $f^{-1}(y)$ for $y \in \mathcal{Y}_0$ is equal to $\dim \mathcal{X} - \dim \mathcal{Y}$. Furthermore, if $\mathcal{X}_0 \subseteq \mathcal{X}$ is a nonempty Zariski-open subset then f is dominant if and only if its restriction f to \mathcal{X}_0 is dominant.

We also recall for future reference the following results: If \mathcal{X} and \mathcal{Y} are smooth (irreducible algebraic) varieties and $f: \mathcal{X} \rightarrow \mathcal{Y}$ is dominant then the set of regular values (i.e., the points y such that df_x is surjective for all preimages $x \in f^{-1}(y)$) contains a Zariski-open set. Also, if df_x is surjective for every x then the image by f of any Zariski-open set in \mathcal{X} is a Zariski-open set in \mathcal{Y} . In particular this is the case when $f: \mathcal{V} \rightarrow \mathcal{B}$ is a vector bundle.

In the present context, the maps $\omega_{\mathcal{I}}^0$ and $\omega_{\mathcal{I}}$ are morphisms of irreducible algebraic varieties and so the preceding discussion applies. Furthermore, the domain of $\omega_{\mathcal{I}}$ is the closure of the domain of $\omega_{\mathcal{I}}^0$ in $\mathrm{GL}(n) \times \mathrm{Flag}(n)^s$. Therefore, $\omega_{\mathcal{I}}$ is dominant if and only if $\omega_{\mathcal{I}}^0$ is dominant.

Lemma 4.2.4. *Let $\mathcal{I} \in \text{Subsets}(r, n, s)$. Then $\mathcal{I} \in \text{Intersecting}(r, n, s)$ if and only if $\omega_{\mathcal{I}}$ or $\omega_{\mathcal{I}}^0$ is dominant.*

Proof. On the one hand, Lemma 4.2.3 shows that \mathcal{I} is intersecting if and only if $\omega_{\mathcal{I}}$ is surjective. On the other hand, we have just observed that $\omega_{\mathcal{I}}^0$ is dominant if and only if $\omega_{\mathcal{I}}$ is dominant. Thus it remains to show that $\omega_{\mathcal{I}}$ is automatically surjective if it is dominant. For this, we observe that the space $\text{Flag}_{I_1}(r, n) \times \cdots \times \text{Flag}_{I_s}(r, n)$ is left invariant by the diagonal action of the parabolic $P(r, n)$, as can be seen from (3.3.2). Thus $\omega_{\mathcal{I}}$ factors over a map

$$(4.2.5) \quad \bar{\omega}_{\mathcal{I}}: \begin{cases} \text{GL}(n) \times_{P(r,n)} \text{Flag}_{I_1}(r, n) \times \cdots \times \text{Flag}_{I_s}(r, n) \rightarrow \text{Flag}(n)^s \\ [\gamma, E_1, \dots, E_s] \mapsto (\gamma E_1, \dots, \gamma E_s). \end{cases}$$

Clearly, $\omega_{\mathcal{I}}$ and $\bar{\omega}_{\mathcal{I}}$ have the same image. If $\bar{\omega}_{\mathcal{I}}$ is dominant, then its image contains a nonempty Zariski-open set and therefore is dense in the Euclidean topology. But the domain of $\bar{\omega}_{\mathcal{I}}$ is compact in the Euclidean topology and hence the image is also closed in the Euclidean topology. It follows that $\bar{\omega}_{\mathcal{I}}$ is automatically surjective if $\bar{\omega}_{\mathcal{I}}$ is dominant. □

A first, obvious condition for \mathcal{I} to be intersecting is therefore that the dimension of the domain of $\omega_{\mathcal{I}}$ is no smaller than the dimension of the target space. If we apply this argument to the factored map (4.2.5), which has the same image, we obtain that the expected dimension introduced in Definition 2.14 is nonnegative:

Lemma 4.2.6. *If $\mathcal{I} \in \text{Intersecting}(r, n, s)$ then*

$$(4.2.7) \quad \text{edim } \mathcal{I} = r(n - r) - \sum_{k=1}^s (r(n - r) - \dim I_k) \geq 0.$$

Proof. Let $\mathcal{X} := \text{GL}(n) \times_{P(r,n)} \text{Flag}_{I_1}(r, n) \times \cdots \times \text{Flag}_{I_s}(r, n)$ and $\mathcal{Y} := \text{Flag}(n)^s$. If \mathcal{I} is intersecting then the map $\bar{\omega}_{\mathcal{I}}: \mathcal{X} \rightarrow \mathcal{Y}$ in (4.2.5) is dominant, hence $\dim \mathcal{X} \geq \dim \mathcal{Y}$. But

$$(4.2.8) \quad \begin{aligned} \dim \mathcal{X} - \dim \mathcal{Y} &= (\dim \text{GL}(n)/P(r, n)) + \sum_{k=1}^s (\dim \text{Flag}_{I_k}(r, n) - \dim \text{Flag}(n)) \\ &= \dim \text{Gr}(r, n) - \sum_{k=1}^s (\dim \text{Gr}(r, n) - \dim I_k) = \text{edim } \mathcal{I} \end{aligned}$$

where the first equality is obvious and the second is Eq. (3.3.8). □

At this point, we have established all facts that we used in Section 2 to prove Corollary 2.17. That is, the proof of Corollary 2.17 is now complete.

We conclude this section by recording the following rules for the expected dimension,

$$(4.2.9) \quad \text{edim } \mathcal{I}/\mathcal{J} = \text{edim } \mathcal{I} + \text{edim } \mathcal{J} - \text{edim } \mathcal{I}\mathcal{J},$$

$$(4.2.10) \quad \text{edim } \mathcal{I}^{\mathcal{J}}\mathcal{K} - \text{edim } \mathcal{K} = \text{edim } \mathcal{I}(\mathcal{J}\mathcal{K}) - \text{edim } \mathcal{J}\mathcal{K},$$

$$(4.2.11) \quad \text{edim } \mathcal{I}^{\mathcal{J}} = \text{edim } \mathcal{I}\mathcal{J} - \text{edim } \mathcal{J},$$

which hold for all $\mathcal{I} \in \text{Subsets}(r, n, s)$, $\mathcal{J} \in \text{Subsets}(d, r, s)$, and $\mathcal{K} \in \text{Subsets}(m, d, s)$. Equations (4.2.9) to (4.2.11) are direct consequences of Lemmas 3.2.13 and 3.2.14. Eq. (4.2.10) in particular will play a crucial role in Section 5.3, as we will use it to show that if \mathcal{I} satisfies the Horn inequalities and \mathcal{J} is intersecting then so does $\mathcal{I}^{\mathcal{J}}$. This will be key to establishing Belkale’s theorem on the sufficiency of the Horn inequalities by induction (Theorem 5.3.4).

4.3. Slopes and Horn inequalities. We are now interested in proving a strengthened version of Corollary 2.17 (see Corollary 4.3.11 below). As a first step, we introduce the promised ‘good set’ of s -tuples of flags which are sufficiently generic to detect when an s -tuple \mathcal{I} is intersecting: Define in analogy to (4.2.5) the map

$$\bar{\omega}_{\mathcal{I}}^0: \begin{cases} \text{GL}(n) \times_{P(r,n)} \text{Flag}_{I_1}^0(r, n) \times \cdots \times \text{Flag}_{I_s}^0(r, n) \rightarrow \text{Flag}(n)^s \\ [\gamma, E_1, \dots, E_s] \mapsto (\gamma E_1, \dots, \gamma E_s) \end{cases}$$

Lemma 4.3.1. *There exists a nonempty Zariski-open subset $\text{Good}(n, s) \subseteq \text{Flag}(n)^s$ that satisfies the following three properties for all $r \in [n]$:*

- (a) $\text{Good}(n, s)$ consists of regular values (in the image) of $\bar{\omega}_{\mathcal{I}}^0$ for every $\mathcal{I} \in \text{Intersecting}(r, n, s)$.
- (b) For every $\mathcal{I} \in \text{Subsets}(r, n, s)$, the following are equivalent:
 - (i) $\mathcal{I} \in \text{Intersecting}(r, n, s)$.
 - (ii) For all $\mathcal{E} \in \text{Good}(n, s)$, $\Omega_{\mathcal{I}}^0(\mathcal{E}) \neq \emptyset$.
 - (iii) There exists $\mathcal{E} \in \text{Good}(n, s)$ such that $\Omega_{\mathcal{I}}^0(\mathcal{E}) \neq \emptyset$.
- (c) If $\mathcal{I} \in \text{Intersecting}(r, n, s)$, then for every $\mathcal{E} \in \text{Good}(n, s)$ the variety $\Omega_{\mathcal{I}}^0(\mathcal{E})$ has the same number of irreducible components, each connected component is of dimension $\text{edim } \mathcal{I}$, and $\Omega_{\mathcal{I}}^0(\mathcal{E})$ is dense in $\Omega_{\mathcal{I}}(\mathcal{E})$.

Proof. Let us construct $\text{Good}(n, s)$ satisfying the properties above. Let $\mathcal{I} \in \text{Subsets}(r, n, s)$, where $r \in [n]$. If $\mathcal{I} \notin \text{Intersecting}(r, n, s)$ then by Lemma 4.2.4 the map $\omega_{\mathcal{I}}^0$ is not dominant, and we define $U_{\mathcal{I}}$ as the complement of the Zariski-closure of $\text{im } \omega_{\mathcal{I}}^0$. Thus $U_{\mathcal{I}}$ is a nonempty Zariski-open subset of $\text{Flag}(n)^s$. Otherwise, if $\mathcal{I} \in \text{Intersecting}(r, n, s)$ then $\omega_{\mathcal{I}}^0$ is dominant by Lemma 4.2.4. The map $\bar{\omega}_{\mathcal{I}}^0$ has the same image as $\omega_{\mathcal{I}}^0$ and is therefore also a dominant map between smooth irreducible varieties. Thus its image contains a nonempty Zariski-open subset $U_{\mathcal{I}}$ of $\text{Flag}(n)^s$ consisting of regular values, such that the fibers $(\bar{\omega}_{\mathcal{I}}^0)^{-1}(\mathcal{E})$ for $\mathcal{E} \in U_{\mathcal{I}}$ all have the same number of irreducible components, each of dimension equal to $\text{edim } \mathcal{I}$, by the calculation in (4.2.8). We now define the good set as

$$\text{Good}(n, s) := \bigcap_{\mathcal{I}} U_{\mathcal{I}},$$

where the intersection is over all s -tuples \mathcal{I} , intersecting or not. As a finite intersection of nonempty Zariski-open subsets, $\text{Good}(n, s)$ is again nonempty and Zariski-open. By construction, it satisfies property (a).

We now show that $\text{Good}(n, s)$ satisfies (b). To see that (bi) implies (bii), note that for any $\mathcal{I} \in \text{Intersecting}(r, n, s)$ and $\mathcal{E} \in \text{Good}(n, s)$, $\mathcal{E} \in U_{\mathcal{I}} \subseteq \text{im } \bar{\omega}_{\mathcal{I}}^0 = \text{im } \omega_{\mathcal{I}}^0$. Thus Lemma 4.2.3 shows that $\Omega_{\mathcal{I}}^0(\mathcal{E}) \neq \emptyset$. Clearly, (bii) implies (biii) since $\text{Good}(n, s)$ is nonempty. Lastly, suppose that (biii) holds. By Lemma 4.2.3, $\Omega_{\mathcal{I}}^0(\mathcal{E}) \neq \emptyset$ implies that $\mathcal{E} \in \text{im } \omega_{\mathcal{I}}^0$. But $\mathcal{E} \in \text{Good}(n, s) \subseteq (\text{im } \omega_{\mathcal{I}})^c \subseteq (\text{im } \omega_{\mathcal{I}}^0)^c$ unless \mathcal{I} is intersecting; this establishes (bi).

Lastly, we verify c. Observe that, for any $\mathcal{E} \in \text{Flag}(n)^s$, the fiber $(\bar{\omega}_{\mathcal{I}}^0)^{-1}(\mathcal{E})$ is equal to the set of $[\gamma, \gamma^{-1}E_1, \dots, \gamma^{-1}E_s]$ such that $\gamma^{-1}E_k \in \text{Flag}_{I_k}^0(r, n)$ for all $k \in [s]$. It can therefore by $\gamma \mapsto \gamma V_0$ be identified with $\Omega_{\mathcal{I}}^0(\mathcal{E})$. Now assume that $\mathcal{I} \in \text{Intersecting}(r, n, s)$. As we vary $\mathcal{E} \in \text{Good}(n, s)$, $\mathcal{E} \in U_{\mathcal{I}}$ and so $(\bar{\omega}_{\mathcal{I}}^0)^{-1}(\mathcal{E}) \cong \Omega_{\mathcal{I}}^0(\mathcal{E})$ has the same number of irreducible components, each of dimension $\text{edim } \mathcal{I}$. We still need to show that $\Omega_{\mathcal{I}}^0(\mathcal{E})$ is dense in $\Omega_{\mathcal{I}}(\mathcal{E})$. This will follow if we can show that $\Omega_{\mathcal{I}}^0(\mathcal{E})$ meets any irreducible component \mathcal{Z} of $\Omega_{\mathcal{I}}(\mathcal{E})$. Let us assume that this is not the case, so that $\mathcal{Z} \subseteq \Omega_{\mathcal{I}}(\mathcal{E}) \setminus \Omega_{\mathcal{I}}^0(\mathcal{E})$. But

$$\Omega_{\mathcal{I}}(\mathcal{E}) \setminus \Omega_{\mathcal{I}}^0(\mathcal{E}) = \bigcup_{k=1}^s \left((\Omega_{I_k}(E_k) \setminus \Omega_{I_k}^0(E_k)) \cap \bigcap_{l \neq k} \Omega_{I_l}(E_l) \right) = \bigcup_{\substack{I'_1 \leq I_1, \dots, I'_s \leq I_s \\ \exists k \in [s]: I'_k \neq I_k}} \Omega_{\mathcal{I}'}^0$$

by Lemma 3.1.5. That is, $\Omega_{\mathcal{I}}(\mathcal{E}) \setminus \Omega_{\mathcal{I}}^0(\mathcal{E})$ is a union of varieties $\Omega_{\mathcal{I}'}^0(\mathcal{E})$ with $\text{edim } \mathcal{I}' < \text{edim } \mathcal{I}$. If \mathcal{I}' is intersecting then any irreducible component of $\Omega_{\mathcal{I}'}^0(\mathcal{E})$ has dimension equal to $\text{edim } \mathcal{I}'$. Otherwise, if \mathcal{I}' is not intersecting, then $\Omega_{\mathcal{I}'}^0(\mathcal{E}) = \emptyset$. It follows that any irreducible component of $\Omega_{\mathcal{I}}(\mathcal{E}) \setminus \Omega_{\mathcal{I}}^0(\mathcal{E})$ has dimension strictly smaller than $\text{edim } \mathcal{I}$. But this is a contradiction, since the dimension of \mathcal{Z} is equal to at least $\text{edim } \mathcal{I}$. \square

The following is a direct consequence of the equivalence between (bi) and (biii) in Lemma 4.3.1:

$$(4.3.2) \quad \text{Intersecting}(r, n, s) = \{ \text{Pos}(V, \mathcal{E}) : V \subseteq \mathbb{C}^n, \dim V = r \}$$

for every $\mathcal{E} \in \text{Good}(n, s)$.

We now study the numerical inequalities satisfied by intersecting s -tuples more carefully. Recall that a weight θ for $\text{GL}(r)$ is *antidominant* if $\theta(1) \leq \dots \leq \theta(r)$. For example, given a subset $I \subseteq [n]$ of cardinality r , the weight $\theta(a) := I(a) - a$ is antidominant. It is convenient to introduce the following definition:

Definition 4.3.3. Given an s -tuple $\vec{\theta} = (\theta_1, \dots, \theta_s)$ of antidominant weights for $\text{GL}(r)$, we define the *slope* of a tuple $\mathcal{J} \in \text{Subsets}(d, r, s)$ as

$$\mu_{\vec{\theta}}(\mathcal{J}) := \frac{1}{d} \sum_{k=1}^s \sum_{a \in J_k} \theta_k(a) = \frac{1}{d} \sum_{k=1}^s (T_{J_k}, \theta_k).$$

For any nonzero subspace $\{0\} \neq S \subseteq \mathbb{C}^r$ and s -tuple of flags \mathcal{F} on \mathbb{C}^r , we further define

$$\mu_{\vec{\theta}}(S, \mathcal{F}) := \mu_{\vec{\theta}}(\text{Pos}(S, \mathcal{F})).$$

Here and in the following, we write $\text{Pos}(S, \mathcal{F})$ for the s -tuple of positions $(\text{Pos}(S, F_k))_{k \in [s]}$.

Note that we can interpret $\mu_{\vec{\theta}}(\mathcal{J})$ as a sum of averages of the nowhere decreasing functions θ_k for uniform choice of $a \in J_k$.

The following lemma asserts that there is a unique slope-minimizing subspace of maximal dimension:

Lemma 4.3.4 (Harder–Narasimhan, [Bell]). *Let $\vec{\theta}$ be an s -tuple of antidominant weights for $\text{GL}(r)$, and $\mathcal{F} \in \text{Flag}(r)^s$. Let $m_* := \min_{\{0\} \neq S \subseteq \mathbb{C}^r} \mu_{\vec{\theta}}(S, \mathcal{F})$ and $d_* := \max\{\dim S : \mu_{\vec{\theta}}(S, \mathcal{F}) = m_*\}$. Then there exists a unique subspace $S_* \subseteq \mathbb{C}^r$ such that $\mu_{\vec{\theta}}(S_*, \mathcal{F}) = m_*$ and $\dim S_* = d_* > 0$.*

Proof. Existence is immediate, so it remains to show uniqueness. Thus suppose for sake of finding a contradiction that there are two such subspaces, $S_1 \neq S_2$, such that $\mu_{\vec{\theta}}(S_j, \mathcal{F}) = m_*$ and $\dim S_j = d_*$ for $j = 1, 2$. We note that $d_* > 0$ and that the inclusions $S_1 \cap S_2 \subsetneq S_1$ and $S_2 \subsetneq S_1 + S_2$ are strict.

Let $\mathcal{J} = \text{Pos}(S_1, \mathcal{F})$ and $\mathcal{K} = \text{Pos}(S_1 \cap S_2, \mathcal{F}^{S_1})$. Then $\text{Pos}(S_1 \cap S_2, \mathcal{F}) = \mathcal{J}\mathcal{K}$ by the chain rule (Lemma 3.2.8). Let us first assume that $S_1 \cap S_2 \neq \{0\}$, so that $\mu_{\vec{\theta}}(\mathcal{J}\mathcal{K})$ is well-defined. Then,

$$\mu_{\vec{\theta}}(\mathcal{J}\mathcal{K}) = \mu_{\vec{\theta}}(S_1 \cap S_2, \mathcal{F}) \geq m_* = \mu_{\vec{\theta}}(S_1, \mathcal{F}) = \mu_{\vec{\theta}}(\mathcal{J}),$$

where the equalities hold by definition, and the inequality holds as m_* is the minimal slope. On the other hand, note that $J_k = J_k K_k \cup J_k K_k^c$ for each $k \in [s]$, hence we can write

$$\mu_{\bar{\theta}}(\mathcal{J}) = \frac{d}{d_*} \mu_{\bar{\theta}}(\mathcal{JK}) + \frac{d_* - d}{d_*} \mu_{\bar{\theta}}(\mathcal{JK}^c),$$

where $d := \dim S_1 \cap S_2 < \dim S_1 = d_*$. It follows that

$$(4.3.5) \quad m_* = \mu_{\bar{\theta}}(\mathcal{J}) \geq \mu_{\bar{\theta}}(\mathcal{JK}^c).$$

If $S_1 \cap S_2 = \{0\}$ then $\mathcal{J} = \mathcal{JK}^c$ and so (4.3.5) holds with equality.

Likewise, let $\mathcal{L} = \text{Pos}(S_1 + S_2, \mathcal{F})$ and $\mathcal{M} = \text{Pos}(S_2, \mathcal{F}^{S_1+S_2})$. Since $S_2 \subsetneq S_1 + S_2$, but S_2 was assumed to be a maximal-dimensional subspace with minimal slope, it follows that the slope of $S_1 + S_2$ is strictly larger than m_* :

$$\mu_{\bar{\theta}}(\mathcal{LM}) = \mu_{\bar{\theta}}(S_2, \mathcal{F}) = m_* < \mu_{\bar{\theta}}(S_1 + S_2, \mathcal{F}) = \mu_{\bar{\theta}}(\mathcal{L}).$$

Just as before, we decompose

$$\mu_{\bar{\theta}}(\mathcal{L}) = \frac{d_*}{d'} \mu_{\bar{\theta}}(\mathcal{LM}) + \frac{d' - d_*}{d'} \mu_{\bar{\theta}}(\mathcal{LM}^c),$$

where now $d' := \dim S_1 + S_2 > \dim S_2 = d_* > 0$. Thus we obtain the strict inequality

$$(4.3.6) \quad \mu_{\bar{\theta}}(\mathcal{LM}^c) > \mu_{\bar{\theta}}(\mathcal{LM}) = m_*.$$

At last, we apply Corollary 3.2.12, which shows that $J_k K_k^c(b) \geq L_k M_k^c(b)$ for all b and k , and hence

$$\mu_{\bar{\theta}}(\mathcal{JK}^c) \geq \mu_{\bar{\theta}}(\mathcal{LM}^c).$$

Together with (4.3.5) and (4.3.6), we obtain the desired contradiction:

$$m_* \geq \mu_{\bar{\theta}}(\mathcal{JK}^c) \geq \mu_{\bar{\theta}}(\mathcal{LM}^c) > m_*. \quad \square$$

We will now use Lemmas 4.3.1 and 4.3.4 to show that the conditions in Corollary 2.17 with $\text{edim } \mathcal{J} = 0$ imply those for general intersecting \mathcal{J} .

Definition 4.3.7. Let $I \subseteq [n]$ be a subset of cardinality r . We define $\lambda_I \in \Lambda_+(r)$ by

$$\lambda_I(a) := a - I(a) \quad (a \in [r]).$$

Any highest weight λ with $\lambda(1) \leq 0$, $\lambda(r) \geq r - n$ can be written in this form. Moreover, if I^c denotes the complement of I in $[n]$ then the dominant weight $\lambda_{I^c} \in \Lambda_+(n - r)$ can be written as

$$(4.3.8) \quad \begin{aligned} \lambda_{I^c}(b) &= b - I^c(b) = -\#\{a \in [r] : I(a) < I^c(b)\} \\ &= -\#\{a \in [r] : I(a) - a < b\}. \end{aligned}$$

Remark. This equation has a pleasant interpretation in terms of Young diagrams. Consider the Young diagram Y_I corresponding to λ_I^* , which has $I(r + 1 - a) - (r + 1 - a)$ boxes in its a -th row. By definition, its transpose Y_I^t is the Young diagram such that the number of boxes in the b -th row is equal to the number of boxes in the b -th column of T_I . Thus (4.3.8) asserts that $Y_I^t = r\mathbb{1}_{n-r} + \lambda_{I^c}$, i.e., the two Young diagrams Y_I^t and Y_{I^c} (the latter with rows in reverse order) make up a rectangle of size $r \times (n - r)$.

Lemma 4.3.9. *Let $\mathcal{I} \in \text{Subsets}(r, n, s)$. Set $\lambda_k = \lambda_{I_k} + (n - r)\mathbb{1}_r$ for $k \in [s - 1]$ and $\lambda_s = \lambda_{I_s}$. Then we have that $\text{edim } \mathcal{I} = -\sum_{k=1}^s |\lambda_k|$. More generally, for every $\mathcal{J} \in \text{Subsets}(d, r, s)$,*

$$\text{edim } \mathcal{I}\mathcal{J} - \text{edim } \mathcal{J} = -\sum_{k=1}^s (T_{J_k}, \lambda_k) = d\mu_{-\vec{\lambda}}(\mathcal{J}),$$

where we recall that $(T_J, \xi) = \sum_{j \in J} \xi(j)$ for any $J \subseteq [r]$ and $\xi \in \text{it}(r)$.

Proof. It suffices to prove the second statement, which follows from

$$\begin{aligned} \text{edim } \mathcal{I}\mathcal{J} - \text{edim } \mathcal{J} &= d(n - r)(1 - s) + \sum_{k=1}^s \sum_{a \in J_k} (I_k(a) - a) \\ &= d(n - r)(1 - s) - \sum_{k=1}^s (T_{J_k}, \lambda_{I_k}) = -\sum_{k=1}^s (T_{J_k}, \lambda_k) \\ &= d\mu_{-\vec{\lambda}}(\mathcal{J}). \end{aligned}$$

□

It follows that minimizing $\mu_{-\vec{\lambda}}(\mathcal{J})$ and $\frac{1}{d}(\text{edim } \mathcal{I}\mathcal{J} - \text{edim } \mathcal{J})$ as a function of \mathcal{J} are equivalent. We then have the following result:

Proposition 4.3.10. *Let $\mathcal{I} \in \text{Subsets}(r, n, s)$ such that $\text{edim } \mathcal{I} \geq 0$ and, for any $0 < d < r$ and $\mathcal{J} \in \text{Intersecting}(d, r, s)$ with $\text{edim } \mathcal{J} = 0$ we have that $\text{edim } \mathcal{I}\mathcal{J} \geq 0$. Then we have for any $0 < d < r$ and $\mathcal{J} \in \text{Intersecting}(d, r, s)$ that*

$$\text{edim } \mathcal{I}\mathcal{J} \geq \text{edim } \mathcal{J}.$$

Proof. Suppose for sake of finding a contradiction that there exists $\mathcal{J} \in \text{Intersecting}(d, r, s)$ with $0 < d < r$ and $\text{edim } \mathcal{I}\mathcal{J} < \text{edim } \mathcal{J}$, so that $\mu_{-\vec{\lambda}}(\mathcal{J}) < 0$ according to Lemma 4.3.9. Fix some $\mathcal{F} \in \text{Good}(r, s)$. Then $\Omega_{\mathcal{J}}^0(\mathcal{F}) \neq \emptyset$ by Lemma 4.3.1, (bii). Thus there exists a subspace $\{0\} \neq S \subseteq \mathbb{C}^r$ such that $\mu_{-\vec{\lambda}}(S, \mathcal{F}) = \mu_{-\vec{\lambda}}(\mathcal{J}) < 0$.

Now let S_* be the unique subspace of minimal slope $m_* < 0$ and maximal dimension $d_* > 0$ from Lemma 4.3.4 and denote by $\mathcal{J}_* := \text{Pos}(S_*, \mathcal{F})$ its s -tuple of positions. The uniqueness statement implies that $\Omega_{\mathcal{J}_*}^0(\mathcal{F}) = \{S_*\}$, since slope and dimension are fully determined by the position. Moreover, \mathcal{J}_* is intersecting by (4.3.2), and therefore $\text{edim } \mathcal{J}_* = \dim \Omega_{\mathcal{J}_*}^0(\mathcal{F}) = 0$ by Lemma 4.3.1. Thus we have found an s -tuple $\mathcal{J}_* \in \text{Intersecting}(d_*, r, s)$ with $d_* > 0$, $\text{edim } \mathcal{J}_* = 0$, and

$$\text{edim } \mathcal{I}\mathcal{J}_* = \text{edim } \mathcal{I}\mathcal{J}_* - \text{edim } \mathcal{J}_* = d_*m_* < 0,$$

where we have used Lemma 4.3.9 once again in the last equality. Since $\text{edim } \mathcal{I} \geq 0$, this also implies that $d_* < r$. This is the desired contradiction. \square

Proposition 4.3.10 will be useful to prove Belkale’s Theorem 5.3.4 in Section 5 below, since it allows us to work with a larger set of inequalities.

Remark. The proof of Proposition 4.3.10 shows that we may in fact restrict to \mathcal{J} such that $\Omega_{\mathcal{J}}^0(\mathcal{F})$ is a point for all s -tuples of good flags $\mathcal{F} \in \text{Good}(r, s)$ – or also to those for which $\Omega_{\mathcal{J}}(\mathcal{F})$ is a point, which is equivalent by the last statement in Lemma 4.3.1. See the remark after Corollary 6.3.3 for the implications of this on the description of the Kirwan cone.

We also record the following corollary which follows together with and improves over Corollary 2.17.

Corollary 4.3.11. *If $\mathcal{I} \in \text{Intersecting}(r, n, s)$ then for any $0 < d < r$ and any s -tuple $\mathcal{J} \in \text{Intersecting}(d, r, s)$ we have that $\text{edim } \mathcal{I}\mathcal{J} \geq \text{edim } \mathcal{J}$.*

We remark that for $d = r$ there is only one s -tuple, $\mathcal{J} = ([r], \dots, [r])$, and it is intersecting and satisfies $\text{edim } \mathcal{J} = 0$. In this case, $\text{edim } \mathcal{I}\mathcal{J} - \text{edim } \mathcal{J} = \text{edim } \mathcal{I}$, and so we may safely allow for $d = r$ in Corollaries 2.17 and 4.3.11 and Proposition 4.3.10.

We conclude this section with some simple examples of the Horn inequalities of Corollary 4.3.11. We refer to Appendix A for lists of all Horn triples $\mathcal{I} = (I_1, I_2, I_3)$ up to $n = 4$.

Example 4.3.12 ($r = 1$). The only condition for $\mathcal{I} \in \text{Intersecting}(1, n, s)$ is the dimension condition, $\text{edim } \mathcal{I} \geq 0$. Indeed, the Grassmannian $\text{Gr}(1, n)$ is the projective space $\mathbb{P}(\mathbb{C}^n)$, whose Schubert varieties are given by $\Omega_{\{i\}}(E) = \{[v] \in \mathbb{P}(\mathbb{C}^n) : v \in E(i)\}$. Thus $\mathcal{I} = (\{i_1\}, \dots, \{i_s\})$ is intersecting if and only if for any s -tuple of flags \mathcal{E} , $E_1(i_1) \cap \dots \cap E_s(i_s) \neq \{0\}$. By linear algebra, it is certainly sufficient that $\sum_{k=1}^s (n - i_k) \leq n - 1$, which is equivalent to $\text{edim } \mathcal{I} \geq 0$. This also establishes Theorem 5.3.4 in the case $r = 1$.

Example ($s = 2, r = 2$). Let $\mathcal{I} = (I_1, I_2)$. Then the condition $\text{edim } \mathcal{I} \geq 0$ is $I_1(1) + I_1(2) + I_2(1) + I_2(2) \geq 2n + 2$. However, there are two additional conditions coming from the $\mathcal{J} \in \text{Intersecting}(1, 2, 2)$ with $\text{edim } \mathcal{J} = 0$. By the preceding example, there are two such pairs, $(\{1\}, \{2\})$ and $(\{2\}, \{1\})$. The corresponding conditions are $I_1(1) + I_2(2) \geq n + 1$ and $I_1(2) + I_2(1) \geq n + 1$.

For example, if $n = 4$ then $\mathcal{I} = (\{1, 4\}, \{2, 4\})$ satisfies all Horn inequalities. On the other hand, $\mathcal{I} = (\{1, 4\}, \{2, 3\})$ fails one the Horn inequalities. Indeed, if we consider $\mathcal{J} = (\{1\}, \{2\})$ then $\mathcal{I}\mathcal{J} = (\{1\}, \{3\})$ is such that $\text{edim } \mathcal{I}\mathcal{J} = -1 < 0$.

5. Sufficiency of Horn inequalities

In this section we prove that the Horn inequalities are also sufficient to characterize intersections of Schubert varieties.

5.1. Tangent maps. In Lemma 4.2.4, we established that an s -tuple \mathcal{I} is intersecting if and only if the corresponding morphism $\omega_{\mathcal{I}}$ defined in (4.2.2) is dominant. Now it is a general fact that a morphism $f: \mathcal{X} \rightarrow \mathcal{Y}$ between smooth and irreducible varieties is dominant if and only if there exists a point $p \in \mathcal{X}$ where the differential $T_p f$ is surjective. This will presently allow us to reduce the intersecting of Schubert varieties to an infinitesimal question about tangent maps. Later, in Section 6, we will also use the determinant of the tangent map to construct explicit nonzero tensor product invariants and establish the saturation property.

Lemma 5.1.1. *Let $\mathcal{I} \in \text{Subsets}(r, n, s)$. Then $\mathcal{I} \in \text{Intersecting}(r, n, s)$ if and only if there exist $\vec{g} = (g_1, \dots, g_s) \in \text{GL}(V_0)^s$ and $\vec{h} = (h_1, \dots, h_s) \in \text{GL}(Q_0)^s$ such that the linear map*

(5.1.2)

$$\Delta_{\mathcal{I}, \vec{g}, \vec{h}}: \begin{cases} \text{Hom}(V_0, Q_0) \times H_{I_1}(F_0, G_0) \times \dots \times H_{I_s}(F_0, G_0) \rightarrow \text{Hom}(V_0, Q_0)^s \\ (\zeta, \phi_1, \dots, \phi_s) \mapsto (\zeta + h_1 \phi_1 g_1^{-1}, \dots, \zeta + h_s \phi_s g_s^{-1}) \end{cases}$$

is surjective.

Proof. Using the isomorphisms $\text{Flag}_{I_k}^0(r, n) = G_{I_k}(r, n)E_0 \cong G_{I_k}(r, n)/B(n)$ (Definition 3.3.3 and Section 4.1) and $\text{Flag}(n) \cong \text{GL}(n)/B(n)$, we find that $\omega_{\mathcal{I}}^0$ is dominant if and only if

(5.1.3)

$$\text{GL}(n) \times G_{I_1}(r, n) \times \dots \times G_{I_s}(r, n) \rightarrow \text{GL}(n)^s, (\gamma, \gamma_1, \dots, \gamma_s) \mapsto (\gamma\gamma_1, \dots, \gamma\gamma_s)$$

is dominant. This is again a morphism between smooth and irreducible varieties and thus dominance is equivalent to surjectivity of the differential at some point

$(\gamma, \gamma_1, \dots, \gamma_s)$. The map (5.1.3) is $GL(n)$ -equivariant on the left and $B(n)^s$ -equivariant on the right. By the former, we may assume that $\gamma = 1$, and by the latter that $\gamma_k = p_k w_{I_k}$ for some $p_k = \begin{pmatrix} g_k & b_k \\ 0 & h_k \end{pmatrix}$, since $G_{I_k}(r, n) = P(r, n)w_{I_k}B(n)$ according Lemma 3.3.4.

We now compute the differential. Thus we consider an arbitrary curve $1 + \varepsilon X$ tangent to $\gamma = 1$, where $X \in \mathfrak{gl}(n)$, and curves $(1 + \varepsilon Y_k)p_k w_{I_k}$ through the $\gamma_k = p_k w_k$, where $Y_k \in \mathfrak{gl}(n)$. If we write $Y_k = \begin{pmatrix} A_k & B_k \\ C_k & D_k \end{pmatrix}$ with $A_k \in \mathfrak{gl}(r)$ etc., then we see from (4.1.1) that $(1 + \varepsilon Y_k)p_k w_{I_k}$ is tangent to $G_{I_k}(r, n)$ precisely if $h_k^{-1}C_k g_k \in H_{I_k}(F_0, G_0)$, that is, if $C_k \in h_k H_{I_k}(F_0, G_0)g_k^{-1}$. Lastly, the calculation $(1 + \varepsilon X)(1 + \varepsilon Y_k)\gamma_k = \gamma_k + \varepsilon(X + Y_k)\gamma_k + O(\varepsilon^2)$ shows that the differential of (5.1.3) at $(g, \gamma_1, \dots, \gamma_s)$ can be identified with $(X, Y_1, \dots, Y_s) \mapsto (X + Y_1, \dots, X + Y_s)$.

We may check for surjectivity block by block. Since there are no constraints on the A_k , B_k , and D_k , it is clear that the differential is surjective on the three blocks corresponding to $\mathfrak{p}(r, n)$. Thus we only need to check surjectivity on the last block of the linear map, corresponding to $\text{Hom}(V_0, Q_0)$. This block can plainly be identified with (5.1.2), since the C_k are constrained to be elements of $h_k H_{I_k}(F_0, G_0)g_k^{-1}$. Thus we obtain that $\omega_{\mathcal{I}}$ is dominant if and only if (5.1.2) is surjective. \square

Remark 5.1.4. The map $\Delta_{\mathcal{I}, \vec{g}, \vec{h}}$ can be identified with the differential of $\bar{\delta}_{\mathcal{I}}^0$ at the point $[1, \mathcal{E}]$, where $E_k \in \text{Flag}_{\mathfrak{gl}(r, n)}^0$ is such that $(E_k)^{V_0} = g_k \cdot F_0$ and $(E_k)^{Q_0} = h_k \cdot G_0$ for $k \in [s]$. This follows from the proof of Lemma 5.1.1 and justifies calling $\Delta_{\mathcal{I}, \vec{g}, \vec{h}}$ a *tangent map*.

By the rank-nullity theorem and using Lemma 3.1.7, the kernel of the linear map $\Delta_{\mathcal{I}, \vec{g}, \vec{h}}$ defined in (5.1.2) is of dimension at least

$$(5.1.5) \quad \begin{aligned} \dim(\text{Hom}(V_0, Q_0) \times H_{I_1}(F_0, G_0) \times \dots \times H_{I_s}(F_0, G_0)) - \dim \text{Hom}(V_0, Q_0)^s \\ = r(n - r)(1 - s) + \sum_{k=1}^s \dim I_k = \text{edim } \mathcal{I}, \end{aligned}$$

and $\Delta_{\mathcal{I}, \vec{g}, \vec{h}}$ is surjective if and only if equality holds. On the other hand, it is immediate that

$$(5.1.6) \quad \ker \Delta_{\mathcal{I}, \vec{g}, \vec{h}} = \bigcap_{k=1}^s h_k H_{I_k}(F_0, G_0)g_k = \bigcap_{k=1}^s H_{I_k}(F_k, G_k),$$

where $F_k = g_k F_0$ and $G_k = h_k G_0$. As we vary g_k and h_k , the F_k and G_k are arbitrary flags on V_0 and Q_0 , respectively. Thus we obtain the following characterization:

Definition 5.1.7. Let $\mathcal{I} \in \text{Subsets}(r, n, s)$. We define the *true dimension* of \mathcal{I} as

$$(5.1.8) \quad \text{tdim } \mathcal{I} := \min_{\mathcal{F}, \mathcal{G}} \dim H_{\mathcal{I}}(\mathcal{F}, \mathcal{G}) = \min_{\vec{g}, \vec{h}} \dim \ker \Delta_{\mathcal{I}, \vec{g}, \vec{h}},$$

where the first side minimization is over all s -tuples of flags \mathcal{F} on V_0 and \mathcal{G} on Q_0 , the second one over $\vec{g} \in \text{GL}(r)^s$, $\vec{h} \in \text{GL}(n - r)^s$, and where

$$H_{\mathcal{I}}(\mathcal{F}, \mathcal{G}) := \bigcap_{k=1}^s H_{I_k}(F_k, G_k) \subseteq \text{Hom}(V_0, Q_0).$$

Corollary 5.1.9. Let $\mathcal{I} \in \text{Subsets}(r, n, s)$. Then we have $\text{tdim } \mathcal{I} \geq \text{edim } \mathcal{I}$, with equality if and only if $\mathcal{I} \in \text{Intersecting}(r, n, s)$.

We note that for the purpose of computing true dimensions we may always assume that F_1 and G_1 are the standard flags on V_0 and Q_0 , respectively (by equivariance).

Example ($s=2, r=2, n=4$). We verify the example at the end of Section 4 by using Corollary 5.1.9. We first consider $\mathcal{I} = (\{1, 4\}, \{2, 4\})$. Then $\text{edim } \mathcal{I} = 1$. To bound $\text{tdim } \mathcal{I}$, we let $\mathcal{F} = (F_1, F_2)$ and $\mathcal{G} = (G_1, G_2)$, where F_1 is the standard flag on V_0 , F_2 the flag with adapted basis $(e(1) + e(2), e(2))$, and $G_1 = G_2$ the standard flags on Q_0 . Then

$$H_{\mathcal{I}}(\mathcal{F}, \mathcal{G}) = \left\{ \begin{pmatrix} 0 & * \\ 0 & * \end{pmatrix} \right\} \cap H_{I_2}(F_2, G_2) = \left\{ \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix} \right\}$$

is one-dimensional, which shows that $\text{tdim } \mathcal{I} \leq 1$. Since always $\text{tdim } \mathcal{I} \geq \text{edim } \mathcal{I}$, it follows that, in fact, $\text{tdim } \mathcal{I} = \text{edim } \mathcal{I}$ and so \mathcal{I} is intersecting.

We now consider $\mathcal{I} = (\{1, 4\}, \{2, 3\})$. Then $\text{edim } \mathcal{I} = 0$. Let \mathcal{F} and \mathcal{G} be pairs of flags on V_0 and Q_0 , respectively. Without loss of generality, we shall assume that F_1 and G_1 are the standard flags. Then

$$H_{\mathcal{I}}(\mathcal{F}, \mathcal{G}) = \left\{ \begin{pmatrix} 0 & * \\ 0 & * \end{pmatrix} \right\} \cap H_{I_2}(F_2, G_2) = \mathbb{C} \begin{pmatrix} 0 & x \\ 0 & y \end{pmatrix},$$

where $\mathbb{C} \begin{pmatrix} x \\ y \end{pmatrix} := G_2(1)$. Indeed, $H_{I_2}(F_2, G_2)$ consists of those linear maps that map any vector in V_0 into $G_2(1)$. In particular, $H_{\mathcal{I}}(\mathcal{F}, \mathcal{G})$ is one-dimensional for any choice of F_2 and G_2 . Thus $\text{tdim } \mathcal{I} = 1 > 0 = \text{edim } \mathcal{I}$, and we conclude that \mathcal{I} is not intersecting.

Example ($s=2,r=3,n=6$). Let $\mathcal{I} = (\{3, 4, 6\}, \{2, 4, 5\})$. Then $\text{edim } \mathcal{I} = 3$. We now establish that \mathcal{I} is intersecting by verifying that $\text{tdim } \mathcal{I} = 3$. Again we choose F_1 and G_1 to be the standard flags on V_0 and Q_0 , respectively, while F_2 and G_2 are defined as follows in terms of adapted bases:

$$\begin{aligned} F_2: & \quad e(1) + z_{21}e(2) + z_{31}e(3), \quad e(2) + z_{32}e(3), \quad e(3), \\ G_2: & \quad \bar{e}(1) + u_{21}\bar{e}(2) + u_{31}\bar{e}(3), \quad \bar{e}(2) + u_{32}\bar{e}(3), \quad \bar{e}(3). \end{aligned}$$

Then a basis for $H_{\mathcal{I}}(\mathcal{F}, \mathcal{G})$ is on the open set where $u_{31}u_{32} \neq 0$ given by

$$\begin{aligned} \phi_1 &= \begin{pmatrix} -z_{21}u_{32} & u_{32} & 0 \\ -z_{21}(u_{32}u_{21} - u_{31}) & u_{32}u_{21} - u_{31} & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\ \phi_2 &= \begin{pmatrix} z_{31}u_{32} & 0 & 0 \\ z_{31}(u_{32}u_{21} - u_{31}) & 0 & u_{31} \\ 0 & 0 & u_{32}u_{31} \end{pmatrix}, \quad \phi_3 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & u_{21} \\ 0 & 0 & u_{31} \end{pmatrix} \end{aligned}$$

as can be checked by manual inspection.

5.2. Kernel dimension and position. Let us consider a tuple $\mathcal{I} \in \text{Subsets}(r, n, s)$, where we always assume that $r \in [n]$. To prove sufficiency of the Horn inequalities, we aim to use Corollary 5.1.9, which states that $\text{tdim } \mathcal{I} \geq \text{edim } \mathcal{I}$, with equality if and only if \mathcal{I} is intersecting.

If $\text{tdim } \mathcal{I} = 0$ then, necessarily, $\text{tdim } \mathcal{I} = \text{edim } \mathcal{I} = 0$, since $\text{edim } \mathcal{I}$ is nonnegative by assumption (part of the Horn inequalities). Hence in this case \mathcal{I} is intersecting.

Thus the interesting case is when $\text{tdim } \mathcal{I} > 0$. To study the spaces $H_{\mathcal{I}}(\mathcal{F}, \mathcal{G})$ in a unified fashion, we consider the space

$$P(\mathcal{I}) := \{(\mathcal{F}, \mathcal{G}, \phi) \in \text{Flag}(V_0)^s \times \text{Flag}(Q_0)^s \times \text{Hom}(V_0, Q_0) : \phi \in H_{\mathcal{I}}(\mathcal{F}, \mathcal{G})\}.$$

We caution that $P(\mathcal{I})$ is not in general irreducible, as the following example shows:

Example. Let $s = 2, n = 3, r = 1$, and consider $I_1 = I_2 = \{2\}$. There is only a single flag on $V_0 \cong \mathbb{C}$, while any flag G on $Q_0 \cong \mathbb{C}^2$ is determined by a line $L = G(1) \in \mathbb{P}(\mathbb{C}^2)$. Thus we can identify $P(\mathcal{I}) \cong \{(L_1, L_2, \phi) \in \mathbb{P}(\mathbb{C}^2)^2 \times \text{Hom}(\mathbb{C}^1, \mathbb{C}^2) : \phi(e(1)) \in L_1 \cap L_2\}$. If we consider the map $(L_1, L_2, \phi) \mapsto (L_1, L_2)$, then the fiber for any $L_1 = L_2$ is a one-dimensional line, while for any $L_1 \neq L_2$ the fiber is just $\phi = 0$. In particular, we note that $P(\mathcal{I})$ is not irreducible.

We now restrict to those $(\mathcal{F}, \mathcal{G})$ such that the intersection $H_{\mathcal{I}}(\mathcal{F}, \mathcal{G})$ is of dimension $\text{tdim } \mathcal{I}$. Thus we introduce

$$\begin{aligned} P_t(\mathcal{I}) &:= \{(\mathcal{F}, \mathcal{G}, \phi) \in P(\mathcal{I}) : \dim H_{\mathcal{I}}(\mathcal{F}, \mathcal{G}) = \text{tdim } \mathcal{I}\}, \\ B_t(\mathcal{I}) &:= \{(\mathcal{F}, \mathcal{G}) \in \text{Flag}(V_0)^s \times \text{Flag}(Q_0)^s : \dim H_{\mathcal{I}}(\mathcal{F}, \mathcal{G}) = \text{tdim } \mathcal{I}\}. \end{aligned}$$

The subscripts in $P_t(\mathcal{I})$ and $B_t(\mathcal{I})$ stands for the true dimension, $\text{tdim } \mathcal{I}$. We use similar subscripts throughout this section when we fix various other dimensions and positions.

Since $\text{tdim } \mathcal{I}$ is the minimal possible dimension, this is the generic case. Moreover, this restriction makes $P_t(\mathcal{I})$ irreducible, as it is a vector bundle over $B_t(\mathcal{I})$. We record this in the following lemma:

Lemma 5.2.1. *The space $P(\mathcal{I})$ is a closed subvariety of $\text{Flag}(V_0)^s \times \text{Flag}(Q_0)^s \times \text{Hom}(V_0, Q_0)$, and $P_t(\mathcal{I})$ is a nonempty Zariski-open subset of $P(\mathcal{I})$. Moreover, $B_t(\mathcal{I})$ is a nonempty Zariski-open subset of $\text{Flag}(V_0)^s \times \text{Flag}(Q_0)^s$, and the map $(\mathcal{F}, \mathcal{G}, \phi) \mapsto (\mathcal{F}, \mathcal{G})$ turns $P_t(\mathcal{I})$ into a vector bundle over $B_t(\mathcal{I})$. In particular, $P_t(\mathcal{I})$ is an irreducible and smooth variety.*

In particular:

$$(5.2.2) \quad \dim P_t(\mathcal{I}) = s(\dim \text{Flag}(V_0) + \dim \text{Flag}(Q_0)) + \text{tdim } \mathcal{I}$$

Belkale’s insight is now to consider the behavior of generic kernels of maps $\phi \in H_{\mathcal{I}}(\mathcal{F}, \mathcal{G})$, where $(\mathcal{F}, \mathcal{G}) \in B_t(\mathcal{I})$. We start with the following definition:

Definition 5.2.3. Let $\mathcal{I} \in \text{Subsets}(r, n, s)$. We define the *kernel dimension* of \mathcal{I} as

$$\text{kdim } \mathcal{I} := \min \{ \dim \ker \phi : \phi \in H_{\mathcal{I}}(\mathcal{F}, \mathcal{G}) \text{ where } (\mathcal{F}, \mathcal{G}) \in B_t(\mathcal{I}) \}$$

There are two special cases that we can treat right away. If $\text{kdim } \mathcal{I} = r$ then any morphism in $H_{\mathcal{I}}(\mathcal{F}, \mathcal{G})$ for $(\mathcal{F}, \mathcal{G}) \in B_t(\mathcal{I})$ is zero, and hence $\text{tdim } \mathcal{I} = 0$. This is the case that we had discussed initially and we record this observation for future reference:

Lemma 5.2.4. *Let $\mathcal{I} \in \text{Subsets}(r, n, s)$ such that $\text{edim } \mathcal{I} \geq 0$. If $\text{kdim } \mathcal{I} = r$ then $\text{tdim } \mathcal{I} = \text{edim } \mathcal{I} = 0$, and hence $\mathcal{I} \in \text{Intersecting}(r, n, s)$.*

Likewise, the case where $\text{kdim } \mathcal{I} = 0$ can easily be treated directly. The idea is to compute the dimension of $P_t(\mathcal{I})$ in a second way and compare the result with (5.2.2).

Lemma 5.2.5. *Let $\mathcal{I} \in \text{Subsets}(r, n, s)$. If $\text{kdim } \mathcal{I} = 0$ then*

$$\dim P_t(\mathcal{I}) = s(\dim \text{Flag}(V_0) + \dim \text{Flag}(Q_0)) + \text{edim } \mathcal{I}.$$

Proof. We first note that $\text{kdim } \mathcal{I} = 0$ implies that there exists an injective map $\phi \in H_{\mathcal{I}}(\mathcal{F}, \mathcal{G})$ for some $(\mathcal{F}, \mathcal{G}) \in B_t(\mathcal{I})$. In particular, $I_k(a) - a \geq a$ for all $k \in [s]$ and $a \in [r]$ (a fact that we use further below in the proof). Now define

$$P_k := \{(\mathcal{F}, \mathcal{G}, \phi) \in P(\mathcal{I}) : \dim \ker \phi = 0\}$$

Then P_k is a nonempty Zariski-open subset of $P(\mathcal{I})$ that intersects $P_t(\mathcal{I})$. By Lemma 5.2.1, the latter is irreducible. Thus it suffices to show that P_k is likewise irreducible and to compute its dimension.

For this, we consider the map

$$\pi : P_k \rightarrow M_k := \text{Flag}(V_0)^s \times \text{Hom}^\times(V_0, Q_0), \quad (\mathcal{F}, \mathcal{G}, \phi) \mapsto (\mathcal{F}, \phi)$$

where we write $\text{Hom}^\times(V_0, Q_0)$ for the Zariski-open subset of injective linear maps in $\text{Hom}(V_0, Q_0)$. The fibers of π are given by

$$\pi^{-1}(\mathcal{F}, \phi) \cong \prod_{k=1}^s \text{Flag}_{I_k}^0(F_k, \phi)$$

which according to Lemma 3.3.10 are smooth irreducible varieties of dimension $s \dim \text{Flag}(Q_0) - sr(n-r) + \sum_{k=1}^s \dim I_k$. It is not hard to see that π gives P_k the structure of a fiber bundle over M_k . Therefore, P_k is irreducible. Moreover, the space M_k has dimension $s \dim \text{Flag}(V_0) + r(n-r)$. By adding the dimension of the fibers, we obtain that the dimension of P_k , and hence of $P_t(\mathcal{I})$, is indeed the one claimed in the lemma. □

Corollary 5.2.6. *Let $\mathcal{I} \in \text{Subsets}(r, n, s)$. If $\text{kdim } \mathcal{I} = 0$ then $\text{tdim } \mathcal{I} = \text{edim } \mathcal{I}$, and hence $\mathcal{I} \in \text{Intersecting}(r, n, s)$.*

Proof. This follows directly by comparing Eq. (5.2.2) and Lemma 5.2.5. □

We now consider the general case, where $0 < d := \text{kdim } \mathcal{I} < r$. We first note that the kernel dimension is attained generically. Thus we define

$$P_{kt}(\mathcal{I}) := \{(\mathcal{F}, \mathcal{G}, \phi) \in P_t(\mathcal{I}) : \dim \ker \phi = \text{kdim } \mathcal{I}\},$$

$$B_{kt}(\mathcal{I}) := \{(\mathcal{F}, \mathcal{G}) : \exists \phi \text{ s.th. } (\mathcal{F}, \mathcal{G}, \phi) \in P_{kt}(\mathcal{I})\} \subseteq B_t(\mathcal{I}),$$

where the subscripts denote that we fix both the true dimension as well as the kernel dimension. We have the following lemma:

Lemma 5.2.7. *The set $P_{kt}(\mathcal{I})$ is a nonempty Zariski-open subset of $P_t(\mathcal{I})$, hence also irreducible. Moreover, $B_{kt}(\mathcal{I})$ is a nonempty Zariski-open subset of $\text{Flag}(V_0)^s \times \text{Flag}(Q_0)^s$.*

Proof. The first claim holds since $P_{kt}(\mathcal{I})$ can be defined by the nonvanishing of certain minors. The second claim now follows as $B_{kt}(\mathcal{I})$ is the image of the Zariski-open subset $P_{kt}(\mathcal{I})$ of the vector bundle $P_t(\mathcal{I}) \rightarrow B_t(\mathcal{I})$. \square

Belkale’s insight is to consider the positions of generic kernels for an induction:

Definition 5.2.8. Let $\mathcal{I} \in \text{Subsets}(r, n, s)$. Then we define the *kernel position* of \mathcal{I} as the tuple $\mathcal{J} \in \text{Subsets}(d, r, s)$ defined by

$$J_k(b) := \min \{ \text{Pos}(\ker \phi, F_k)(b) : (\mathcal{F}, \mathcal{G}, \phi) \in P_{kt}(\mathcal{I}) \}$$

for $b \in [d]$ and $k \in [s]$. We write $k\text{Pos}(\mathcal{I}) = \mathcal{J}$.

The goal in the remainder of this subsection is to prove the following equality:

$$\text{tdim } \mathcal{I} = \text{edim } \mathcal{J} + \text{edim } \mathcal{I} / \mathcal{J},$$

where $\mathcal{J} = k\text{Pos}(\mathcal{I})$. This will again be accomplished by computing the dimension of $P_t(\mathcal{I})$ in a second way and comparing the result with (5.2.2). Specifically, we consider the spaces

$$\begin{aligned} P_{kpt}(\mathcal{I}) &:= \{ (\mathcal{F}, \mathcal{G}, \phi) \in P_{kt}(\mathcal{I}) : \text{Pos}(\ker \phi, \mathcal{F}) = k\text{Pos}(\mathcal{I}) \}, \\ B_{kpt}(\mathcal{I}) &:= \{ (\mathcal{F}, \mathcal{G}) : \exists \phi \text{ s.th. } (\mathcal{F}, \mathcal{G}, \phi) \in P_{kpt}(\mathcal{I}) \} \subseteq B_{kt}(\mathcal{I}). \end{aligned}$$

Then $P_{kpt}(\mathcal{I})$ is Zariski-open in $P_{kt}(\mathcal{I})$, since it can again be defined by demanding that certain minors are nonzero. We obtain the following lemma, the second claim in which is proved as before:

Lemma 5.2.9. *Let $\mathcal{I} \in \text{Subsets}(r, n, s)$ such that $0 < k\dim \mathcal{I} < r$. Then $P_{kpt}(\mathcal{I})$ is a nonempty Zariski-open subset of $P_{kt}(\mathcal{I})$, hence also irreducible. Moreover, $B_{kpt}(\mathcal{I})$ is a nonempty Zariski-open subset of $\text{Flag}(V_0)^s \times \text{Flag}(Q_0)^s$.*

Corollary 5.2.10. *Let $\mathcal{I} \in \text{Subsets}(r, n, s)$ such that $0 < k\dim \mathcal{I} < r$. Then $k\text{Pos}(\mathcal{I}) \in \text{Intersecting}(d, r, s)$.*

Proof. According to Lemma 5.2.9, $B_{kpt}(\mathcal{I})$ is a nonempty Zariski-open subset of $\text{Flag}(V_0)^s \times \text{Flag}(Q_0)^s$, hence Zariski-dense. It follows that its image under the projection $(\mathcal{F}, \mathcal{G}) \mapsto \mathcal{F}$ is likewise Zariski-dense. For any such \mathcal{F} , there exists a \mathcal{G} and ϕ such that $(\mathcal{F}, \mathcal{G}, \phi) \in P_{kpt}(\mathcal{I})$, and hence $\ker \phi \in \Omega_{k\text{Pos}(\mathcal{I})}^0(\mathcal{F})$; in particular, $\Omega_{k\text{Pos}(\mathcal{I})}^0(\mathcal{F})$ is nonempty. Thus Lemmas 4.2.3 and 4.2.4 show that $k\text{Pos}(\mathcal{I})$ is intersecting. \square

We now compute the dimension of $P_{\text{kpt}}(\mathcal{I})$. As in the proof of Lemma 5.2.5, it will be useful to consider an auxiliary space where we do *not* enforce the true dimension:

$$P_{\text{kp}}(\mathcal{I}) := \{(\mathcal{F}, \mathcal{G}, \phi) \in P(\mathcal{I}) : \text{Pos}(\ker \phi, \mathcal{F}) = \text{kPos}(\mathcal{I})\}$$

Note that constraint on the position of the kernel implies that its dimension is $\text{kdim } \mathcal{I}$.

Lemma 5.2.11. *Let $\mathcal{I} \in \text{Subsets}(r, n, s)$ such that $0 < \text{kdim } \mathcal{I} < r$. Then $P_{\text{kp}}(\mathcal{I})$ is nonempty, smooth, irreducible, and satisfies*

$$\dim P_{\text{kp}}(\mathcal{I}) = s(\dim \text{Flag}(V_0) + \dim \text{Flag}(Q_0)) + \text{edim } \mathcal{J} + \text{edim } \mathcal{I}/\mathcal{J},$$

where $\mathcal{J} := \text{kPos}(\mathcal{I})$.

Proof. Clearly, $P_{\text{kp}}(\mathcal{I})$ is nonempty since it contains $P_{\text{kpt}}(\mathcal{I})$. We now introduce

$$M_{\text{kp}} := \{(\mathcal{F}, \phi) \in \text{Flag}(V_0)^s \times \text{Hom}(V_0, Q_0) : \text{Pos}(\ker \phi, \mathcal{F}) = \text{kPos}(\mathcal{I})\}$$

and consider the map

$$\pi : P_{\text{kp}}(\mathcal{I}) \rightarrow M_{\text{kp}}, \quad (\mathcal{F}, \mathcal{G}, \phi) \mapsto (\mathcal{F}, \phi).$$

Its fibers are given by

$$\pi^{-1}(\mathcal{F}, \phi) \cong \prod_{k=1}^s \{G_k \in \text{Flag}(Q_0) : \phi \in H_{I_k}(F_k, G_k)\}$$

To understand the right-hand side, define $S := \ker \phi$ and let $\bar{\phi} : V_0/S \rightarrow Q_0$ the corresponding injective map. By Lemma 3.2.15, $\phi \in H_{I_k}(F_k, G_k)$ if and only if $\bar{\phi} \in H_{I_k/J_k}((F_k)_{V_0/S}, G_k)$, that is, $G_k \in \text{Flag}_{I_k/J_k}^0((F_k)_{V_0/S}, \bar{\phi})$ as introduced in Definition 3.3.9. Thus we find that the fibers of π can be identified as

$$\pi^{-1}(\mathcal{F}, \phi) \cong \prod_{k=1}^s \text{Flag}_{I_k/J_k}^0((F_k)_{V_0/S}, \bar{\phi}).$$

By Lemma 3.3.10, the k -th factor on the right-hand side is a smooth irreducible variety of dimension $\dim \text{Flag}(Q_0) - (r - d)(n - r) + \dim I_k/J_k$, where $d := \dim \ker \phi = \text{kdim } \mathcal{I}$. It is not hard to see that π is a fiber bundle, and we will show momentarily that M_{kp} is irreducible. Hence

$$(5.2.12) \quad \dim P_{\text{kp}}(\mathcal{I}) = \dim M_{\text{kp}} + s \dim \text{Flag}(Q_0) - s(r - d)(n - r) + \sum_{k=1}^s \dim I_k/J_k.$$

It remains to show that M_{kp} is smooth and irreducible and to compute its dimension. For this, we consider the map

$$\tau: M_{kp} \rightarrow \text{Gr}(d, V_0), (\mathcal{F}, \phi) \mapsto \ker \phi.$$

Since ϕ can be specified in terms of the kernel $S := \ker \phi$ and the injection $\bar{\phi}: V_0/S \rightarrow Q_0$, it is clear that the fibers of τ are given by

$$\tau^{-1}(S) = \text{Hom}^\times(V_0/S, Q_0) \times \prod_{k=1}^s \text{Flag}_{J_k}^0(S, V_0).$$

Since τ is likewise a fiber bundle, we obtain that M_{kp} is smooth and irreducible and, using (3.3.7), that

$$\begin{aligned} \dim M_{kp} &= \dim \text{Gr}(d, V_0) + (r-d)(n-r) + \sum_{k=1}^s \dim \text{Flag}_{J_k}^0(S, V_0) \\ &= d(r-d) + (r-d)(n-r) + s(\dim \text{Flag}(S) + \dim \text{Flag}(V_0/S)) \\ &\qquad\qquad\qquad + \sum_{k=1}^s \dim J_k \\ &= d(r-d)(1-s) + (r-d)(n-r) + s \dim \text{Flag}(V_0) + \sum_{k=1}^s \dim J_k. \end{aligned}$$

By plugging this result into (5.2.12) and simplifying, we obtain the desired result. □

Corollary 5.2.13. *Let $\mathcal{I} \in \text{Subsets}(r, n, s)$ such that $0 < \text{kdim } \mathcal{I} < r$, and $\mathcal{J} = \text{kPos}(\mathcal{I})$. Then,*

$$(5.2.14) \qquad \text{tdim } \mathcal{I} = \text{edim } \mathcal{J} + \text{edim } \mathcal{I}/\mathcal{J}$$

Proof. Recall that $P_{kp}(\mathcal{I}) \subseteq P(\mathcal{I}) \supseteq P_t(\mathcal{I})$. Moreover,

$$P_{kpt}(\mathcal{I}) = P_{kp}(\mathcal{I}) \cap P_t(\mathcal{I}) \subseteq P(\mathcal{I}).$$

All three varieties $P_{kpt}(\mathcal{I})$, $P_{kp}(\mathcal{I})$, $P_t(\mathcal{I})$ are irreducible (Lemmas 5.2.1, 5.2.9 and 5.2.11). Moreover, $P_{kpt}(\mathcal{I})$ is nonempty and Zariski-open in $P(\mathcal{I})$, hence in both $P_{kp}(\mathcal{I})$ and $P_t(\mathcal{I})$. It follows that

$$\dim P_{kp}(\mathcal{I}) = \dim P_{kpt}(\mathcal{I}) = \dim P_t(\mathcal{I}).$$

We now obtain (5.2.14) via Lemma 5.2.11 and Eq. (5.2.2). □

Remark. Purbhoo [Pur2] asserts that if \mathcal{J} denotes the kernel position of \mathcal{I} then \mathcal{I}/\mathcal{J} is intersecting. However, we believe that the proof given therein is incomplete, as it is not clear that the map $(\mathcal{F}, \mathcal{G}, \phi) \mapsto (\mathcal{F}_{V/S}, \mathcal{G})$ is dominant (cf. the remark at [Pur1]). The following argument suggests that the situation is somewhat more delicate.

5.3. The kernel recurrence. To conclude the proof in the case that $0 < \text{kdim } \mathcal{I} < r$, we need to understand the right-hand side of (5.2.14) some more. We start with the calculation

$$(5.3.1) \quad \text{tdim } \mathcal{I} - \text{edim } \mathcal{I} = \text{edim } \mathcal{J} - (\text{edim } \mathcal{I}\mathcal{J} - \text{edim } \mathcal{J}) = \text{edim } \mathcal{J} - \text{edim } \mathcal{I}^{\mathcal{J}},$$

where the first equality is due to Eqs. (4.2.9) and (5.2.14) and the second is Eq. (4.2.11).

The last missing ingredient is to understand the expected dimension of the kernel position, $\text{edim } \mathcal{J}$.

Lemma 5.3.2. *Let $\mathcal{I} \in \text{Subsets}(r, n, s)$ such that $0 < \text{kdim } \mathcal{I} < r$, and let $\mathcal{J} := \text{kPos}(\mathcal{I})$. Then we have $\text{edim } \mathcal{J} \leq \text{tdim } \mathcal{I}^{\mathcal{J}}$.*

Proof. For any $(\mathcal{F}, \mathcal{G}, \phi) \in \text{P}_{\text{kp}}(\mathcal{I})$, the space $H_{\mathcal{J}}(\mathcal{F}^{\text{ker } \phi}, \mathcal{F}_{V_0/\text{ker } \phi})$ injects into $H_{\mathcal{I}\mathcal{J}}(\mathcal{F}^{\text{ker } \phi}, \mathcal{G})$ by composition with the injective map $\bar{\phi}: V_0/\text{ker } \phi \rightarrow Q_0$ induced by ϕ (Lemma 3.2.15). Thus,

$$\text{edim } \mathcal{J} \leq \text{tdim } \mathcal{J} \leq \dim H_{\mathcal{J}}(\mathcal{F}^{\text{ker } \phi}, \mathcal{F}_{V_0/\text{ker } \phi}) \leq \dim H_{\mathcal{I}\mathcal{J}}(\mathcal{F}^{\text{ker } \phi}, \mathcal{G}),$$

where the first inequality is always true (Corollary 5.1.9), the second holds by definition of the true dimension and the third follows from the injection. It thus suffices to prove that there exists $(\mathcal{F}, \mathcal{G}, \phi) \in \text{P}_{\text{kp}}(\mathcal{I})$ such that $\dim H_{\mathcal{I}\mathcal{J}}(\mathcal{F}^{\text{ker } \phi}, \mathcal{G}) \leq \text{tdim } \mathcal{I}^{\mathcal{J}}$.

For this, let $K(d, V_0)$ denote the fiber bundle over $\text{Gr}(d, V_0)$ with fiber over $S \in \text{Gr}(d, V_0)$ given by $\text{Flag}(S)^s \times \text{Flag}(Q_0)^s$. It is an irreducible algebraic variety and we denote its elements by $(S, \tilde{\mathcal{F}}, \mathcal{G})$. We consider the morphism

$$\pi: \text{P}_{\text{kp}}(\mathcal{I}) \rightarrow K(d, V_0), \quad (\mathcal{F}, \mathcal{G}, \phi) \mapsto (\text{ker } \phi, \mathcal{F}^{\text{ker } \phi}, \mathcal{G}).$$

For any $(\mathcal{F}, \mathcal{G}, \phi) \in \text{P}_{\text{kp}}(\mathcal{I})$, $\dim \text{ker } \phi = d$ and $\text{Pos}(\text{ker } \phi, \mathcal{F}) = \mathcal{J}$, hence π is indeed a morphism.

We first prove that π is dominant. Note that, as a consequence of Lemma 5.2.9, the map $\text{P}_{\text{kp}}(\mathcal{I}) \rightarrow \text{Flag}(Q_0)^s, (\mathcal{F}, \mathcal{G}, \phi) \mapsto \mathcal{G}$ contains a nonempty Zariski-open subset $U \subseteq \text{Flag}(Q_0)^s$. We now show that the image of π contains all elements $(S, \tilde{\mathcal{F}}, \mathcal{G})$ with $S \in \text{Gr}(d, V_0)$, $\tilde{\mathcal{F}} \in \text{Flag}(S)^s$ and $\mathcal{G} \in U$. For this, let $(\mathcal{F}_0, \mathcal{G}, \phi_0) \in \text{P}_{\text{kp}}(\mathcal{I})$ be the preimage of some arbitrary $\mathcal{G} \in U$. Let $S_0 := \text{ker } \phi_0$ and choose some $g \in \text{GL}(V_0)$ such that $g \cdot S_0 = S$. Using the corresponding diagonal action, $\mathcal{F} := g \cdot \mathcal{F}_0$ and $\phi := g \cdot \phi_0$, we obtain that $(\mathcal{F}, \mathcal{G}, \phi) \in \text{P}_{\text{kp}}(\mathcal{I})$ and $\text{ker } \phi = S$. Given $\tilde{\mathcal{F}} \in \text{Flag}(S)^s$, we now choose $\vec{h} \in \text{GL}(V_0)^s$ such that $h_k S \subseteq S$, $h_k \cdot F_k^S = \tilde{F}_k$, and h_k acts trivially on V_0/S for all $k \in [s]$. Then $\text{Pos}(S, \vec{h} \cdot \mathcal{F}) = \text{Pos}(S, \mathcal{F}) = \mathcal{J}$, which shows that $(\vec{h} \cdot \mathcal{F})^S = \vec{h} \cdot \mathcal{F}^S = \tilde{\mathcal{F}}$.

Moreover, $(\vec{h} \cdot \mathcal{F})_{V_0/S} = \mathcal{F}_{V_0/S}$. Thus $\phi \in H_{\mathcal{I}}(\mathcal{F}, \mathcal{G})$ implies that $\phi \in H_{\mathcal{I}}(\vec{h} \cdot \mathcal{F}, \mathcal{G})$ by Lemma 3.2.15. Together, we find that the triple $(\vec{h} \cdot \mathcal{F}, \mathcal{G}, \phi)$ is in $\text{P}_{\text{kp}}(\mathcal{I})$ and mapped by π to $(S, \tilde{\mathcal{F}}, \mathcal{G})$. We thus obtain that π is dominant.

To conclude the proof, we note that the subset $W \subseteq K(d, V_0)$ consisting of those $(S, \tilde{\mathcal{F}}, \mathcal{G})$ with $\dim H_{\mathcal{I}^{\mathcal{J}}}(\tilde{\mathcal{F}}, \mathcal{G}) = \text{tdim } \mathcal{I}^{\mathcal{J}}$ is a nonempty Zariski-open subset, and hence Zariski-dense since $K(d, V_0)$ is irreducible. For each fixed choice of S , this is the claim in Lemma 5.2.1 for $B_t(\mathcal{I})$, with $\mathcal{I}^{\mathcal{J}}$ instead of \mathcal{I} . The ‘parametrized version’ is proved in the same way. Since π is dominant, the preimage $\pi^{-1}(W)$ is a nonempty Zariski-open subset of $\text{P}_{\text{kp}}(\mathcal{I})$. In particular, any $(\mathcal{F}, \mathcal{G}, \pi) \in \pi^{-1}(W) \subseteq \text{P}_{\text{kp}}(\mathcal{I})$ satisfies $\dim H_{\mathcal{I}^{\mathcal{J}}}(\mathcal{F}^S, \mathcal{G}) \leq \text{tdim } \mathcal{I}^{\mathcal{J}}$. \square

We thus obtain the following fundamental recurrence relation, due to Sherman [She], as a consequence of Eq. (5.3.1) and Lemma 5.3.2:

$$(5.3.3) \quad \text{tdim } \mathcal{I} - \text{edim } \mathcal{I} \leq \text{tdim } \mathcal{I}^{\mathcal{J}} - \text{edim } \mathcal{I}^{\mathcal{J}}$$

Now we have assembled all ingredients to prove Belkale’s theorem:

Theorem 5.3.4 (Belkale [Bel3], restated).

For $r \in [n]$ and $s \geq 2$, $\text{Intersecting}(r, n, s) = \text{Horn}(r, n, s)$.

Proof. We proceed by induction on r . The base case, $r = 1$, is Example 4.3.12. Thus we have $\text{Intersecting}(1, n, s) = \text{Horn}(1, n, s)$ for all $n \geq 1$.

Now let $r > 1$. By the induction hypothesis, $\text{Horn}(d, n', s) = \text{Intersecting}(d, n', s)$ for all $0 < d < r$ and $d \leq n'$. In particular, $\text{Horn}(r, n, s)$ from Definition 2.18 can be written in the following form:

$$\begin{aligned} & \text{Horn}(r, n, s) \\ &= \{ \mathcal{I} : \text{edim } \mathcal{I} \geq 0, \forall \mathcal{J} \in \text{Intersecting}(d, r, s), 0 < d < r, \\ & \hspace{20em} \text{edim } \mathcal{J} = 0 : \text{edim } \mathcal{I}\mathcal{J} \geq 0 \} \\ &= \{ \mathcal{I} : \text{edim } \mathcal{I} \geq 0, \forall \mathcal{J} \in \text{Intersecting}(d, r, s), 0 < d < r, \text{edim } \mathcal{I}\mathcal{J} \geq \text{edim } \mathcal{J} \} \end{aligned}$$

where the second equality is due to Proposition 4.3.10. Hence it is a direct consequence of Corollary 2.17 that $\text{Intersecting}(r, n, s) \subseteq \text{Horn}(r, n, s)$. We now prove the converse.

Thus let $\mathcal{I} \in \text{Horn}(r, n, s)$. Let $d := \text{kdim } \mathcal{I}$. If $d = 0$ or $d = r$ then we know from Lemma 5.2.4 and Corollary 5.2.6, respectively, that \mathcal{I} is intersecting. We now discuss the case where $0 < d < r$. By Eq. (5.3.3), we have that

$$\text{tdim } \mathcal{I} - \text{edim } \mathcal{I} \leq \text{tdim } \mathcal{I}^{\mathcal{J}} - \text{edim } \mathcal{I}^{\mathcal{J}},$$

where $\mathcal{J} := \text{kPos}(\mathcal{I})$ denotes the kernel position of \mathcal{I} . If we can show that $\mathcal{I}^{\mathcal{J}}$ is intersecting then the right-hand side is zero by Corollary 5.1.9, hence so is the

left-hand side, since $\text{tdim } \mathcal{I} - \text{edim } \mathcal{I} \geq 0$, and thus \mathcal{I} is intersecting, which is what we set out to prove.

To see that $\mathcal{I}^{\mathcal{J}}$ is intersecting, we note that $\text{Intersecting}(d, n - r + d, s) = \text{Horn}(d, n - r + d, s)$ by the induction hypothesis, hence it remains to verify that $\mathcal{I}^{\mathcal{J}}$ satisfies the Horn inequalities. Let $\mathcal{K} \in \text{Horn}(m, d, s) = \text{Intersecting}(m, d, s)$ for any $0 < m \leq d$, where we have used the induction hypothesis one last time. Thus $\mathcal{JK} \in \text{Intersecting}(m, r, s)$ by Corollary 5.2.10 and Lemma 2.16. It follows that

$$\text{edim } \mathcal{I}^{\mathcal{J}} \mathcal{K} - \text{edim } \mathcal{K} = \text{edim } \mathcal{I}(\mathcal{JK}) - \text{edim } \mathcal{JK} \geq 0$$

where the first step is (4.2.10) and the second step holds because by assumption $\mathcal{I} \in \text{Horn}(r, n, s)$ and $\mathcal{JK} \in \text{Intersecting}(m, r, s) = \text{Horn}(m, r, s)$, as explained above. We remark that these inequalities include $\text{edim } \mathcal{I}^{\mathcal{J}} \geq 0$ (corresponding to $m = d$). Thus we have shown that $\mathcal{I}^{\mathcal{J}}$ satisfies the Horn inequalities. This is what remained to be proved. \square

6. Invariants and Horn inequalities

In this section, we show that the Horn inequalities not only characterize intersections, but also the existence of corresponding nonzero invariants and, thereby, the Kirwan cone for the eigenvalues of sums of Hermitian matrices.

6.1. Borel–Weil construction. For any dominant weight $\lambda \in \Lambda_+(r)$ there exists an irreducible representation $L(\lambda)$ of $\text{GL}(r)$ with highest weight λ , unique up to isomorphism. Following Borel and Weil, it can be constructed as follows:

For any weight $\mu \in \Lambda(r)$, let us denote by $\chi_\mu: B(r) \rightarrow \mathbb{C}^*$ the character of $B(r)$ such that $\chi_\mu(t) = t^\mu = t(1)^{\mu(1)} \dots t(r)^{\mu(r)}$ for all $t \in H(r) \subseteq B(r)$. Here, we recall that $B(r)$ is the group of upper-triangular invertible matrices and $H(r) \subseteq B(r)$ the Cartan subgroup, which consists of invertible matrices $t \in \text{GL}(r)$ that are diagonal in the standard basis, with diagonal entries $t(1), \dots, t(r)$. Lastly, we write $\mathbb{1}_r = (1, \dots, 1) \in \Lambda(r)$ for the highest weight of the determinant representation of $\text{GL}(r)$, denoted \det_r . It is clear that $L(\lambda + k\mathbb{1}_r) = L(\lambda) \otimes \det_r^k$ for any $\lambda \in \Lambda_+(r)$ and $k \in \mathbb{Z}$.

Definition 6.1.1. Let $\lambda \in \Lambda_+(r)$. Then we define the *Borel–Weil realization* of $L(\lambda)$ as

$$L_{BW}(\lambda) = \left\{ s: \text{GL}(r) \rightarrow \mathbb{C} \text{ holomorphic} : \begin{aligned} s(gb) &= s(g)\chi_{\lambda^*}(b) \\ \forall g \in \text{GL}(r), b \in B(r) \end{aligned} \right\}$$

with the action of $GL(r)$ given by $(g \cdot s)(h) := s(g^{-1}h)$. We recall that $\lambda^* = (-\lambda(r), \dots, -\lambda(1))$.

The Borel–Weil theorem asserts that $L_{BW}(\lambda)$ is an irreducible $GL(r)$ -representation of highest weight λ . Note that, by definition, a holomorphic function is in $L_{BW}(\lambda)$ if it is a highest weight vector of weight λ^* with respect to the *right* multiplication representation, $(g \star s)(h) := s(hg)$.

The space $L_{BW}(\lambda)$ can also be interpreted as the space of holomorphic sections of the $GL(r)$ -equivariant line bundle $\mathcal{L}_{BW}(\lambda) := GL(r) \times_{B(r)} \mathbb{C}_{-\lambda^*}$ over $\text{Flag}(r) \cong GL(r)/B(r)$, where we write \mathbb{C}_μ for the one-dimensional representation of $B(r)$ given by the character χ_μ .

It is useful to observe that we have a $GL(r)$ -equivariant isomorphism

$$(6.1.2) \quad L(\lambda)^* \rightarrow L_{BW}(\lambda^*), \quad f \mapsto (s_f: GL(r) \rightarrow \mathbb{C}, g \mapsto f(g \cdot v_\lambda))$$

where v_λ denotes a fixed highest weight vector in $L(\lambda)$.

The tensor product of several Borel–Weil representations can again be identified with a space of functions. E.g., if $\lambda \in \Lambda_+(r)$ and $\lambda' \in \Lambda_+(r')$ then

$$\begin{aligned} &L_{BW}(\lambda) \otimes L_{BW}(\lambda') \\ &\cong \{s: GL(r) \times GL(r') \rightarrow \mathbb{C} \text{ holomorphic, } s(gb, g'b') = s(g, g')\chi_{\lambda^*}(b)\chi_{\lambda'^*}(b') \\ &\quad \forall g \in GL(r), g' \in GL(r'), b \in B(r), b' \in B(r')\}. \end{aligned}$$

We will use this below to obtain a nonzero vector in a tensor product space by exhibiting a corresponding holomorphic function with the appropriate equivariance properties.

6.2. Invariants from intersecting tuples. Let us consider the tangent map (5.1.2),

$$\Delta_{\mathcal{I}, \vec{g}, \vec{h}}: \begin{cases} \text{Hom}(V_0, Q_0) \times H_{I_1}(F_0, G_0) \times \dots \times H_{I_s}(F_0, G_0) \rightarrow \text{Hom}(V_0, Q_0)^s \\ (\zeta, \phi_1, \dots, \phi_s) \mapsto (\zeta + h_1\phi_1g_1^{-1}, \dots, \zeta + h_s\phi_s g_s^{-1}) \end{cases}$$

If $\text{edim } \mathcal{I} = 0$ then (5.1.5) implies that the dimension of the domain and target space are the same. Thus we may consider the determinant of $\Delta_{\mathcal{I}, \vec{g}, \vec{h}}$, as in the following definition:

Definition 6.2.1. Let $\mathcal{I} \in \text{Subsets}(r, n, s)$ such that $\text{edim } \mathcal{I} = 0$. Then we define the *determinant function* as the holomorphic function

$$\delta_{\mathcal{I}}: \begin{cases} GL(r)^s \times GL(n-r)^s \rightarrow \mathbb{C}, \\ (\vec{g}, \vec{h}) \mapsto \det \Delta_{\mathcal{I}, \vec{g}, \vec{h}} \end{cases}$$

where the determinant is evaluated with respect to two arbitrary bases.

Here, and throughout the following, we identify $V_0 \cong \mathbb{C}^r$ and $Q_0 \cong \mathbb{C}^{n-r}$, so that $\mathrm{GL}(V_0) \cong \mathrm{GL}(r)$ and $\mathrm{GL}(Q_0) \cong \mathrm{GL}(n-r)$ and the discussion in Section 6.1 is applicable.

If \mathcal{I} is intersecting then also $\mathrm{tdim} \mathcal{I} = \mathrm{edim} \mathcal{I}$ by Corollary 5.1.9. Hence by (5.1.8) there exist \vec{g}, \vec{h} such that $\delta_{\mathcal{I}}(\vec{g}, \vec{h}) \neq 0$. That is, $\delta_{\mathcal{I}}$ is a nonvanishing holomorphic function of $\mathrm{GL}(r)^s \times \mathrm{GL}(n-r)^s$. Our goal is to show that $\delta_{\mathcal{I}}$ can be interpreted as an invariant in a tensor product of irreducible $\mathrm{GL}(r) \times \mathrm{GL}(n-r)$ -representations.

We now consider the representation of $\mathrm{GL}(r) \times \mathrm{GL}(n-r)$ on $\mathrm{Hom}(V_0, Q_0)$ given by $(a, d) \cdot \phi := d\phi a^{-1}$. Since $\mathrm{Hom}(V_0, Q_0) = V_0^* \otimes Q_0$, it is clear that for $g \in \mathrm{GL}(V_0)$, $g' \in \mathrm{GL}(Q_0)$,

$$(6.2.2) \quad \det(\mathrm{Hom}(V_0, Q_0) \ni \phi \mapsto g' \phi g^{-1} \in \mathrm{Hom}(V_0, Q_0)) \\ = \det(g)^{-(n-r)} \det(g')^r.$$

We now restrict to the subspaces $H_I(F_0, G_0)$:

Lemma 6.2.3. *Let $I \subseteq [n]$ be a subset of cardinality r . Then $H_I(F_0, G_0) \subseteq \mathrm{Hom}(V_0, Q_0)$ is $B(r) \times B(n-r)$ -stable. Furthermore, for $b \in B(r)$ and $b' \in B(n-r)$ we have*

$$\det(H_I(F_0, G_0) \ni \phi \mapsto b' \phi b^{-1} \in H_I(F_0, G_0)) = \chi_{\lambda_I}(b) \chi_{\lambda_{I^c} + r \mathbb{1}_{n-r}}(b'),$$

where we recall that λ_I was defined in Definition 4.3.7.

Proof. For the first claim, we use Lemma 3.2.7: Since the flag F_0 is stabilized by $B(r)$ and the flag G_0 is stabilized by $B(n-r)$, it is clear that $H_I(F_0, G_0)$ is stable under the action of $B(r) \times B(n-r)$.

For the second claim, we note that unipotent elements always act by representation matrices of determinant one. Hence it suffices to verify the formula for the determinant for $t \in H(r)$ and $t' \in H(n-r)$. For this, we work in the weight basis of $H_I(F_0, G_0)$ given by the elementary matrices $E_{b,a}$ that send $e(a) \mapsto \bar{e}(b)$, where $a \in [r]$ and $b \in [I(a) - a]$, and all other basis vectors to zero. Then:

$$\det(H_I(F_0, G_0) \ni \phi \mapsto t' \phi t^{-1} \in H_I(F_0, G_0)) \\ = \prod_{a=1}^r \prod_{b=1}^{I(a)-a} t'(b) t(a)^{-1} = \left(\prod_{a=1}^r t(a)^{a-I(a)} \right) \left(\prod_{b=1}^{n-r} t'(b)^{r-\#\{a: I(a)-a < b\}} \right) \\ = t^{\lambda_I} t'^{r \mathbb{1}_{n-r} + \lambda_{I^c}},$$

where we have used (4.3.8) in the last step. □

We now show that the $\delta_{\mathcal{I}}$ can be interpreted as an invariant:

Theorem 6.2.4. *Let $\mathcal{I} \in \text{Subsets}(r, n, s)$ such that $\text{edim } \mathcal{I} = 0$, and let $\delta_{\mathcal{I}}$ denote the corresponding determinant function (Definition 6.2.1). Then $\delta_{\mathcal{I}}$ belongs to $\otimes_{k=1}^s (L_{BW}(\lambda_{I_k}^*) \otimes L_{BW}(\lambda_{I_k^c}^* - r \mathbb{1}_{n-r}))$. Moreover, it transforms under the diagonal action of $\text{GL}(r) \times \text{GL}(n-r)$ by the character $\det_r^{(n-r)(s-1)} \otimes \det_{n-r}^{r(1-s)}$.*

Proof. For the first claim, we note that if $\vec{g}' \in B(r)^s$, $\vec{h}' \in B(n-r)^s$ then we can write $\Delta_{\mathcal{I}, \vec{g}, \vec{g}', \vec{h}, \vec{h}'}$ as a composition of $\Delta_{\mathcal{I}, \vec{g}, \vec{h}}$ with the automorphisms on $H_{I_k}(F_0, G_0)$ that send $\phi_k \mapsto h'_k \phi_k (g'_k)^{-1}$. Using Lemma 6.2.3, we obtain

$$\delta_{\mathcal{I}}(\vec{g}, \vec{g}', \vec{h}, \vec{h}') = \delta_{\mathcal{I}}(\vec{g}, \vec{h}) \prod_{k=1}^s \chi_{\lambda_{\mathcal{I}}}(g'_k) \chi_{\lambda_{I_k^c} + r \mathbb{1}_{n-r}}(h'_k).$$

In view of the discussion at the end of Section 6.1 this establishes the first claim.

For the second claim, let $g \in \text{GL}(r)$ and $g' \in \text{GL}(n-r)$. Thus $\Delta_{\mathcal{I}, g^{-1}\vec{g}, g'^{-1}\vec{h}}$ maps $(\zeta, \phi_1, \dots, \phi_s)$ to

$$\begin{aligned} &(\zeta + g'^{-1}h_1\phi_1g_1^{-1}g, \dots, \zeta + g'^{-1}h_s\phi_s g_s^{-1}g) \\ &= g'^{-1}(g'\zeta g^{-1} + h_1\phi_1g_1^{-1}, \dots, g'\zeta g^{-1} + h_s\phi_s g_s^{-1})g \end{aligned}$$

Thus we can write $\Delta_{\mathcal{I}, g^{-1}\vec{g}, g'^{-1}\vec{h}}$ as a composition of three maps: The automorphism $\zeta \mapsto g'\zeta g^{-1}$ of $\text{Hom}(V_0, Q_0)$, the map $\Delta_{\mathcal{I}, \vec{g}, \vec{h}}$ and the automorphism $\vec{\phi} \mapsto g'^{-1}\vec{\phi}g$ on $\text{Hom}(V_0, Q_0)^s$. Thus, using Eq. (6.2.2),

$$((g, g') \cdot \delta_{\mathcal{I}})(\vec{g}, \vec{h}) = \delta_{\mathcal{I}}(g^{-1}\vec{g}, g'^{-1}\vec{h}) = \det(g)^{-(n-r)(1-s)} \det(g')^{r(1-s)} \delta_{\mathcal{I}}(\vec{g}, \vec{h}),$$

which establishes the second claim. □

If \mathcal{I} is intersecting then we had argued before that $\delta_{\mathcal{I}}$ is nonzero. By dualizing and simplifying, we obtain the following corollary of Theorem 6.2.4:

Corollary 6.2.5. *Let $\mathcal{I} \in \text{Intersecting}(r, n, s)$ and $\text{edim } \mathcal{I} = 0$. Then,*

$$\left(\det_r^{(s-1)(n-r)} \otimes \otimes_{k=1}^s L(\lambda_{I_k}) \right)^{\text{GL}(r)} \neq 0 \quad \text{and} \quad \left(\det_{n-r}^r \otimes \otimes_{k=1}^s L(\lambda_{I_k^c}) \right)^{\text{GL}(n-r)} \neq 0.$$

Let us correspondingly define

$$c(\mathcal{I}) := \dim \left(\det_r^{(s-1)(n-r)} \otimes \otimes_{k=1}^s L(\lambda_{I_k}) \right)^{\text{GL}(r)}.$$

Then Corollary 6.2.5 states that, if \mathcal{I} is intersecting and $\text{edim}\mathcal{I} = 0$ then $c(\mathcal{I}) > 0$. This relationship between generic intersections of Schubert cells and tensor product multiplicities can be made quantitative. While we do not use this in the following Section 6.3 to describe the Kirwan cone and prove the saturation property for tensor product multiplicities, we will give a brief sketch later on in Section 6.4 and use it to establish the Fulton conjecture.

6.3. Kirwan cone and saturation. We now show that the existence of nonzero invariants is characterized by the Horn inequalities. For this, recall that we defined $c(\vec{\lambda})$ as the dimension of the space of $\text{GL}(r)$ -invariants in the tensor product $\bigotimes_{k=1}^s L(\lambda_k)$. Thus, if we define $\lambda_k = \lambda_{I_k} + (n-r)\mathbb{1}_r$ for $k \in [s-1]$ and $\lambda_s = \lambda_{I_s}$, then Corollary 6.2.5 shows that

$$(6.3.1) \quad c(\vec{\lambda}) = c(\mathcal{I}) > 0$$

whenever \mathcal{I} is intersecting and $\text{edim}\mathcal{I} = 0$. Here, we have somewhat arbitrarily selected the first $s-1$ highest weights $\lambda_1, \dots, \lambda_{s-1}$ to have nonnegative entries no larger than $n-r$, while λ_s has nonpositive entries no smaller than $r-n$. Conversely, any s -tuple of highest weights $\vec{\lambda}$ with these properties can be obtained in this way from some $\mathcal{I} \in \text{Subsets}(r, n, s)$ (recall discussion below Definition 4.3.7).

Proposition 6.3.2. *Let $\vec{\lambda} \in \Lambda_+(r)^s$ be an s -tuple such that $\sum_{k=1}^s |\lambda_k| = 0$, and for any $0 < d < r$ and any s -tuple $\mathcal{J} \in \text{Horn}(d, r, s)$ with $\text{edim}\mathcal{J} = 0$ we have that $\sum_{k=1}^s (T_{J_k}, \lambda_k) \leq 0$. Then $c(\vec{\lambda}) > 0$.*

Proof. By adding/removing suitable multiples of $\mathbb{1}_r$, the highest weight of the determinant representation, we may assume that $\lambda_1(r), \dots, \lambda_{s-1}(r) \geq 0$ and $\lambda_s(1) \leq 0$. Let $n := r + q$, where $q := \max\{\lambda_1(1), \dots, \lambda_{s-1}(1), -\lambda_s(r)\}$. Then $\vec{\lambda}$ is associated to an s -tuple $\mathcal{I} \in \text{Subsets}(r, n, s)$ as in Lemma 4.3.9.

We now show that $\text{edim}\mathcal{I} = 0$ and that \mathcal{I} is intersecting. The former follows from the first statement in Lemma 4.3.9, which gives that $\text{edim}\mathcal{I} = -\sum_{k=1}^s |\lambda_k| = 0$. To see that \mathcal{I} is intersecting, we may use Theorem 5.3.4 and show instead that \mathcal{I} satisfies the Horn inequalities $\text{edim}\mathcal{I}\mathcal{J} \geq 0$ for any $\mathcal{J} \in \text{Horn}(d, r, s)$ with $\text{edim}\mathcal{J} = 0$ and $0 < d < r$. But the second statement in Lemma 4.3.9 implies that these are equivalent to the linear inequalities $\sum_{k=1}^s (T_{J_k}, \lambda_k) \leq 0$, which hold by assumption. Thus \mathcal{I} is indeed intersecting and satisfies $\text{edim}\mathcal{I} = 0$. Now (6.3.1) shows that $c(\vec{\lambda}) = c(\mathcal{I}) > 0$. \square

At last we can prove the saturation property and characterize of the Kirwan cone in terms of Horn inequalities.

Corollary 6.3.3 (Knutson–Tao, [KT], restated). (a) *Horn inequalities:* The Kirwan cone $\text{Kirwan}(r, s)$ is the convex polyhedral cone of $\vec{\xi} \in C_+(r)^s$ such that $\sum_{k=1}^s |\xi_k| = 0$, and for any $0 < d < r$ and any s -tuple $\mathcal{J} \in \text{Horn}(d, r, s)$ with $\text{edim } \mathcal{J} = 0$ we have that $\sum_{k=1}^s (T_{J_k}, \xi_k) \leq 0$.

(b) *Saturation property:* For a dominant weight $\vec{\lambda} \in \Lambda_+(r)^s$, the space of invariants $(L(\lambda_1) \otimes \cdots \otimes L(\lambda_s))^{\text{GL}(r)}$ is nonzero if and only if $\vec{\lambda} \in \text{Kirwan}(r, s)$.

In particular, $c(\vec{\lambda}) := \dim(L(\lambda_1) \otimes \cdots \otimes L(\lambda_s))^{\text{GL}(r)} > 0$ if and only if $c(N\vec{\lambda}) > 0$ for some integer $N > 0$.

Proof. The two statements are closely interlinked. For clarity, we give separate proofs that do not refer to each other.

(a) Any $\vec{\xi} \in \text{Kirwan}(r, s)$ satisfies the Horn inequalities (Corollary 2.13). We now observe that $\text{Kirwan}(r, s)$ is a closed subset of $C_+(r)^s$ which, moreover, is invariant under rescaling by nonnegative real numbers. Thus it suffices to prove the converse only for $\vec{\lambda} \in \Lambda_+(r)^s$. For this, we use that if $\vec{\lambda}$ satisfies the Horn inequalities then $c(\vec{\lambda}) > 0$ by Proposition 6.3.2, hence $\vec{\lambda} \in \text{Kirwan}(r, s)$ by Proposition 2.3.

(b) Let $\vec{\lambda} \in \Lambda_+(r)^s$. If $c(\vec{\lambda}) > 0$ then $\vec{\lambda} \in \text{Kirwan}(r, s)$ by Proposition 2.3. Conversely, if $\vec{\lambda} \in \text{Kirwan}(r, s)$ then it satisfies the Horn inequalities by Corollary 2.13, hence $c(\vec{\lambda}) > 0$ by Proposition 6.3.2.

□

Remark 6.3.4. As follows from the discussion below Proposition 4.3.10, the Kirwan cone is in fact already defined by those \mathcal{J} such that $\Omega_{\mathcal{J}}(\mathcal{G})$ is a point for all $\mathcal{G} \in \text{Good}(r, s)$. Ressayre has shown that the corresponding inequalities are irredundant and can be computed by an inductive algorithm [Res2]. Demanding that $\Omega_{\mathcal{J}}(\mathcal{G})$ is a point for all good \mathcal{G} is a more stringent requirement than $\text{edim } \mathcal{J} = 0$, and indeed the set of inequalities $\text{edim } \mathcal{I}\mathcal{J} \geq 0$ for $\mathcal{J} \in \text{Horn}(d, r, s)$ with $\text{edim } \mathcal{J} = 0$ is in general still redundant. However, from a practical point of view we prefer the latter criterion since it is much easier to check numerically.

6.4. Invariants and intersection theory. We now explain how the relationship between generic intersections of Schubert cells and tensor product multiplicities can be made more quantitative. Specifically, we shall relate the dimension $c(\mathcal{I})$ of the space of $\text{GL}(r)$ -invariants to the number of points in a generic intersection $\Omega_{\mathcal{I}}(\mathcal{E})$, as in the following definition:

Definition 6.4.1. Let $\mathcal{I} \in \text{Subsets}(r, n, s)$ such that $\text{edim } \mathcal{I} = 0$. We define the corresponding *intersection number* as

$$c_{\text{int}}(\mathcal{I}) := \#\Omega_{\mathcal{I}}^0(\mathcal{E}) = \#\Omega_{\mathcal{I}}(\mathcal{E}),$$

where \mathcal{E} is an arbitrary s -tuple of flags in $\text{Good}(n, s)$. By Lemma 4.3.1, the right-hand side is finite and independent of the choice of \mathcal{E} in $\text{Good}(n, s)$. Moreover, $c_{\text{int}}(\mathcal{I}) > 0$ if and only if \mathcal{I} is intersecting.

In Section 6.2 we showed that if \mathcal{I} is intersecting then $c(\mathcal{I}) > 0$. Indeed, in this case the determinant function $\delta_{\mathcal{I}}$ on $\text{GL}(r)^s \times \text{GL}(n-r)^s$ is nonzero, so that for some suitable $\vec{h} \in \text{GL}(n-r)^s$ the function

$$(6.4.2) \quad \delta_{\mathcal{I}, \vec{h}}: \text{GL}(r)^s \rightarrow \mathbb{C}, \quad \delta_{\mathcal{I}, \vec{h}}(\vec{g}) := \delta_{\mathcal{I}}(\vec{g}, \vec{h})$$

is a nonzero vector in $\bigotimes_{k=1}^s L_{BW}(\lambda_{I_k}^*)$ that transforms as the character $\det_r^{(n-r)(s-1)}$ with respect to the diagonal action of $\text{GL}(r)$.

In the following we show that, as we vary \vec{h} , the functions $\delta_{\mathcal{I}, \vec{h}}$ span a vector space of dimension at least $c_{\text{int}}(\mathcal{I})$, which will imply that $c(\mathcal{I}) \geq c_{\text{int}}(\mathcal{I})$. More precisely, we shall construct elements $(\vec{g}_{\alpha}, \vec{h}_{\alpha}) \in \text{GL}(r)^s \times \text{GL}(n-r)^s$ for $\alpha \in [c_{\text{int}}(\mathcal{I})]$ such that $\delta_{\mathcal{I}}(\vec{g}_{\alpha}, \vec{h}_{\alpha}) \neq 0$ while $\delta_{\mathcal{I}}(\vec{g}_{\alpha}, \vec{h}_{\beta}) = 0$ if $\alpha \neq \beta$. The construction, due to Belkale [Bel2], depends on a choice of good flags \mathcal{E} and goes as follows.

Let \mathcal{E} be an s -tuple of good flags and consider the intersection

$$\Omega_{\mathcal{I}}^0(\mathcal{E}) = \{V_1, \dots, V_{c_{\text{int}}(\mathcal{I})}\}.$$

Let $\gamma_{\alpha} \in \text{GL}(n)$ such that $V_{\alpha} = \gamma_{\alpha} \cdot V_0$ for each $\alpha \in [c_{\text{int}}(\mathcal{I})]$, and consider the s -tuple of flags $\mathcal{E}_{\alpha} = (E_{\alpha,1}, \dots, E_{\alpha,s})$ defined by $E_{\alpha,k} = \gamma_{\alpha}^{-1} \cdot E_k$. Then $\bar{\omega}_{\mathcal{I}}^0([\gamma_{\alpha}, \mathcal{E}_{\alpha}]) = \mathcal{E}$. According to Lemma 4.3.1, \mathcal{E} is a regular value of $\bar{\omega}_{\mathcal{I}}^0$, since \mathcal{I} is intersecting. Since $\text{edim } \mathcal{I} = 0$, this implies that the differential of $\bar{\omega}_{\mathcal{I}}^0$ is bijective at $[\gamma_{\alpha}, \mathcal{E}_{\alpha}]$, and, by equivariance, so is its differential at $[1, \mathcal{E}_{\alpha}]$. By Remark 5.1.4, its determinant is precisely $\delta_{\mathcal{I}}(\vec{g}_{\alpha}, \vec{h}_{\alpha})$, where $\vec{g}_{\alpha} = (g_{\alpha,1}, \dots, g_{\alpha,s}) \in \text{GL}(r)$ and $\vec{h}_{\alpha} = (h_{\alpha,1}, \dots, h_{\alpha,s}) \in \text{GL}(n-r)$ are such that $g_{\alpha,k} \cdot F_0 = (E_{\alpha,k})^{V_0}$ and $h_{\alpha,k} \cdot G_0 = (E_{\alpha,k})^{Q_0}$ for all α and k . In particular, $\delta_{\mathcal{I}}(\vec{g}_{\alpha}, \vec{h}_{\alpha}) \neq 0$.

Using $\text{edim } \mathcal{I} = 0$, Eqs. (5.1.5) and (5.1.6) imply that

$$(6.4.3) \quad \delta_{\mathcal{I}}(\vec{g}, \vec{h}) \neq 0 \Leftrightarrow \dim H_{\mathcal{I}}(\vec{g} \cdot F_0, \vec{h} \cdot G_0) = 0.$$

Then we have the following lemma:

Lemma 6.4.4. *Let \mathcal{I} be intersecting, $\text{edim } \mathcal{I} = 0$, and $\mathcal{E} \in \text{Good}(n, s)$. As above, choose γ_{α} , \vec{g}_{α} and \vec{h}_{α} for $\alpha \in [c_{\text{int}}(\mathcal{I})]$. Define $\delta_{\mathcal{I}, \alpha}(\vec{g}) := \det \Delta_{\mathcal{I}, \vec{g}, \vec{h}_{\alpha}}$. Then $\delta_{\mathcal{I}, \alpha}(\vec{g}_{\alpha}) \neq 0$ for all α , while $\delta_{\mathcal{I}, \beta}(\vec{g}_{\alpha}) = 0$ for all $\alpha \neq \beta$.*

Proof. We only need to consider the case that $\alpha \neq \beta$. In view of (6.4.3), it suffices to show that $H_{\mathcal{I}}((\mathcal{E}_\alpha)^{V_0}, (\mathcal{E}_\beta)_{Q_0}) \neq \{0\}$. For this, we define the map

$$\phi_{\alpha,\beta}: V_0 \rightarrow \mathbb{C}^n / V_0 \cong Q_0, v \mapsto (\gamma_\alpha)^{-1} \gamma_\beta v + V_0,$$

which is nonzero since $\gamma_\alpha V_0 = V_\alpha \neq V_\beta = \gamma_\beta V_0$. Then $\phi_{\alpha,\beta}$ is a nonzero element in $H_{\mathcal{I}}((\mathcal{E}_\alpha)^{V_0}, (\mathcal{E}_\beta)_{Q_0})$, since

$$\phi_{\alpha,\beta}((E_{\beta,k})^{V_0}(a)) = \phi_{\alpha,\beta}(E_{\beta,k}(I_k(a))) = E_{\alpha,k}(I_k(a)) + V_0 = (E_{\alpha,k})_{Q_0}(I_k(a) - a)$$

for all $a \in [r]$ and $k \in [s]$, using that $\mathcal{I} = \text{Pos}(V_0, \mathcal{E}_\alpha)$. □

Lemma 6.4.4 shows that the functions $\delta_{\mathcal{I},1}, \dots, \delta_{\mathcal{I},c_{\text{int}}(\mathcal{I})}$ are linearly independent. If we identify them with $\text{GL}(r)$ -invariants as before, we obtain the following corollary:

Corollary 6.4.5. *Let $\text{edim } \mathcal{I} = 0$. Then, $c(\mathcal{I}) \geq c_{\text{int}}(\mathcal{I})$.*

In fact, it is a classical result that

$$(6.4.6) \quad c(\mathcal{I}) = c_{\text{int}}(\mathcal{I})$$

(see, e.g., [Full]). Thus Corollary 6.4.5 shows that we can produce a basis of the tensor product invariants from Belkale’s determinants $\delta_{\mathcal{I},\vec{h}}(\vec{g}) = \det \Delta_{\mathcal{I},\vec{g},\vec{h}}$. These invariants can be identified with the construction of Howe, Tan and Willenbring [HTW], as described in [VW].

7. Proof of Fulton’s conjecture

We now revisit the conjecture by Fulton which states that if $c(\vec{\lambda}) = 1$ for an s -tuple of highest weights then $c(N\vec{\lambda}) = 1$ for all $N \geq 1$. We note that its converse is also true and holds as a direct consequence of the saturation property and the bound $c(N\vec{\lambda}) \geq c(\vec{\lambda})$, which follows from the semigroup property of the Littlewood-Richardson coefficients. Fulton’s conjecture was first proved by Knutson, Tao and Woodward [KTW]. We closely follow Belkale’s geometric proof [Bel2, Bel3, Bel4], in its simplified form due to Sherman [She], which in turn was in part inspired by the technique of Schofield [Sch].

7.1. Nonzero invariants and intersections. Let $c(\vec{\lambda}) = 1$. Equivalently, $c(\vec{\lambda}^*) = 1$ and so there exists a nonzero $\text{GL}(r)$ -invariant holomorphic function f in $L_{BW}(\lambda_1^*) \otimes \dots \otimes L_{BW}(\lambda_s^*)$, which is unique up to rescaling.

Suppose for a moment that there exists a nowhere vanishing function g in $L_{BW}(\lambda_1^*) \otimes \dots \otimes L_{BW}(\lambda_s^*)$ (not necessarily $GL(r)$ -invariant). In this case, if g' is any other holomorphic function in $L_{BW}(\lambda_1^*) \otimes \dots \otimes L_{BW}(\lambda_s^*)$, then g'/g is right $B(r)^s$ -invariant and therefore descends to a holomorphic function on $\text{Flag}(r)^s$. But this is a compact space, hence any such function is constant. It then follows that each $L_{BW}(\lambda_k^*)$ is one-dimensional and hence that the λ_k are just characters, i.e., $\lambda_k = m_k \mathbb{1}_r$ and $L(\lambda_k) = \det_r^{m_k}$ for some $m_k \in \mathbb{Z}$. In this case, Fulton's conjecture is certainly true.

We now consider the nontrivial case when f has zeros. For any function in $L_{BW}(\lambda_1^*) \otimes \dots \otimes L_{BW}(\lambda_s^*)$, the zero set is right $B(r)^s$ -stable. Accordingly, we shall write $f(\mathcal{F}) = 0$ for the condition that $f(\vec{g}) = 0$, where $\mathcal{F} = \vec{g} \cdot F_0$, and consider

$$Z_f := \{ \mathcal{F} \in \text{Flag}(r)^s : f(\mathcal{F}) = 0 \}.$$

Without loss of generality, we may assume that there exists an s -tuple \mathcal{I} with $\text{edim } \mathcal{I} = 0$ that is related to $\vec{\lambda}$ as in Lemma 4.3.9, i.e.,

$$(7.1.1) \quad \lambda_k = \lambda_{I_k} + (n-r)\mathbb{1}_r \text{ for } k \in [s-1], \quad \lambda_s = \lambda_{I_s}$$

(otherwise we may add/remove suitable multiples of $\mathbb{1}_r$, as in the proof of Proposition 6.3.2). Now recall from (6.4.2) that the functions $\delta_{\mathcal{I}, \vec{h}} = \delta_{\mathcal{I}}(-, \vec{h})$ are in $\bigotimes_{k=1}^s L_{BW}(\lambda_{I_k}^*)$ and transform as the character $\det_r^{(n-r)(s-1)}$ with respect to the diagonal action of $GL(r)$. It follows that each $\tilde{\delta}_{\mathcal{I}, \vec{h}}(\vec{g}) := \det_r^{-(n-r)}(g_1) \dots \det_r^{-(n-r)}(g_{s-1}) \delta_{\mathcal{I}, \vec{h}}(\vec{g})$ must be proportional to f . Hence,

$$(7.1.2) \quad \delta_{\mathcal{I}}(\vec{g}, \vec{h}) = \det_r^{(n-r)}(g_1) \dots \det_r^{(n-r)}(g_{s-1}) f(\vec{g}) \hat{f}(\vec{h}),$$

for some function $\hat{f}: GL(n-r)^s \rightarrow \mathbb{C}$, which is nonzero due to (6.4.6). In view of (6.4.3), we obtain the following lemma:

Lemma 7.1.3. *Let f, \mathcal{I} as above. If $\mathcal{F} \in Z_f$ then $H_{\mathcal{I}}(\mathcal{F}, \mathcal{G}) \neq \{0\}$ for all $\mathcal{G} \in \text{Flag}(Q_0)^s$. Conversely, if $\hat{f}(\mathcal{G}) \neq 0$ then $H_{\mathcal{I}}(\mathcal{F}, \mathcal{G}) \neq \{0\}$ implies that $\mathcal{F} \in Z_f$.*

For sake of finding a contradiction, let us assume that $c(N\vec{\lambda}) > 1$ for some N . Then there exists an invariant $f' \in L_{BW}(N\lambda_1^*) \otimes \dots \otimes L_{BW}(N\lambda_s^*)$ that is linearly independent from f^N .

Lemma 7.1.4. *Let \mathcal{L} be a holomorphic line bundle over a smooth irreducible variety. Then two linearly independent holomorphic sections f_1, f_2 are automatically algebraically independent.*

Proof. Let us suppose that f_1 and f_2 satisfy a nontrivial relation $\sum_{i,j} c_{i,j} f_1^i f_2^j = 0$. Each $f_1^i f_2^j$ is a section of the line bundle $\mathcal{L}^{\otimes(i+j)}$. The relation holds degree by degree, and so we may assume that $i + j$ is the same for each nonzero $c_{i,j}$. But any homogeneous polynomial in two variables is a product of linear factors. Thus we have $\prod_i (a_i f_1 + b_i f_2) = 0$ for some $a_i, b_i \in \mathbb{C}$, and one of the factors has to vanish identically. This shows that f_1 and f_2 are linearly dependent, in contradiction to our assumption. \square

Lemma 7.1.4 implies that f^N and f' , and therefore f and f' are algebraically independent. As a consequence, there exists a nonempty Zariski-open subset of $\mathcal{F} \in Z_f$ such that $f'(\mathcal{F}) \neq 0$.

Our strategy in the below will be as follows. As before, we consider the kernel position \mathcal{J} of a generic map $0 \neq \phi \in H_{\mathcal{I}}(\mathcal{F}, \mathcal{G})$, with now \mathcal{F} varying in Z_f . Although \mathcal{J} is not necessarily intersecting, the condition $f'(\mathcal{F}) \neq 0$ will be sufficient to show that the tuple $\mathcal{I}^{\mathcal{J}}$ is intersecting. In Section 7.2 we will then prove Sherman's refined version of his recurrence relation (5.3.3), which will allow us to show that $H_{\mathcal{I}}(\mathcal{F}, \mathcal{G}) = \{0\}$ for generic $\mathcal{F} \in Z_f$. In view of Lemma 7.1.3, this will give a contradiction.

We first prove a general lemma relating semistable vectors and moment maps. Let M be a complex vector space equipped with a $GL(r)$ -representation and $U(r)$ -invariant Hermitian inner product $\langle \cdot, \cdot \rangle$, complex linear in the second argument, and denote by $\rho_M: \mathfrak{gl}(r) \rightarrow \mathfrak{gl}(M)$ the Lie algebra representation. We define the corresponding *moment map* $\Phi_M: \mathbb{P}(M) \rightarrow i\mathfrak{u}(r)$ by

$$\mathrm{tr}(\Phi_M([m])A) = \frac{\langle m, \rho_M(A)m \rangle}{\|m\|^2}$$

for all $A \in \mathfrak{gl}(r)$; cf. Eq. (2.2).

Lemma 7.1.5. *Let $A \in i\mathfrak{u}(r)$ and $0 \neq m \in M$. If $\exp(At) \cdot m \not\rightarrow 0$ as $t \rightarrow -\infty$ then*

$$\lim_{t \rightarrow -\infty} \mathrm{tr}(\Phi_M([\exp(At) \cdot m])A) \leq 0.$$

Proof. Write $m = \sum_{i=1}^k m_i$ where the m_i are nonzero eigenvectors of $\rho_M(A)$, with eigenvalues $\theta_1 < \dots < \theta_k$. Then,

$$\exp(At) \cdot m = \sum_{i=1}^k e^{\theta_i t} m_i \not\rightarrow 0$$

as $t \rightarrow -\infty$ if and only if $\theta_1 \leq 0$. In this case,

$$\lim_{t \rightarrow -\infty} \mathrm{tr}(\Phi_M([\exp(At) \cdot m])A) = \lim_{t \rightarrow -\infty} \frac{\sum_i \theta_i e^{2\theta_i t} \|m_i\|^2}{\sum_i e^{2\theta_i t} \|m_i\|^2} = \theta_1 \leq 0. \quad \square$$

We now relate the position of subspaces to components of the moment map:

Lemma 7.1.6. *Let $\lambda \in \Lambda_+(r)$, $F = g \cdot F_0$ a flag on V_0 , S a nonzero subspace of \mathbb{C}^r , and P_S the orthogonal projector. Then,*

$$\lim_{t \rightarrow -\infty} \operatorname{tr}(\Phi_{L(\lambda)}([\exp(P_S t)g \cdot v_\lambda])P_S) = \langle T_J, \lambda \rangle,$$

where $J = \operatorname{Pos}(S, F)$.

Proof. Let $d = \#J$. We may assume that $S = S_0$ is generated by the first d vectors $e(1), \dots, e(d)$ of the standard basis of V_0 , and also that $g = u$ is unitary. Thus P_{S_0} is the diagonal matrix with d ones and $r - d$ zeros, and we need to show that

$$\lim_{t \rightarrow -\infty} \operatorname{tr}(\Phi_{L(\lambda)}([\exp(P_{S_0} t)u \cdot v_\lambda])P_{S_0}) = \langle T_J, \lambda \rangle.$$

Let R_0 denote the orthogonal complement of S_0 in V_0 . The action of $U(S_0) \times U(R_0)$ commutes with P_{S_0} and hence we can assume that F^{S_0} is the standard flag on S_0 , while F_{V_0/S_0} has the adapted basis $e(J^c(b)) + S_0$ for $b \in [r - d]$. Thus we see that $\lim_{t \rightarrow -\infty} \exp(P_{S_0} t)F = w_J F_0$. It follows that $\lim_{t \rightarrow -\infty} [\exp(P_{S_0} t)u \cdot v_\lambda] = [w_J \cdot v_\lambda]$ and hence, using (2.2), that

$$\lim_{t \rightarrow -\infty} \operatorname{tr}(\Phi_{L(\lambda)}([\exp(P_{S_0} t)u \cdot v_\lambda])P_{S_0}) = \frac{\langle v_\lambda, \rho_\lambda(w_J^{-1} P_{S_0} w_J) v_\lambda \rangle}{\|v_\lambda\|^2} = \langle T_J, \lambda \rangle. \quad \square$$

We now use the preceding lemma to obtain from any nonzero invariant an s -tuple of flags with nonnegative slope:

Lemma 7.1.7. *Let $p \in (L(N\lambda_1) \otimes \dots \otimes L(N\lambda_s))^*$ a $\operatorname{GL}(r)$ -invariant homogeneous polynomial such that $p(g_1 \cdot v_{N\lambda_1} \otimes \dots \otimes g_s \cdot v_{N\lambda_s}) \neq 0$, and define $\mathcal{F} = (g_1 F_0, \dots, g_s F_0)$. Then $\sum_{k=1}^s (T_{J_k}, \lambda_k) \leq 0$ for all $\mathcal{J} = \operatorname{Pos}(S, \mathcal{F})$, where S is an arbitrary nonzero subspace of \mathbb{C}^r .*

Proof. Consider the representation $M = L(N\lambda_1) \otimes \dots \otimes L(N\lambda_s)$ with its moment map Φ_M , and $m := g_1 \cdot v_{N\lambda_1} \otimes \dots \otimes g_s \cdot v_{N\lambda_s}$. Let P_S denote the orthogonal projector onto the subspace S . As p is $\operatorname{GL}(r)$ -invariant, $p(\exp(P_S t) \cdot m) = p(m) \neq 0$, which implies that $\exp(P_S t) \cdot m \not\rightarrow 0$ as $t \rightarrow -\infty$. Thus Lemma 7.1.5 implies that

$$\lim_{t \rightarrow -\infty} \operatorname{tr}(\Phi_M([\exp(P_S t) \cdot m])P_S) \leq 0.$$

On the other hand, Lemma 7.1.6 shows that the left-hand side of this inequality is equal to

$$\sum_{k=1}^s \lim_{t \rightarrow -\infty} \operatorname{tr}(\Phi_{L(N\lambda_k)}([\exp(P_S t)g_k \cdot v_{N\lambda_k}]P_S) = \sum_{k=1}^s \lim_{t \rightarrow -\infty} (T_{J_k}, \lambda_k). \quad \square$$

Corollary 7.1.8. *Let $\vec{\lambda}$ and \mathcal{I} as in (7.1.1), $f' \in (L_{BW}(N\lambda_1^*) \otimes \cdots \otimes L_{BW}(N\lambda_s^*))^{\text{GL}(r)}$. Let $\mathcal{F} \in \text{Flag}(V_0)^s$, $\{0\} \neq S \subseteq \mathbb{C}^r$, and $\mathcal{J} = \text{Pos}(S, \mathcal{F})$. If $f'(\mathcal{F}) \neq 0$ then $\mathcal{I}^{\mathcal{J}}$ is intersecting.*

Proof. Write $\mathcal{F} = (g_1 \cdot F_0, \dots, g_s \cdot F_0)$ for suitable $g_1, \dots, g_s \in \text{GL}(r)$. Then, using (6.1.2), there exists a $\text{GL}(r)$ -invariant homogeneous polynomial $p \in (L(N\lambda_1) \otimes \cdots \otimes L(N\lambda_s))^*$ such that $p(g_1 \cdot v_{N\lambda_1} \otimes \cdots \otimes g_s \cdot v_{N\lambda_s}) = f'(g_1, \dots, g_s) \neq 0$. Thus the assumptions of Lemma 7.1.7 are satisfied.

We now show that $\mathcal{I}^{\mathcal{J}}$ is intersecting. For this, we use Theorem 5.3.4 and verify the Horn inequalities. Thus let $0 < m \leq d = \dim S$ and $\mathcal{K} \in \text{Horn}(m, d, s) = \text{Intersecting}(m, d, s)$: Since \mathcal{K} is intersecting, there exists some subspace $S' \in \Omega_{\mathcal{K}}(\mathcal{F}^S)$. Hence $S' \in \Omega_{\mathcal{J}\mathcal{K}}(\mathcal{F})$ by the chain rule (3.2.9). According to Lemma 3.1.5, $\mathcal{J}' = \text{Pos}(S', \mathcal{F})$ is such that $J'_k(a) \leq J_k K_k(a)$ for all $k \in [s]$ and $a \in [m]$. Thus we obtain the first inequality in

$$\begin{aligned} \text{edim } \mathcal{I}^{\mathcal{J}} \mathcal{K} - \text{edim } \mathcal{K} &= \text{edim } \mathcal{I}(\mathcal{J}\mathcal{K}) - \text{edim } \mathcal{J}\mathcal{K} \\ &= - \sum_{k=1}^s (T_{J_k K_k}, \lambda_k) \geq - \sum_{k=1}^s (T_{J'_k}, \lambda_k) \geq 0; \end{aligned}$$

the first equality is (4.2.11), the second is Lemma 4.3.9, and the last inequality is Lemma 7.1.7, applied to S' . This concludes the proof. \square

7.2. Sherman’s refined lemma. We now study the behavior of $\dim H_{\mathcal{I}}(\mathcal{F}, \mathcal{G})$ in more detail. We proceed as in Section 5, but for a *fixed* s -tuple of flags $\mathcal{F} \in \text{Flag}(V_0)^s$. Specifically, we consider the following refinement of the true dimension (5.1.8) for fixed \mathcal{F} :

$$\text{tdim}_{\mathcal{F}} \mathcal{I} := \min_{\mathcal{G}} \dim H_{\mathcal{I}}(\mathcal{F}, \mathcal{G})$$

Thus we study the variety

$$P_{\mathcal{F}}(\mathcal{I}) := \{(\mathcal{G}, \phi) \in \text{Flag}(Q_0)^s \times \text{Hom}(V_0, Q_0) : \phi \in H_{\mathcal{I}}(\mathcal{F}, \mathcal{G})\}.$$

Restricting to those \mathcal{G} such that $\dim H_{\mathcal{I}}(\mathcal{F}, \mathcal{G}) = \text{tdim}_{\mathcal{F}} \mathcal{I}$, we obtain open sets $B_{\mathcal{F},t}(\mathcal{I}) \subseteq \text{Flag}(Q_0)^s$ and $P_{\mathcal{F},t}(\mathcal{I}) \subseteq P_{\mathcal{F}}(\mathcal{I})$. Let $\text{kdim}_{\mathcal{F}}(\mathcal{I})$ denote the minimal (and hence generic) dimension of $\ker \phi$ for $(\mathcal{G}, \phi) \in P_{\mathcal{F},t}(\mathcal{I})$. The following lemma is proved just like Corollary 5.2.6:

Lemma 7.2.1. *If $\text{kdim}_{\mathcal{F}} \mathcal{I} = 0$ then $\text{tdim}_{\mathcal{F}} \mathcal{I} = \text{edim } \mathcal{I}$.*

Let us now assume that $\text{kdim}_{\mathcal{F}} \mathcal{I} > 0$. Let $\text{kPos}_{\mathcal{F}}(\mathcal{I})$ denote the kernel position, defined as in Definition 5.2.8 but for fixed \mathcal{F} . We thus obtain an irreducible

variety $P_{\mathcal{F},\text{kpt}}(\mathcal{I})$ over a Zariski-open subset $B_{\mathcal{F},\text{kpt}}(\mathcal{I})$ of $B_{\mathcal{F},t}(\mathcal{I})$. To compute its dimension, we again define $P_{\mathcal{F},\text{kp}}(\mathcal{I}) \subseteq P_{\mathcal{F}}(\mathcal{I})$, where we fix the kernel dimension and position, but *not* the dimension of $H_{\mathcal{I}}(\mathcal{F}, \mathcal{G})$. In contrast to Lemma 5.2.11, the variety $P_{\mathcal{F},\text{kp}}(\mathcal{I})$ is in general neither smooth nor irreducible. However, we can describe it similarly as before: We first constrain $S = \ker \phi$ to be in $\Omega_{\mathcal{J}}^0(\mathcal{F})$ (which may not be irreducible), then ϕ is determined by $\bar{\phi} \in \text{Hom}^{\times}(V_0/S, Q_0)$ and \mathcal{G} by $G_k \in \text{Flag}_{I_k/J_k}^0((F_k)_{V_0/S}, \bar{\phi})$. Thus we obtain for each irreducible component $C \subseteq \Omega_{\mathcal{J}}^0(\mathcal{F})$ a corresponding irreducible component $P_{\mathcal{F},\text{kp},C}(\mathcal{I})$. In particular, there exists some component $C_{\mathcal{F}}$ such that $P_{\mathcal{F},\text{kp},C_{\mathcal{F}}}(\mathcal{I})$ is the closure of $P_{\mathcal{F},\text{kpt}}(\mathcal{I})$ in $P_{\mathcal{F},\text{kp}}(\mathcal{I})$, namely the irreducible component containing the elements $S = \ker \phi$ for (ϕ, \mathcal{G}) varying in the irreducible variety $P_{\mathcal{F},\text{kpt}}(\mathcal{I})$. As a consequence, $\dim P_{\mathcal{F},\text{kpt}}(\mathcal{I}) = \dim P_{\mathcal{F},\text{kp},C_{\mathcal{F}}}(\mathcal{I})$, and so we obtain, using completely analogous dimension computations, the following refinement of (5.3.1):

$$(7.2.2) \quad \text{tdim}_{\mathcal{F}} \mathcal{I} - \text{edim} \mathcal{I} = \dim C_{\mathcal{F}} - \text{edim} \mathcal{I}^{\mathcal{J}}$$

Indeed, when we apply (7.2.2) to generic $\mathcal{F} \in \text{Flag}(V_0)^s$ then \mathcal{J} is intersecting and $\dim C_{\mathcal{F}} = \text{edim} \mathcal{J}$, so we recover (5.3.1). We now instead apply the above to generic \mathcal{F} in a component of the zero set Z_f of the unique nonzero invariant f . Thus we obtain the following variant of the key recursion relation (5.3.3):

Lemma 7.2.3 (Sherman). *Let f, \mathcal{I} as above in Section 7.1, and $Z \subseteq Z_f$ an irreducible component such that $\text{kdim}_{\mathcal{F}} \mathcal{I} \neq 0$ for all $\mathcal{F} \in Z$. Then there exists \mathcal{J} and a nonempty Zariski-open subset of $\mathcal{F} \in Z$ such that $\text{kPos}_{\mathcal{F}} \mathcal{I} = \mathcal{J}$ and*

$$\text{tdim}_{\mathcal{F}} \mathcal{I} - \text{edim} \mathcal{I} \leq \text{tdim} \mathcal{I}^{\mathcal{J}} - \text{edim} \mathcal{I}^{\mathcal{J}}.$$

Proof. We choose d and \mathcal{J} as the kernel dimension and position for generic $\mathcal{F} \in Z$. We note that $d < r$, since $d = r$ would imply that $H_{\mathcal{I}}(\mathcal{F}, \mathcal{G}) = \{0\}$, in contradiction to Lemma 7.1.3. Let $U \subseteq Z$ denote the Zariski-open subset such that $\text{kPos}_{\mathcal{F}} \mathcal{I} = \mathcal{J}$ for all $\mathcal{F} \in U$. We proceed as in Lemma 5.3.2. Let

$$\begin{aligned} \mathcal{X} &:= \{(\mathcal{F}, \mathcal{G}, \phi) : \mathcal{F} \in U, (\mathcal{G}, \phi) \in P_{\mathcal{F},\text{kp},C_{\mathcal{F}}}(\mathcal{I})\}, \\ \mathcal{Y} &:= \{(S, \tilde{\mathcal{F}}, \mathcal{G}) : S \in \text{Gr}(d, V_0), \tilde{\mathcal{F}} \in \text{Flag}(S)^s, \mathcal{G} \in \text{Flag}(Q_0)^s\}. \end{aligned}$$

Both \mathcal{X} and \mathcal{Y} are irreducible varieties and we have a morphism

$$\pi : \mathcal{X} \rightarrow \mathcal{Y}, (\mathcal{F}, \mathcal{G}, \phi) \mapsto (\ker \phi, \mathcal{F}^{\ker \phi}, \mathcal{G}).$$

As before, we argue that π is dominant. By construction, the image of \mathcal{X} by the map $(\mathcal{F}, \mathcal{G}, \phi) \mapsto \mathcal{G}$ contains a Zariski-open subset U' of $\text{Flag}(Q_0)^s$. We may also assume that $\hat{f}(\mathcal{G}) \neq 0$ for all $\mathcal{G} \in U'$, where \hat{f} is the map from (7.1.2). We now show that the image of π contains all elements $(S, \tilde{\mathcal{F}}, \mathcal{G})$

with $S \in \text{Gr}(d, V_0)$, $\tilde{\mathcal{F}} \in \text{Flag}(S)^s$, and $\mathcal{G} \in U'$. For this, let $(\mathcal{F}_0, \mathcal{G}, \phi_0) \in \mathcal{X}$ be the preimage of some arbitrary $\mathcal{G} \in U'$. Let $S_0 := \ker \phi_0$ and choose some $g \in \text{GL}(V_0)$ such that $g \cdot S_0 = S$. Using the corresponding diagonal action, define $\mathcal{F} := g \cdot \mathcal{F}_0$ and $\phi := g \cdot \phi_0$. Then $(\mathcal{F}, \mathcal{G}, \phi) \in \mathcal{X}$, since Z is stable under the diagonal action of $\text{GL}(V_0)$, and $\ker \phi = S$. Now consider the group $G \subseteq \text{GL}(V_0)^s$ consisting of all elements $\vec{h} \in \text{GL}(V_0)^s$ such that $h_k S \subseteq S$ and h_k acts trivially on V_0/S for all $k \in [s]$. Note that G is an irreducible algebraic group. By construction, $\phi \in H_{\mathcal{I}}(\vec{h} \cdot \mathcal{F}, \mathcal{G})$, while $d < r$ implies that $\phi \neq 0$. This means that $H_{\mathcal{I}}(\vec{h} \cdot \mathcal{F}, \mathcal{G}) \neq 0$, and so we obtain from Lemma 7.1.3 that $\vec{h} \cdot \mathcal{F} \in Z_f$. It follows that, in fact, $\vec{h} \cdot \mathcal{F} \in Z$, as it is obtained by the action of the irreducible algebraic group G on $\mathcal{F} \in Z$, and so stays in the same irreducible component. For given $\tilde{\mathcal{F}} \in \text{Flag}(S)^s$, we now choose $\vec{h} \in G$ such that $h_k \cdot F_k^S = \tilde{F}_k$ for $k \in [s]$. Then $(\vec{h} \cdot \mathcal{F}, \mathcal{G}, \phi) \in \mathcal{X}$ is a preimage of $(S, \tilde{\mathcal{F}}, \mathcal{G})$, and we conclude that π is dominant.

As before, the dominance implies that we can find a nonempty Zariski-open set of $(\mathcal{F}, \mathcal{G}, \phi) \in \mathcal{X}$ such that $\dim H_{\mathcal{I}^{\mathcal{J}}}(\mathcal{F}^{\ker \phi}, \mathcal{G}) = \text{tdim } \mathcal{I}^{\mathcal{J}}$. We may assume in addition that $\ker \phi \in C_{\mathcal{F}}$ is a smooth point. For any \mathcal{F} in this set, $\bar{\phi}$ injects $H_{\mathcal{J}}(\mathcal{F}^{\ker \phi}, \mathcal{F}_{V_0/\ker \phi})$ into $H_{\mathcal{I}^{\mathcal{J}}}(\mathcal{F}^{\ker \phi}, \mathcal{G})$ (Lemma 3.2.15). Thus,

$$\dim C_{\mathcal{F}} = \dim T_{\ker \phi} C_{\mathcal{F}} \leq H_{\mathcal{J}}(\mathcal{F}^{\ker \phi}, \mathcal{F}_{V_0/\ker \phi}) \leq \dim H_{\mathcal{I}^{\mathcal{J}}}(\mathcal{F}^{\ker \phi}, \mathcal{G}) = \text{tdim } \mathcal{I}^{\mathcal{J}},$$

where in the first inequality we have used that the intersection $\Omega_{\mathcal{J}}^0(\mathcal{F})$ is not necessarily transversal at S and so the tangent space of the intersection is in general only a subspace of the intersection of the tangent spaces (3.2.5). In view of (7.2.2), we obtain the desired inequality. \square

Theorem 7.2.4 (Belkale). *Let $c(\vec{\lambda}) = 1$. Then $c(N\vec{\lambda}) = 1$ for all $N > 1$.*

Proof. Let f, \mathcal{I} as in Section 7.1 and recall that $\text{edim } \mathcal{I} = 0$. Assume for sake of finding a contradiction that $c(N\vec{\lambda}) \neq 1$ for some $N > 0$. Then there exists another invariant f' , as in Section 7.1, such that $f'(\mathcal{F}) \neq 0$ for a nonempty Zariski-open subset of some irreducible component $Z \subseteq Z_f$. If $\text{kdim}_{\mathcal{F}} \mathcal{I} = 0$ for some $\mathcal{F} \in Z$ then $\text{tdim}_{\mathcal{F}} \mathcal{I} = 0$ by Lemma 7.2.1. Otherwise, we may apply Lemma 7.2.3 to the component Z . We find that there exists some \mathcal{J} and another Zariski-open subset of \mathcal{F} in Z such that $\text{kPos}_{\mathcal{F}} \mathcal{I} = \mathcal{J}$ and

$$(7.2.5) \quad \text{tdim}_{\mathcal{F}} \mathcal{I} - \text{edim } \mathcal{I} \leq \text{tdim } \mathcal{I}^{\mathcal{J}} - \text{edim } \mathcal{I}^{\mathcal{J}}.$$

As a consequence, there exists some $\mathcal{F} \in Z_f$ for which all three of the properties $f'(\mathcal{F}) \neq 0$, $\text{kPos}_{\mathcal{F}} \mathcal{I} = \mathcal{J}$ and (7.2.5) hold true. By Corollary 7.1.8, the first two properties imply that $\mathcal{I}^{\mathcal{J}} = 0$, and hence the right-hand side of (7.2.5) is equal to zero by Corollary 5.1.9. This again implies that $\text{tdim}_{\mathcal{F}} \mathcal{I} = 0$.

It follows that in either case there exist some \mathcal{G} such that $H_{\mathcal{I}}(\mathcal{F}, \mathcal{G}) = \{0\}$. According to Lemma 7.1.3, this can only be if $\mathcal{F} \notin Z_f$. But $\mathcal{F} \in Z_f$. This is the desired contradiction. \square

Remark. It likewise holds that $c(\vec{\lambda}) = 2$ implies that $c(N\vec{\lambda}) = N + 1$ [Ike, She]. However, in general it is *not* true that $c(\vec{\lambda}) = c$ implies $c(N\vec{\lambda}) = O(N^{c-1})$. Belkale has found a counterexample for $c = 6$.

A. Horn triples in low dimensions

In this appendix, we list all Horn triples $\mathcal{J} \in \text{Horn}(d, r, 3)$ for $d < r \leq 4$, as defined in Definition 2.18, as well as the expected dimensions $\text{edim } \mathcal{J}$. The triples with $\text{edim } \mathcal{J} = 0$ are highlighted in bold.

Example A.1 ($d = 1$). As discussed in Example 4.3.12, only the dimension condition $\text{edim } \mathcal{J} \geq 0$ is necessary. The following are the triples in $\text{Horn}(1, r, 3)$ (up to permutations):

r	J_1	J_2	J_3	$\text{edim } \mathcal{J}$
2	{1}	{2}	{2}	0
	{2}	{2}	{2}	1
3	{1}	{3}	{3}	0
	{2}	{2}	{3}	0
	{2}	{3}	{3}	1
	{3}	{3}	{3}	2
4	{1}	{4}	{4}	0
	{2}	{3}	{4}	0
	{2}	{4}	{4}	1
	{3}	{3}	{3}	0
	{3}	{3}	{4}	1
	{3}	{4}	{4}	2
	{4}	{4}	{4}	3

Example A.2 ($d = 2$). The dimension condition $\text{edim } \mathcal{J} \geq 0$ reads

$$(J_1(1) + J_1(2)) + (J_2(1) + J_2(2)) + (J_3(1) + J_3(2)) \geq 4r + 1.$$

In addition, we have to satisfy three Horn inequalities, corresponding to $\mathcal{K} = (\{1\}, \{2\}, \{2\})$ and its permutations, which are the only elements in

Horn(1, 2, 3) with $\dim \mathcal{K} = 0$. The resulting Horn inequalities, $\text{edim } \mathcal{JK} \geq 0$, are

$$J_1(1) + J_2(2) + J_3(2) \geq 2r + 1,$$

$$J_1(2) + J_2(1) + J_3(2) \geq 2r + 1,$$

$$J_1(2) + J_2(2) + J_3(1) \geq 2r + 1.$$

Thus we obtain the following triples in Horn(2, r , 3) (up to permutations):

r	J_1	J_2	J_3	$\text{edim } \mathcal{J}$
3	{1, 2}	{2, 3}	{2, 3}	0
	{1, 3}	{1, 3}	{2, 3}	0
	{1, 3}	{2, 3}	{2, 3}	1
	{2, 3}	{2, 3}	{2, 3}	2
4	{1, 2}	{3, 4}	{3, 4}	0
	{1, 3}	{2, 4}	{3, 4}	0
	{1, 3}	{3, 4}	{3, 4}	1
	{1, 4}	{1, 4}	{3, 4}	0
	{1, 4}	{2, 4}	{2, 4}	0
	{1, 4}	{2, 4}	{3, 4}	1
	{1, 4}	{3, 4}	{3, 4}	2
	{2, 3}	{2, 3}	{3, 4}	0
	{2, 3}	{2, 4}	{2, 4}	0
	{2, 3}	{2, 4}	{3, 4}	1
	{2, 3}	{3, 4}	{3, 4}	2
	{2, 4}	{2, 4}	{2, 4}	1
	{2, 4}	{2, 4}	{3, 4}	2
	{2, 4}	{3, 4}	{3, 4}	3
{3, 4}	{3, 4}	{3, 4}	4	

Example A.3. We find the following triples in Horn(3, 4, 3) (up to permutations):

r	J_1	J_2	J_3	$\text{edim } \mathcal{J}$
4	{1, 2, 3}	{2, 3, 4}	{2, 3, 4}	0
	{1, 2, 4}	{1, 3, 4}	{2, 3, 4}	0
	{1, 2, 4}	{2, 3, 4}	{2, 3, 4}	1
	{1, 3, 4}	{1, 3, 4}	{1, 3, 4}	0
	{1, 3, 4}	{1, 3, 4}	{2, 3, 4}	1
	{1, 3, 4}	{2, 3, 4}	{2, 3, 4}	2
	{2, 3, 4}	{2, 3, 4}	{2, 3, 4}	3

Remark. It is not an accident that both $\text{Horn}(1, 4, 3)$ and $\text{Horn}(3, 4, 3)$ have the same number of elements. In fact, we can identify $\text{Intersecting}(d, r, s) \cong \text{Intersecting}(r - d, r, s)$ via $J_k \mapsto \{r + 1 - a : a \in J_k^c\}$. This can be seen by using the canonical isomorphism $\text{Gr}(d, \mathbb{C}^r) \cong \text{Gr}(r - d, (\mathbb{C}^r)^*)$. However, the corresponding Horn inequalities are distinct (see below).

B. Kirwan cones in low dimensions

In this appendix, we list necessary and sufficient conditions on highest weights $\lambda, \mu, \nu \in \Lambda_+(r)$ such that $(L(\lambda) \otimes L(\mu) \otimes L(\nu))^{U(r)} \neq \{0\}$, up to $r = 4$. That is, these conditions describe the Kirwan cones as in Corollary 6.3.3. We use the abbreviation $\text{Horn}_0(d, r, s)$ for the set of Horn triples in $\mathcal{J} \in \text{Horn}(d, r, s)$ such that $\text{edim } \mathcal{J} = 0$ (highlighted bold in Appendix A).

Example B.1 ($r = 1$). Clearly, the only condition is $\lambda(1) + \mu(1) + \nu(1) = 0$.

Example B.2 ($r = 2$). We always have the Weyl chamber inequalities $\lambda(1) \geq \lambda(2)$, $\mu(1) \geq \mu(2)$, and $\nu(1) \geq \nu(2)$, and the equation

$$(\lambda(1) + \lambda(2)) + (\mu(1) + \mu(2)) + (\nu(1) + \nu(2)) = 0.$$

Using Example A.1, we obtain three Horn inequalities, namely

$$\lambda(1) + \mu(2) + \nu(2) \leq 0,$$

corresponding to the triple $(\{1\}, \{2\}, \{2\}) \in \text{Horn}_0(d, r, s)$, and its permutations. These are the well-known conditions for the existence of nonzero invariants in a triple tensor product of irreducible $U(2)$ -representations. We remark that the Weyl chamber inequalities are redundant.

Example B.3 ($r = 3$). In addition to the Weyl chamber inequalities and $|\lambda| + |\mu| + |\nu| = 0$, we obtain the following two inequalities from $\text{Horn}_0(1, 3, 3)$ and Example A.1,

$$\lambda(1) + \mu(3) + \nu(3) \leq 0,$$

$$\lambda(2) + \mu(2) + \nu(3) \leq 0,$$

and the following from $\text{Horn}_0(2, 3, 3)$ and Example A.2,

$$(\lambda(1) + \lambda(2)) + (\mu(2) + \mu(3)) + (\nu(2) + \nu(3)) \leq 0,$$

$$(\lambda(1) + \lambda(3)) + (\mu(1) + \mu(3)) + (\nu(2) + \nu(3)) \leq 0,$$

as well as their permutations.

Example B.4 ($r = 4$). Again we have the Weyl chamber inequalities and $|\lambda| + |\mu| + |\nu| = 0$. We have the following two inequalities and their permutations from $\text{Horn}_0(1, 4, 3)$ and Example A.1,

$$\lambda(1) + \mu(4) + \nu(4) \leq 0,$$

$$\lambda(2) + \mu(3) + \nu(4) \leq 0,$$

$$\lambda(3) + \mu(3) + \nu(3) \leq 0,$$

the following six and their permutations from $\text{Horn}_0(2, 4, 3)$ and Example A.2,

$$(\lambda(1) + \lambda(2)) + (\mu(3) + \mu(4)) + (\nu(3) + \nu(4)) \leq 0,$$

$$(\lambda(1) + \lambda(3)) + (\mu(2) + \mu(4)) + (\nu(3) + \nu(4)) \leq 0,$$

$$(\lambda(1) + \lambda(4)) + (\mu(1) + \mu(4)) + (\nu(3) + \nu(4)) \leq 0,$$

$$(\lambda(1) + \lambda(4)) + (\mu(2) + \mu(4)) + (\nu(2) + \nu(4)) \leq 0,$$

$$(\lambda(2) + \lambda(3)) + (\mu(2) + \mu(3)) + (\nu(3) + \nu(4)) \leq 0,$$

$$(\lambda(2) + \lambda(3)) + (\mu(2) + \mu(4)) + (\nu(2) + \nu(4)) \leq 0,$$

and the following three and their permutations from $\text{Horn}_0(3, 4, 3)$ and Example A.3,

$$(\lambda(1) + \lambda(2) + \lambda(3)) + (\mu(2) + \mu(3) + \mu(4)) + (\nu(2) + \nu(3) + \nu(4)) \leq 0,$$

$$(\lambda(1) + \lambda(2) + \lambda(4)) + (\mu(1) + \mu(3) + \mu(4)) + (\nu(2) + \nu(3) + \nu(4)) \leq 0,$$

$$(\lambda(1) + \lambda(3) + \lambda(4)) + (\mu(1) + \mu(3) + \mu(4)) + (\nu(1) + \nu(3) + \nu(4)) \leq 0.$$

Remark. In low dimensions, all Horn triples with $\text{edim } \mathcal{J} = 0$ are such that the intersection is one point, i.e., $c(\mathcal{I}) = c_{\text{int}}(\mathcal{I}) = 1$. This implies that the equations are irredundant [Bel3, Res1] (cf. Remark 6.3.4), and it can also be explicitly checked in the examples above. In general, however, this is not the case, and so the Horn inequalities are still redundant. An example of such a Horn triple is the one given in Example 4.2.1.

Acknowledgements

We would like to thank Velleda Baldoni and Vladimir Popov for interesting discussions. We also thank Prakash Belkale and Cass Sherman for their helpful remarks. We would also like to thank an anonymous referee for their suggestions. MW acknowledges financial support by the Simons Foundation and AFOSR grant FA9550-16-1-0082.

References

- [BCM] P. BÜRGISSER, M. CHRISTANDL, K. D. MULMULEY, and M. WALTER, Membership in moment polytopes is in NP and coNP. *SIAM Journal on Computing* **46** (2017), 972–991. arXiv:1511.03675. Zbl1371.68105 MR 3662037
- [Bel1] P. BELKALE, Local systems on $\mathbb{P}^1 - S$ for S a finite set. *Compositio Mathematica* **129** (2001), 67–86. Zbl1042.14031 MR 1856023
- [Bel2] — Invariant theory of $GL(n)$ and intersection theory of Grassmannians. *International Mathematics Research Notices* **2004** (2004), 3709–3721. Zbl1082.14050 MR 2099498
- [Bel3] — Geometric proofs of Horn and saturation conjectures. *Journal of Algebraic Geometry* **15** (2006), 133–173. arXiv:math/0208107. Zbl1090.14014 MR 2177198
- [Bel4] — Geometric proof of a conjecture of Fulton. *Advances in Mathematics* **216** (2007), 346–357. arXiv:math/0511664. Zbl1129.14063 MR 2353260
- [BLMW] P. BÜRGISSER, J. M. LANDSBERG, L. MANIVEL and J. WEYMAN, An Overview of Mathematical Issues Arising in the Geometric Complexity Theory Approach to $VP \neq VNP$. *SIAM Journal on Computing* **40** (2011), 1179–1209. arXiv:0907.2850. Zbl1252.68134 MR 2861717
- [DW] H. DERKSEN and J. WEYMAN, Semi-invariants of quivers and saturation for Littlewood-Richardson coefficients. *Journal of the American Mathematical Society* **13** (2000), 467–479. Zbl0993.16011 MR 1758750
- [Ful1] W. FULTON, *Young tableaux: with applications to representation theory and geometry*, vol. 35. Cambridge University Press (1997). Zbl0878.14034 MR 1464693
- [Ful2] — Eigenvalues, invariant factors, highest weights, and Schubert calculus. *Bulletin of the American Mathematical Society* **37** (2000), 209–249. arXiv:math/9908012. Zbl0994.15021 MR 1754641
- [Hor] A. HORN, Eigenvalues of sums of Hermitian matrices. *Pacific Journal of Mathematics* **12** (1962), 225–241. Zbl0112.01501 MR 0140521
- [HTW] R. E. HOWE, E.-C. TAN and J. F. WILLENBRING, A basis for the GL_n tensor product algebra. *Advances in Mathematics* **196** (2005), 531–564. arXiv:math/0407468. Zbl1072.22007 MR 2166314
- [Ike] C. IKENMEYER, Small Littlewood-Richardson coefficients. *Journal of Algebraic Combinatorics* **44** (2012), 1–29. arXiv:1209.1521.
- [IMW] C. IKENMEYER, K. D. MULMULEY and M. WALTER, On vanishing of Kronecker coefficients. *Computational Complexity* **26** (2017), 949–992. arXiv:1507.02955. Zbl1382.68093 MR 3723353
- [Kly] A. A. KLYACHKO, Stable bundles, representation theory and Hermitian operators. *Selecta Mathematica, New Series* **4** (1998), 419–445. Zbl0915.14010 MR 1654578

- [KN] G. KEMPF and L. NESS, The length of vectors in representation spaces. In *Algebraic geometry*. Springer (1979). pp. 233–243. Zbl 0407.22012 MR 0555701
- [Knu] A. KNUTSON, The symplectic and algebraic geometry of Horn’s problem. *Linear Algebra and its Applications* **319** (2000), 61–81. arXiv:math/9911088. Zbl 0981.15010 MR 1799624
- [KT] A. KNUTSON and T. TAO, The honeycomb model of GL_n tensor products I: Proof of the saturation conjecture. *Journal of the American Mathematical Society* **12** (1999), 1055–1090. arXiv:math/9807160. Zbl 0944.05097 MR 1671451
- [KTW] A. KNUTSON, T. TAO and C. WOODWARD, The honeycomb model of GL_n tensor products II: Puzzles determine facets of the Littlewood–Richardson cone. *Journal of the American Mathematical Society* **17** (2004), 19–48. arXiv:math/0107011. Zbl 1043.05111 MR 2015329
- [MNS] K. D. MULMULEY, H. NARAYANAN and M. SOHONI, Geometric complexity theory III: on deciding nonvanishing of a Littlewood–Richardson coefficient. *Journal of Algebraic Combinatorics* **36** (2012), 103–110. Zbl 1271.03055 MR 2927658
- [Mul] K. D. MULMULEY, On P vs. NP and geometric complexity theory. *Journal of the ACM (JACM)* **58** (2011). Zbl 1327.68111 MR 2786586
- [Per] D. PERRIN, *Algebraic Geometry: An Introduction*. Springer Science & Business Media (2007).
- [Pur1] K. PURBHOO, <http://www.math.uwaterloo.ca/~kpurbhoo/papers/>.
- [Pur2] — Two step flag manifolds and the Horn conjecture. arXiv:math/0603131 (2006).
- [Res1] N. RESSAYRE, Geometric invariant theory and the generalized eigenvalue problem. *Inventiones mathematicae* **180** (2010), 389–441. arXiv:0704.2127. Zbl 1197.14051 MR 2609246
- [Res2] — A cohomology-free description of eigencones in types A , B , and C . *International Mathematics Research Notices* **2012** (2011), 4966–5005. arXiv:0908.4557. Zbl 1270.14026 MR 2993441
- [Res3] — A short geometric proof of a conjecture of Fulton. *L’Enseignement Mathématique* **57** (2011), 103–115. arXiv:0901.3633. Zbl 1234.14035 MR 2850586
- [Sch] A. SCHOFIELD, General representations of quivers. *Proceedings of the London Mathematical Society* **3** (1992), 46–64. Zbl 0795.16008 MR 1162487
- [She] C. SHERMAN, Geometric proof of a conjecture of King, Tollu, and Toumazet. arXiv:1505.06551 (2015).
- [VW] M. VERGNE and M. WALTER, Inequalities for moment cones of finite-dimensional representations. *Journal of Symplectic Geometry* **15** (2017), 1209–1250. arXiv:1410.8144. Zbl 06865057 MR 3734613

(Reçu le 3 juin 2017)

Nicole BERLINE, Centre de Mathématiques Laurent Schwartz, Ecole Polytechnique,
13 avenue de Tresum, 74000 Annecy, France

e-mail: nicole.berline@math.cnrs.fr

Michèle VERGNE, Université Paris-Diderot Paris 7, IMJ-PRG,
Bâtiment Sophie Germain, Case 7012, 75205 Paris Cedex 13, France

e-mail: michele.vergne@imj-prg.fr

Michael WALTER, Korteweg-de Vries Institute for Mathematics, Institute for
Theoretical Physics, and Institute for Logic, Language and Computation,
University of Amsterdam, P.O. Box 94248, 1090 GE Amsterdam, The Netherlands;
QuSoft, Amsterdam; Stanford Institute for Theoretical Physics, Stanford University,
USA

e-mail: m.walter@uva.nl