

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 64 (2018)
Heft: 3-4

Artikel: An elementary and unified proof of Grothendieck's inequality
Autor: Friedland, Shmuel / Lim, Lek-Heng / Zhang, Jinjie
DOI: <https://doi.org/10.5169/seals-842098>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 22.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

An elementary and unified proof of Grothendieck's inequality

Shmuel FRIEDLAND, Lek-Heng LIM and Jinjie ZHANG

Abstract. We present an elementary, self-contained proof of Grothendieck's inequality that unifies the real and complex cases and yields both the Krivine and Haagerup bounds, the current best-known explicit bounds for the real and complex Grothendieck constants respectively. This article is intended to be pedagogical, combining and streamlining known ideas of Lindenstrauss–Pełczyński, Krivine, and Haagerup into a proof that need only univariate calculus, basic complex variables, and a modicum of linear algebra as prerequisites.

Mathematics Subject Classification (2010). Primary: 47A07; Secondary: 46B28, 46B85

Keywords. Grothendieck inequality, Grothendieck constant, Krivine bound, Haagerup bound.

1. Introduction

We will let $\mathbb{F} = \mathbb{R}$ or \mathbb{C} throughout this article. In 1953, Grothendieck proved a powerful result that he called “the fundamental theorem in the metric theory of tensor products” [Gro]; he showed that there exists a finite constant $K > 0$ such that for every $l, m, n \in \mathbb{N}$ and every matrix $M = (M_{ij}) \in \mathbb{F}^{m \times n}$,

$$(1) \quad \max_{\|x_i\| = \|y_j\| = 1} \left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} \langle x_i, y_j \rangle \right| \leq K \max_{|\varepsilon_i| = |\delta_j| = 1} \left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} \varepsilon_i \delta_j \right|$$

where $\langle \cdot, \cdot \rangle$ is the standard inner product in \mathbb{F}^l , the maximum on the left is taken over all $x_i, y_j \in \mathbb{F}^l$ of unit 2-norm, and the maximum on the right is taken over all $\varepsilon_i, \delta_j \in \mathbb{F}$ of unit absolute value (i.e., $\varepsilon_i = \pm 1$, $\delta_j = \pm 1$ over \mathbb{R} ; $\varepsilon_i = e^{i\theta_i}$, $\delta_j = e^{i\phi_j}$ over \mathbb{C}). The inequality (1) has since been christened *Grothendieck's inequality* and the smallest possible constant K *Grothendieck's*

constant. The value of Grothendieck's constant depends on the choice of \mathbb{F} and we will denote it by $K_G^{\mathbb{F}}$.

Over the last 65 years, there have been many attempts to improve and simplify the proof of Grothendieck's inequality, and also to obtain better bounds for the Grothendieck constant $K_G^{\mathbb{F}}$, whose exact value remains unknown. The following are some major milestones:

- (i) The central result of Grothendieck's original paper [Gro] is that his eponymous inequality holds with $\pi/2 \leq K_G^{\mathbb{R}} \leq \sinh(\pi/2) \approx 2.301$ and $1.273 \approx 4/\pi \leq K_G^{\mathbb{C}}$. Grothendieck relied on the sign function for the real case and obtained the complex case from the real case via a complexification argument.
- (ii) The power of Grothendieck's inequality was not generally recognized until the work of Lindenstrauss and Pełczyński [LP] 15 years later, which connected the inequality to absolutely p -summing operators. They elucidated and improved Grothendieck's proof in the real case by computing expectations of sign functions and using Taylor expansions, although they did not get better bounds for $K_G^{\mathbb{R}}$.
- (iii) Rietz [Rie] obtained a slightly smaller bound $K_G^{\mathbb{R}} \leq 2.261$ in 1974 by averaging over \mathbb{R}^n with normalized Gaussian measure and using a variational argument to determine an optimal scalar map corresponding to the sign function.
- (iv) Our current best known upper bounds for $K_G^{\mathbb{R}}$ and $K_G^{\mathbb{C}}$ are due to Krivine [Kri], who in 1979 used Banach space theory and ideas in [LP] to get

$$K_G^{\mathbb{R}} \leq \frac{\pi}{2 \log(1 + \sqrt{2})} \approx 1.78221;$$

and Haagerup [Haa], who in 1987 extended Krivine's ideas to \mathbb{C} to get

$$K_G^{\mathbb{C}} \leq \frac{8}{\pi(x_0 + 1)} \approx 1.40491,$$

where $x_0 \in [0, 1]$ is the unique solution to:

$$x \int_0^{\pi/2} \frac{\cos^2 t}{\sqrt{1 - x^2 \sin^2 t}} dt = \frac{\pi}{8}(x + 1).$$

- (v) Our current best known lower bounds for $K_G^{\mathbb{R}}$ and $K_G^{\mathbb{C}}$ are due to Davie [Davl, Dav2], who in 1984 used spherical integrals to get

$$K_G^{\mathbb{R}} \geq \sup_{x \in (0,1)} \frac{1 - \rho(x)}{\max(\rho(x), f(x))} \approx 1.67696,$$

where

$$\rho(x) := \sqrt{\frac{2}{\pi}} x e^{-x^2/2}, \quad f(x) := \frac{2}{\pi} e^{-x^2} + \rho(x) \left[1 - \sqrt{\frac{8}{\pi}} \int_x^\infty e^{-t^2/2} dt \right];$$

and

$$K_G^C \geq \sup_{x>0} \frac{1 - \theta(x)}{g(x)} \approx 1.33807,$$

where

$$\theta(x) := \frac{1}{2} \left[1 - e^{-x^2} + x \int_x^\infty e^{-t^2} dt \right],$$

$$g(x) := \left[\frac{1}{x} (1 - e^{-x^2}) + \int_x^\infty e^{-t^2} dt \right]^2 + \theta(x) \left[1 - \frac{2}{x} (1 - e^{-x^2}) \right].$$

- (vi) Progress on improving the aforementioned bounds halted for many years. Believing that Krivine's bound is the exact value of K_G^R , some were spurred to find matrices that yield it as the lower bound of K_G^R [Kon1]. The belief was dispelled in 2011 in a landmark paper [Bra], which demonstrated the existence of a positive constant ε such that $K_G^R < \pi/(2 \log(1 + \sqrt{2})) - \varepsilon$ but the authors did not provide an explicit better bound. To date, Krivine's and Haagerup's bounds remain the best known explicit upper bounds for K_G^R and K_G^C respectively.
- (vii) There have also been many alternate proofs of Grothendieck's inequality employing a variety of techniques, among them factorization of Hilbert spaces [Mau, Jam, Pis1], absolutely summing operators [DJT, LP, Pis2], geometry of Banach spaces [AK, LT], metric theory of tensor product [DFS], basic probability theory [Ble], bilinear forms on C^* -algebra [Kai].

In this article, we will present a proof of Grothendieck's inequality that unifies both the (a) real and (b) complex cases; and yields both the (c) Krivine and (d) Haagerup bounds [Kri, Haa]. It is also elementary in that it requires little more than standard college mathematics. Our proof will rely on Lemma 2.1, which is a variation of known ideas in [LP, Haa, Jam]. In particular, the idea of using the sign function to establish (1) in the real case was due to Grothendieck himself [Gro] and later also appeared in [LP, Kri]; whereas the use of the sign function in the complex case first appeared in [Haa]. To be clear, all the key ideas in our proof were originally due to Lindenstrauss–Pełczyński, Krivine, Haagerup, and König [LP, Kri, Haa, Kon2], our only contribution is pedagogical – combining, simplifying, and streamlining their ideas into what we feel is a more palatable proof. To understand the proof, readers need only know univariate calculus, basic complex variables, and a small amount of linear algebra. We will use some basic Hilbert space theory and tensor product constructions in Section 4 but both notions will be explained in a self-contained and elementary way.

2. Gaussian integral of sign function

Throughout this article, our inner product over \mathbb{C} will be sesquilinear in the second argument, i.e.,

$$\langle x, y \rangle := y^* x \quad \text{for all } x, y \in \mathbb{C}^n.$$

For $z \in \mathbb{F} = \mathbb{R}$ or \mathbb{C} , the sign function is

$$(2) \quad \operatorname{sgn}(z) = \begin{cases} z/|z| & \text{if } z \neq 0, \\ 0 & \text{if } z = 0; \end{cases}$$

and for $z \in \mathbb{F}^n$, the Gaussian function is

$$G_n^{\mathbb{F}}(z) = \begin{cases} (2\pi)^{-n/2} \exp(-\|z\|_2^2/2) & \text{if } \mathbb{F} = \mathbb{R}, \\ \pi^{-n} \exp(-\|z\|_2^2) & \text{if } \mathbb{F} = \mathbb{C}. \end{cases}$$

Lemma 2.1 below is based on [Jam, Haa]; the complex version in particular is a slight variation of [Haa, Lemma 3.2]. It plays an important role in our proof because the right side of (3) depends only on the inner product $\langle u, v \rangle$ and not (explicitly) on the dimension n . In addition, the functions on the right are homeomorphisms and admit Taylor expansions, making it possible to expand them in powers $\langle u, v \rangle^d$, which will come in useful when we prove Theorem 4.1.

Lemma 2.1. *Let $u, v \in \mathbb{F}^n$ with $\|u\|_2 = \|v\|_2 = 1$. Then*

$$(3) \quad \begin{aligned} & \int_{\mathbb{F}^n} \operatorname{sgn}\langle u, z \rangle \operatorname{sgn}\langle z, v \rangle G_n^{\mathbb{F}}(z) dz \\ &= \begin{cases} \frac{2}{\pi} \arcsin\langle u, v \rangle & \text{if } \mathbb{F} = \mathbb{R}, \\ \langle u, v \rangle \int_0^{\pi/2} \frac{\cos^2 t}{(1 - |\langle u, v \rangle|^2 \sin^2 t)^{1/2}} dt & \text{if } \mathbb{F} = \mathbb{C}. \end{cases} \end{aligned}$$

Proof. CASE I: $\mathbb{F} = \mathbb{R}$. Let $\arccos\langle u, v \rangle = \theta$, so that $\theta \in [0, \pi]$ and $\arcsin\langle u, v \rangle = \pi/2 - \theta$. Choose α, β such that $0 < \beta - \alpha < \pi$ and define

$$E(\alpha, \beta) = \{(r \cos \theta, r \sin \theta, x_3, \dots, x_n) : r \geq 0, \alpha \leq \theta \leq \beta\}.$$

The Gaussian measure of a measurable set A is the integral of $G_n^{\mathbb{R}}(x)$ over A . Upon integrating with respect to x_3, \dots, x_n , the following term remains:

$$\frac{1}{2\pi} \int_{E(\alpha, \beta)} e^{-\frac{1}{2}(x_1^2 + x_2^2)} dx_1 dx_2 = \frac{1}{2\pi} \int_{\alpha}^{\beta} d\theta \int_0^{\infty} r e^{-\frac{1}{2}r^2} dr = (\beta - \alpha)/2\pi.$$

Hence the Gaussian measure of $E(\alpha, \beta)$ is $(\beta - \alpha)/2\pi$. Since there is an isometry T of \mathbb{R}^n such that $Tu = e_1$ and $Tv = (\cos \theta, \sin \theta, 0, \dots, 0)$, the left side of (3) may be expressed as

$$\int_{\mathbb{R}^n} \operatorname{sgn}\langle Tu, x \rangle \operatorname{sgn}\langle x, Tv \rangle G_n^{\mathbb{R}}(x) dx.$$

The set of x where $\langle Tu, x \rangle > 0$ and $\langle Tv, x \rangle > 0$ is $E(\theta - \pi/2, \pi/2)$, which has Gaussian measure $(\pi - \theta)/2\pi$; ditto for $\langle Tu, x \rangle < 0$ and $\langle Tv, x \rangle < 0$. The set of x where $\langle Tu, x \rangle < 0$ and $\langle Tv, x \rangle > 0$ is $E(\pi/2, \theta + \pi/2)$, which has Gaussian measure $\theta/2\pi$; ditto for $\langle Tu, x \rangle > 0$ and $\langle Tv, x \rangle < 0$. The set of x where $\langle Tu, x \rangle = 0$ has zero Gaussian measure. Hence the value of this integral is $(\pi - \theta)/2\pi + (\pi - \theta)/2\pi - \theta/2\pi - \theta/2\pi = 2 \arcsin\langle u, v \rangle / \pi$.

CASE II: $\mathbb{F} = \mathbb{C}$. We define vectors $\alpha, \beta \in \mathbb{R}^{2n}$ with $\alpha_{2i-1} = \operatorname{Re}(u_i)$, $\alpha_{2i} = \operatorname{Im}(u_i)$, $\beta_{2i-1} = \operatorname{Re}(v_i)$, $\beta_{2i} = \operatorname{Im}(v_i)$, $i = 1, \dots, n$. Then α and β are unit vectors in \mathbb{R}^{2n} . For any $z = (z_1, \dots, z_n) \in \mathbb{C}^n$, we write

$$x = (\operatorname{Re}(z_1), \operatorname{Im}(z_1), \dots, \operatorname{Re}(z_n), \operatorname{Im}(z_n)) \in \mathbb{R}^{2n}.$$

Then,

$$\begin{aligned} \operatorname{Re}(\langle u, z \rangle) &= \sum_{i=1}^n \operatorname{Re}(u_i \bar{z}_i) = \sum_{i=1}^n (\operatorname{Re}(u_i) \operatorname{Re}(z_i) + \operatorname{Im}(u_i) \operatorname{Im}(z_i)) \\ &= \langle \alpha, x \rangle = \langle x, \alpha \rangle, \end{aligned}$$

and likewise $\operatorname{Re}(\langle z, v \rangle) = \langle x, \beta \rangle$. By a change-of-variables and Case I, we have

$$\begin{aligned} (4) \quad \int_{\mathbb{C}^n} \operatorname{sgn}(\operatorname{Re}\langle u, z \rangle) \operatorname{sgn}(\operatorname{Re}\langle z, v \rangle) G_n^{\mathbb{C}}(z) dz &= \int_{\mathbb{R}^{2n}} \operatorname{sgn}\langle x, \alpha \rangle \operatorname{sgn}\langle x, \beta \rangle G_{2n}^{\mathbb{R}}(x) dx \\ &= \frac{2}{\pi} \arcsin\langle \alpha, \beta \rangle = \frac{2}{\pi} \arcsin(\operatorname{Re}\langle u, v \rangle). \end{aligned}$$

It is easy to verify that for any $z \in \mathbb{C}$,

$$(5) \quad \operatorname{sgn}(z) = \frac{1}{4} \int_0^{2\pi} \operatorname{sgn}(\operatorname{Re}(e^{-i\theta} z)) e^{i\theta} d\theta.$$

By (4), (5), and Fubini's theorem,

$$\begin{aligned} &\int_{\mathbb{C}^n} \operatorname{sgn}\langle u, z \rangle \operatorname{sgn}\langle z, v \rangle G_n^{\mathbb{C}}(z) dz \\ &= \frac{1}{16} \int_0^{2\pi} \int_0^{2\pi} \int_{\mathbb{C}^n} \operatorname{sgn}(\operatorname{Re}(\langle e^{-i\theta} u, z \rangle)) \\ &\quad \operatorname{sgn}(\operatorname{Re}(\langle z, \overline{e^{-i\varphi}} v \rangle)) e^{i(\theta+\varphi)} G_n^{\mathbb{C}}(z) dz d\theta d\varphi \\ &= \frac{1}{8\pi} \int_0^{2\pi} \int_0^{2\pi} \arcsin(\operatorname{Re}(\langle e^{-i\theta} u, e^{i\varphi} v \rangle)) e^{i(\theta+\varphi)} d\theta d\varphi \end{aligned}$$

CASE II(a): $\langle u, v \rangle \in \mathbb{R}$. The integral above becomes

$$\begin{aligned}
 &= \frac{1}{8\pi} \int_0^{2\pi} \left[\int_0^{2\pi} \arcsin(\cos(\theta + \varphi) \langle u, v \rangle) e^{i(\theta+\varphi)} d\theta \right] d\varphi \\
 &= \frac{1}{8\pi} \int_0^{2\pi} \left[\int_{\varphi}^{2\pi+\varphi} \arcsin(\langle u, v \rangle \cos t) e^{it} dt \right] d\varphi \\
 &= \frac{1}{8\pi} \int_0^{2\pi} \left[\int_0^{2\pi} \arcsin(\langle u, v \rangle \cos t) e^{it} dt \right] d\varphi \\
 (6) \quad &= \frac{1}{4} \int_0^{2\pi} \arcsin(\langle u, v \rangle \cos t) e^{it} dt.
 \end{aligned}$$

Since $\arcsin(\langle u, v \rangle \cos t)$ is an even function with period 2π ,

$$\int_0^{2\pi} \arcsin(\langle u, v \rangle \cos t) \sin t dt = 0,$$

the last integral in (6) becomes

$$\frac{1}{4} \int_0^{2\pi} \arcsin(\langle u, v \rangle \cos t) \cos t dt,$$

and as $\arcsin(\langle u, v \rangle \cos t) \cos t$ is an even function with period π , it becomes

$$\int_0^{\pi/2} \arcsin(\langle u, v \rangle \cos t) \cos t dt = \int_0^{\pi/2} \arcsin(\langle u, v \rangle \sin t) \sin t dt,$$

which, upon integrating by parts, becomes

$$(7) \quad \langle u, v \rangle \int_0^{\pi/2} \frac{\cos^2 t}{(1 - |\langle u, v \rangle|^2 \sin^2 t)^{1/2}} dt.$$

CASE II(b): $\langle u, v \rangle \notin \mathbb{R}$. This reduces to Case II(a) by setting $c \in \mathbb{C}$ of unit modulus so that $c\langle u, v \rangle = |\langle u, v \rangle|$ and $\langle cu, v \rangle \in \mathbb{R}$, then by (7),

$$\begin{aligned}
 \int_{\mathbb{C}^n} \operatorname{sgn}\langle u, z \rangle \operatorname{sgn}\langle z, v \rangle G_n^{\mathbb{C}}(z) dz &= \bar{c} \int_{\mathbb{C}^n} \operatorname{sgn}\langle cu, z \rangle \operatorname{sgn}\langle z, v \rangle G_n^{\mathbb{C}}(z) dz \\
 &= \bar{c} \langle cu, v \rangle \int_0^{\pi/2} \frac{\cos^2 t}{(1 - |\langle cu, v \rangle|^2 \sin^2 t)^{1/2}} dt \\
 &= \langle u, v \rangle \int_0^{\pi/2} \frac{\cos^2 t}{(1 - |\langle u, v \rangle|^2 \sin^2 t)^{1/2}} dt.
 \end{aligned}$$

□

We will make a simple but useful observation¹ about the quantities in (1) that we will need for the proof of Corollary 2.3 later.

Lemma 2.2. *Let $\mathbb{F} = \mathbb{R}$ or \mathbb{C} and $d, m, n \in \mathbb{N}$. For any $M \in \mathbb{F}^{m \times n}$, we have*

$$(8) \quad \max_{|\varepsilon_i| \leq 1, |\delta_j| \leq 1} \left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} \varepsilon_i \delta_j \right| = \max_{|\varepsilon_i| = |\delta_j| = 1} \left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} \varepsilon_i \delta_j \right|$$

and for any $x_1, \dots, x_m, y_1, \dots, y_n \in \mathbb{F}^d$,

$$(9) \quad \max_{\|x_i\| \leq 1, \|y_j\| \leq 1} \left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} \langle x_i, y_j \rangle \right| = \max_{\|x_i\| = \|y_j\| = 1} \left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} \langle x_i, y_j \rangle \right|.$$

Proof. We will start with (8). Suppose there exists $M \in \mathbb{F}^{m \times n}$ such that the left-hand side of (8) exceeds the right-hand side. Let the maximum of the left-hand side be attained by $\varepsilon_1^*, \dots, \varepsilon_m^*$ and $\delta_1^*, \dots, \delta_n^*$. By our assumption, at least one ε_i^* or δ_j^* must be less than 1 in absolute value and so let $|\varepsilon_1^*| < 1$ without loss of generality. Fix $\varepsilon_i = \varepsilon_i^*$, $i = 2, \dots, m$ and $\delta_j = \delta_j^*$, $j = 1, \dots, n$, but let ε_1 vary with $|\varepsilon_1| \leq 1$ and consider the maximum of the left hand-side over ε_1 . Since $\max\{|a\varepsilon_1 + b| : |\varepsilon_1| \leq 1\}$ is always attained on the boundary $|\varepsilon_1| = 1$ for any $a, b \in \mathbb{F}$, this contradicts our assumption. The proof for (9) is similar with norm in place of absolute value. \square

In the corollary below, the inequality on the left is the “original Grothendieck inequality”, i.e., as first stated by Grothendieck² in [Gro], and the inequality on the right is due to Haagerup [Haa].

Corollary 2.3. *Let $\mathbb{F} = \mathbb{R}$ or \mathbb{C} and $d, m, n \in \mathbb{N}$. For any $M \in \mathbb{F}^{m \times n}$ with*

$$(10) \quad \max_{|\varepsilon_i| = |\delta_j| = 1} \left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} \varepsilon_i \delta_j \right| \leq 1,$$

any $x_1, \dots, x_m, y_1, \dots, y_n \in \mathbb{F}^d$ of unit 2-norm, we have

$$\left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} \arcsin \langle x_i, y_j \rangle \right| \leq \frac{\pi}{2} \quad \text{if } \mathbb{F} = \mathbb{R},$$

$$\left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} H(\langle x_i, y_j \rangle) \right| \leq 1 \quad \text{if } \mathbb{F} = \mathbb{C},$$

where H denotes the function on the right side of (3) for $\mathbb{F} = \mathbb{C}$.

¹This of course follows from other well-known results but we would like to keep our exposition self-contained.

²The better known modern version (1) is in fact due to Lindenstrauss and Pełczyński in [LP].

Proof. The condition (10) implies that

$$\left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} \operatorname{sgn}\langle x_i, x \rangle \operatorname{sgn}\langle y_j, x \rangle G_d^{\mathbb{R}}(z) \right| \leq G_d^{\mathbb{R}}(z),$$

$$\left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} \operatorname{sgn}\langle z, \bar{x}_i \rangle \overline{\operatorname{sgn}\langle z, \bar{y}_j \rangle} G_d^{\mathbb{C}}(z) \right| \leq G_d^{\mathbb{C}}(z),$$

for any $x \in \mathbb{R}^d$, $z \in \mathbb{C}^d$ respectively. Integrating over \mathbb{R}^d or \mathbb{C}^d respectively and applying Lemma 2.1 give the required results. Note that we have implicitly relied on (8) in Lemma 2.2 as the sgn function is not always of absolute value one and may be zero. \square

Corollary 2.3 already looks a lot like the Grothendieck inequality (1) but the nonlinear functions \arcsin and H are in the way. To obtain the Grothendieck inequality, we linearize them: First by using Taylor series to replace these functions by polynomials; and then using a ‘tensor trick’ to express the polynomials as linear functions on a larger space. This is the gist of the proof in Section 4.

3. Haagerup function

We will need to make a few observations regarding the functions on the right side of (3) for the proof of Grothendieck’s inequality. Let the *complex Haagerup function* of a complex variable z be

$$H(z) := z \int_0^{\pi/2} \frac{\cos^2 t}{(1 - |z|^2 \sin^2 t)^{1/2}} dt, \quad |z| \leq 1,$$

and the *real Haagerup function* h as the restriction of H to $[-1, 1] \subseteq \mathbb{R}$. Observe that $h : [-1, 1] \rightarrow [-1, 1]$ and is a strictly increasing continuous bijection. Since $[-1, 1]$ is compact, h is a homeomorphism of $[-1, 1]$ onto itself. By the Taylor expansion

$$(1 - x^2 \sin^2 t)^{-1/2} = \sum_{k=0}^{\infty} \frac{(2k-1)!!}{(2k)!!} x^{2k} \sin^{2k} t, \quad |x| \leq 1, \quad 0 \leq t < \pi/2,$$

and

$$\int_0^{\pi/2} \cos^2 t \sin^{2k} t dt = \frac{\pi}{4(k+1)} \frac{(2k-1)!!}{(2k)!!},$$

thus we get

$$(11) \quad h(x) = \sum_{k=0}^{\infty} \frac{\pi}{4(k+1)} \left[\frac{(2k-1)!!}{(2k)!!} \right]^2 x^{2k+1}, \quad x \in [-1, 1].$$

Since h is analytic at $x = 0$ and $h'(0) \neq 0$, its inverse function $h^{-1} : [-1, 1] \rightarrow [-1, 1]$ can be expanded in a power series in some neighborhood of 0

$$(12) \quad h^{-1}(x) = \sum_{k=0}^{\infty} b_{2k+1} x^{2k+1}.$$

One may in principle determine the coefficients using the Lagrange inversion formula:

$$b_{2k+1} = \frac{1}{(2k+1)!} \lim_{t \rightarrow 0} \left[\frac{d^{2k}}{dt^{2k}} \left(\frac{t}{h(t)} \right)^{2k+1} \right].$$

For example,

$$b_1 = \frac{4}{\pi}, \quad b_3 = -\frac{1}{8} \left(\frac{4}{\pi} \right)^3, \quad b_5 = 0, \quad b_7 = -\frac{1}{1024} \left(\frac{4}{\pi} \right)^7.$$

But determining b_{2k+1} explicitly becomes difficult as k gets larger. A key step in Haagerup's proof [Haa] requires the nonpositivity of the coefficients beyond the first:

$$(13) \quad b_{2k+1} \leq 0, \quad \text{for all } k \geq 1.$$

This step is in our view the most technical part of [Haa]. We have no insights on how it may be avoided but we simplified Haagerup's proof of (13) in Section 5 to keep to our promise of an elementary proof – using only calculus and basic complex variables.

It follows from (13) that $\tilde{h}(z) := b_1 z - h^{-1}(z)$ has nonnegative Taylor coefficients. Pringsheim's theorem implies that if the radius of convergence of the Taylor series of $\tilde{h}(z)$ is r , then $\tilde{h}(z)$, and thus $h^{-1}(z)$, has a singular point at $z = r$. As $h'(t) > 0$ on $(0, 1)$ and $h(1) = 1$, we must have $r \geq 1$. It also follows from (13) that $h^{-1}(t) \leq \sum_{k=0}^N b_{2k+1} t^{2k+1}$ for any $t \in (0, 1)$ and $N \in \mathbb{N}$. So $\sum_{k=1}^N |b_{2k+1}| t^{2k+1} \leq b_1 t - h^{-1}(t)$ for any $t \in (0, 1)$ and $N \in \mathbb{N}$. So $\sum_{k=1}^N |b_{2k+1}| \leq b_1 - 1$ for any $N \in \mathbb{N}$ and we have $\sum_{k=0}^{\infty} |b_{2k+1}| \leq 2b_1 - 1$. As $h^{-1}(1) = 1$ we deduce that $\sum_{k=0}^{\infty} b_{2k+1} = h^{-1}(1) = 1$, and therefore

$$(14) \quad \sum_{k=0}^{\infty} |b_{2k+1}| = 2b_1 - 1.$$

We now turn our attention back to the complex Haagerup function. Observe that $|H(z)| = h(|z|)$ for all $z \in D := \{z \in \mathbb{C} : |z| \leq 1\}$ and $\arg(H(z)) = \arg(z)$ for $0 \neq z \in D$. So $H : D \rightarrow D$ is a homeomorphism of D onto itself. Let $H^{-1} : D \rightarrow D$ be its inverse function. Since $H(z) = \operatorname{sgn}(z)h(|z|)$, we get

$$(15) \quad H^{-1}(z) = \operatorname{sgn}(z)h^{-1}(|z|) = \operatorname{sgn}(z) \sum_{k=0}^{\infty} b_{2k+1} |z|^{2k+1}.$$

Dini's theorem shows that the function $\varphi(x) := \sum_{k=0}^{\infty} |b_{2k+1}|x^{2k+1}$ is a strictly increasing and continuous on $[0, 1]$, with $\varphi(0) = 0$ and $\varphi(1) = \sum_{k=0}^{\infty} |b_{2k+1}| \geq b_1 = 4/\pi > 1$; note that $\varphi(1)$ is finite by (14). Thus there exists a unique $c_0 \in (0, 1)$ such that $\varphi(c_0) = 1$. So

$$1 = \varphi(c_0) = \sum_{k=0}^{\infty} |b_{2k+1}|c_0^{2k+1} = \frac{8}{\pi}c_0 - h^{-1}(c_0),$$

where the last equality follows from $b_1 = 4/\pi$ and (13). Therefore we obtain $h^{-1}(c_0) = 8c_0/\pi - 1$, and if we let $x_0 := h^{-1}(c_0) \in (0, 1)$, then $h(x_0) - \pi(x_0 + 1)/8 = 0$. From the Taylor expansion of $h(x)$, the function $x \mapsto h(x) - \pi(x + 1)/8$ is increasing and continuous on $[0, 1]$. Hence x_0 is the unique solution in $[0, 1]$ to

$$(16) \quad h(x) - \frac{\pi}{8}(x + 1) = 0$$

and $c_0 = \pi(x_0 + 1)/8$.

As Corollary 2.3 indicates, the Haagerup function H plays the analogue of \arcsin in the complex case. Unlike \arcsin , H is a completely obscure function,³ and any of its properties that we require will have to be established from scratch. The goal of this section is essentially to establish (11)–(16), which we will need later.

4. A unified proof of Grothendieck's inequality

In this section we will need the notions of (i) tensor product and (ii) Hilbert space, but just enough to make sense of $\mathcal{T}(\mathbb{F}^n) = \bigoplus_{k=0}^{\infty} (\mathbb{F}^n)^{\otimes k}$ where $\mathbb{F} = \mathbb{R}$ or \mathbb{C} . In keeping to our promise of an elementary proof, we will briefly introduce these notions in a simple manner. For our purpose, it suffices to regard the *tensor product* of k copies of \mathbb{F}^n , denoted

$$(\mathbb{F}^n)^{\otimes k} = \underbrace{\mathbb{F}^n \otimes \cdots \otimes \mathbb{F}^n}_{k \text{ copies}},$$

as the \mathbb{F} -vector space of k -dimensional *hypermatices*,

$$(\mathbb{F}^n)^{\otimes k} := \{[a_{i_1 \dots i_k}] : a_{i_1 \dots i_k} \in \mathbb{F}, i_1, \dots, i_k \in \{1, \dots, n\}\},$$

where scalar multiplication and vector addition of hypermatrices are defined coordinatewise. Also, we let $(\mathbb{F}^n)^{\otimes 0} := \mathbb{F}$. For k vectors $x, y, \dots, z \in \mathbb{F}^n$, their *tensor product* is the k -dimensional hypermatrix given by

³We are unaware of any other occurrence of H outside its use in Haagerup's proof of his bound in [Haa].

$$x \otimes y \otimes \cdots \otimes z := [x_{i_1} y_{i_2} \cdots z_{i_k}]_{i_1, i_2, \dots, i_k=1}^n \in (\mathbb{F}^n)^{\otimes k}.$$

We write

$$x^{\otimes k} := \underbrace{x \otimes \cdots \otimes x}_{k \text{ copies}}.$$

If $\langle \cdot, \cdot \rangle$ is an inner product on \mathbb{F}^n , then defining

$$(17) \quad \langle x \otimes y \otimes \cdots \otimes z, x' \otimes y' \otimes \cdots \otimes z' \rangle := \langle x, x' \rangle \langle y, y' \rangle \cdots \langle z, z' \rangle$$

and extending bilinearly (if $\mathbb{F} = \mathbb{R}$) or sesquilinearly (if $\mathbb{F} = \mathbb{C}$) to all of $(\mathbb{F}^n)^{\otimes k}$ yields an inner product on the k -dimensional hypermatrices. In particular we have

$$\langle x^{\otimes k}, y^{\otimes k} \rangle = \langle x, y \rangle^k.$$

If $\{e_1, \dots, e_n\}$ is the standard orthonormal basis of \mathbb{F}^n , then

$$(18) \quad \{e_{i_1} \otimes \cdots \otimes e_{i_k} \in (\mathbb{F}^n)^{\otimes k} : i_1, \dots, i_k \in \{1, \dots, n\}\}$$

is an orthonormal basis of $(\mathbb{F}^n)^{\otimes k}$. For more information about hypermatrices see [Lim] and for a more formal definition of tensor products see [FA].

If an \mathbb{F} -vector space \mathcal{H} is equipped with an inner product $\langle \cdot, \cdot \rangle$ such that every Cauchy sequence in \mathcal{H} converges with respect to the induced norm $\|v\| = |\langle v, v \rangle|^{1/2}$, we call \mathcal{H} a *Hilbert space*. Hilbert spaces need not be finite-dimensional; we call \mathcal{H} *separable* if there is a countable set of orthonormal vectors $\{e_j \in \mathcal{H} : j \in J\}$, i.e., J is a countable index set, such that every $v \in \mathcal{H}$ satisfies

$$(19) \quad \|v\|^2 = \sum_{j \in J} |\langle v, e_j \rangle|^2.$$

Let $\langle \cdot, \cdot \rangle_k$ be the inner product on $(\mathbb{F}^n)^{\otimes k}$ as defined in (17), $\|\cdot\|_k$ be its induced norm, and B_k be the orthonormal basis in (18). Let $n \in \mathbb{N}$. The *tensor algebra* of \mathbb{F}^n is the \mathbb{F} -vector space⁴

(20)

$$\mathcal{T}(\mathbb{F}^n) := \bigoplus_{k=0}^{\infty} (\mathbb{F}^n)^{\otimes k} = \left\{ (v_0, v_1, v_2, \dots) : v_k \in (\mathbb{F}^n)^{\otimes k}, \sum_{k=0}^{\infty} \|v_k\|_k^2 < \infty \right\}$$

equipped with the inner product

$$(21) \quad \langle u, v \rangle_* := \sum_{k=0}^{\infty} \langle u_k, v_k \rangle_k.$$

It is a separable Hilbert space since $\bigcup_{k=0}^{\infty} B_k$ is a countable set of orthonormal vectors satisfying (19). We write $\|\cdot\|_*$ for the norm induced by (21).

⁴The direct sum in (20) is a Hilbert space direct sum, i.e., it is the closure of the vector space direct sum.

Theorem 4.1 (Grothendieck inequality with Krivine and Haagerup bounds). *Let $\mathbb{F} = \mathbb{R}$ or \mathbb{C} and $l, m, n \in \mathbb{N}$. For any $M \in \mathbb{F}^{m \times n}$, any $x_1, \dots, x_m, y_1, \dots, y_n \in \mathbb{F}^l$ of unit 2-norm, we have*

$$(22) \quad \max_{\|x_i\| = \|y_j\| = 1} \left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} \langle x_i, y_j \rangle \right| \leq K^{\mathbb{F}} \max_{|\varepsilon_i| = |\delta_j| = 1} \left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} \varepsilon_i \delta_j \right|,$$

where

$$K^{\mathbb{R}} := \frac{\pi}{2 \log(1 + \sqrt{2})} \quad \text{and} \quad K^{\mathbb{C}} := \frac{8}{\pi(x_0 + 1)}$$

are Krivine's and Haagerup's bounds respectively. Recall that x_0 is as defined in (16).

Proof. As we described at the end of Section 2, we will 'linearize' the nonlinear functions arcsin and H in Corollary 2.3 by using Taylor series to replace these functions by polynomials, followed by a 'tensor trick' [Jam, Kri] to express polynomials as linear functions on an infinite-dimensional space.

CASE I: $\mathbb{F} = \mathbb{R}$. Let $c := \text{arcsinh}(1) = \log(1 + \sqrt{2})$. Taylor expansion gives

$$(23) \quad \begin{aligned} \sin(c \langle x_i, y_j \rangle) &= \sum_{k=0}^{\infty} (-1)^k \frac{c^{2k+1}}{(2k+1)!} \langle x_i, y_j \rangle^{2k+1} \\ &= \sum_{k=0}^{\infty} (-1)^k \frac{c^{2k+1}}{(2k+1)!} \langle x_i^{\otimes(2k+1)}, y_j^{\otimes(2k+1)} \rangle_k. \end{aligned}$$

For any $l \in \mathbb{N}$, let $\mathcal{T}(\mathbb{R}^l)$ be as in (20), and $S, T : \mathbb{R}^l \rightarrow \mathcal{T}(\mathbb{R}^l)$ be nonlinear maps defined by

$$\begin{aligned} S(x) &:= (S_k(x))_{k=0}^{\infty}, \quad S_{2k}(x) := 0, \quad S_{2k+1}(x) := (-1)^k \sqrt{\frac{c^{2k+1}}{(2k+1)!}} \cdot x^{\otimes(2k+1)}, \\ T(x) &:= (T_k(x))_{k=0}^{\infty}, \quad T_{2k}(x) := 0, \quad T_{2k+1}(x) := \sqrt{\frac{c^{2k+1}}{(2k+1)!}} \cdot x^{\otimes(2k+1)}, \end{aligned}$$

for any $x \in \mathbb{R}^l$. To justify that S and T are indeed maps into $\mathcal{T}(\mathbb{R}^l)$, we need to demonstrate that $\|S(x)\|_*, \|T(x)\|_* < \infty$ but this follows from

$$\|S(x)\|_*^2 = \sum_{k=0}^{\infty} \|S_k(x)\|_k^2 = \sum_{k=0}^{\infty} \frac{c^{2k+1}}{(2k+1)!} \|x\|^{2(2k+1)} = \sum_{k=0}^{\infty} \|T_k(x)\|_k^2 = \|T(x)\|_*^2$$

and

$$\sum_{k=0}^{\infty} \frac{c^{2k+1}}{(2k+1)!} \|x\|^{2(2k+1)} = \sinh(c\|x\|^2) < \infty$$

for all $x \in \mathbb{R}^l$. Note that

$$\langle S(x), T(y) \rangle_* = \sum_{k=0}^{\infty} (-1)^k \frac{c^{2k+1}}{(2k+1)!} \langle x, y \rangle^{2k+1} = \sin(c \langle x, y \rangle),$$

which is the essence of the ‘tensor trick’. Hence (23) becomes:

$$\sin(c \langle x_i, y_j \rangle) = \langle S(x_i), T(y_j) \rangle_* \quad \text{or} \quad c \langle x_i, y_j \rangle = \arcsin \langle S(x_i), T(y_j) \rangle_*.$$

Moreover, since x_i and y_j are unit vectors in \mathbb{R}^l , we get

$$\|S(x_i)\|^2 = \sinh(c \|x_i\|^2) = 1 \quad \text{and} \quad \|T(y_j)\|^2 = \sinh(c \|y_j\|^2) = 1.$$

As the $m+n$ vectors $S(x_1), \dots, S(x_m), T(y_1), \dots, T(y_n)$ in $\mathcal{T}(\mathbb{R}^l)$ span a subspace $\mathcal{S} \subseteq \mathcal{T}(\mathbb{R}^l)$ of dimension $d \leq m+n$; and since any two finite-dimensional inner product spaces are isometric, \mathcal{S} is isometric to \mathbb{R}^d with the standard inner product. So we may apply Corollary 2.3 to obtain

$$\left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} \langle x_i, y_j \rangle \right| = \frac{1}{c} \left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} \arcsin \langle S(x_i), T(y_j) \rangle_* \right| \leq \frac{\pi}{2c},$$

which is Krivine's bound since $\pi/2c = \pi/(2 \log(1 + \sqrt{2})) = K^{\mathbb{R}}$.

CASE II: $\mathbb{F} = \mathbb{C}$. Let $c_0 \in (0, 1)$ be the unique constant defined in (16) such that $\varphi(c_0) = 1$. By the Taylor expansion in (15) and noting that $\operatorname{sgn}(z)|z|^{2k+1} = \bar{z}^k z^{k+1}$,

$$\begin{aligned} (24) \quad H^{-1}(c_0 \langle x_i, y_j \rangle) &= \operatorname{sgn}(c_0 \langle x_i, y_j \rangle) \sum_{k=0}^{\infty} b_{2k+1} |c_0 \langle x_i, y_j \rangle|^{2k+1} \\ &= \sum_{k=0}^{\infty} b_{2k+1} c_0^{2k+1} \overline{\langle x_i, y_j \rangle}^k \langle x_i, y_j \rangle^{k+1} \\ &= \sum_{k=0}^{\infty} b_{2k+1} c_0^{2k+1} \langle \bar{x}_i, \bar{y}_j \rangle^k \langle x_i, y_j \rangle^{k+1} \\ &= \sum_{k=0}^{\infty} b_{2k+1} c_0^{2k+1} \left\langle \bar{x}_i^{\otimes k} \otimes x_i^{\otimes(k+1)}, \bar{y}_j^{\otimes k} \otimes y_j^{\otimes(k+1)} \right\rangle_k. \end{aligned}$$

For any $l \in \mathbb{N}$, let $D_l = \{x \in \mathbb{C}^l : \|x\| \leq 1\}$ be the unit ball, let $\mathcal{T}(\mathbb{C}^l)$ be as in (20), and let $S, T : D_l \rightarrow \mathcal{T}(\mathbb{C}^l)$ be nonlinear maps defined by

$$\begin{aligned} S(x) &= (S_k(x))_{k=0}^{\infty}, & S_{2k}(x) &:= 0, \\ S_{2k+1}(x) &:= \operatorname{sgn}(b_{2k+1}) \sqrt{|b_{2k+1}| c_0^{2k+1}} \cdot \bar{x}^{\otimes(k)} \otimes x^{\otimes(k+1)}, \end{aligned}$$

$$\begin{aligned} T(x) &= (T_k(x))_{k=0}^{\infty}, & T_{2k}(x) &:= 0, \\ T_{2k+1}(x) &:= \sqrt{|b_{2k+1}| c_0^{2k+1}} \cdot \bar{x}^{\otimes(k)} \otimes x^{\otimes(k+1)}, \end{aligned}$$

for any $x \in D_l$. Then S and T are maps into $\mathcal{T}(\mathbb{C}^l)$ since

$$\begin{aligned}\|S(x)\|_*^2 &= \sum_{k=0}^{\infty} \|S_k(x)\|_k^2 = \sum_{k=0}^{\infty} |b_{2k+1}| c_0^{2k+1} \|x\|^{2(2k+1)} = \sum_{k=0}^{\infty} \|T_k(x)\|_k^2 \\ &= \|T(x)\|_*^2\end{aligned}$$

and, as $b_1 > 0$ and $b_{2k+1} \leq 0$ for all $k \geq 1$ by (13),

$$\sum_{k=0}^{\infty} |b_{2k+1}| c_0^{2k+1} \|x\|^{2(2k+1)} = 2b_1 c_0 \|x\|^2 - H^{-1}(c_0 \|x\|^2) < \infty.$$

As in Case I, the ‘tensor trick’ allows us to rewrite (24) as

$$H^{-1}(c_0 \langle x_i, y_j \rangle) = \langle S(x_i), T(y_j) \rangle_* \quad \text{or} \quad c_0 \langle x_i, y_j \rangle = H(\langle S(x_i), T(y_j) \rangle_*).$$

Moreover, since x_i and y_j are unit vectors in \mathbb{C}^l , we get

$$\|S(x_i)\|^2 = \sum_{k=0}^{\infty} |b_{2k+1}| c_0^{2k+1} = \varphi(c_0) = 1,$$

and similarly $\|T(y_j)\| = 1$. So we may apply Corollary 2.3 to get

$$\left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} \langle x_i, y_j \rangle \right| = \frac{1}{c_0} \left| \sum_{i=1}^m \sum_{j=1}^n M_{ij} H(\langle S(x_i), T(y_j) \rangle_*) \right| \leq \frac{1}{c_0},$$

which is Haagerup’s bound since $1/c_0 = 8/\pi(x_0 + 1) = K^{\mathbb{C}}$. \square

5. Nonpositivity of b_{2k+1}

To make the proof in this article entirely self-contained, we present Haagerup’s proof of the nonpositivity of b_{2k+1} that we used earlier in (13). While the main ideas are all due to Haagerup, our small contribution here is that we avoided the use of any known results of elliptic integrals in order to stay faithful to our claim of an elementary proof, i.e., one that uses only calculus and basic complex variables. To be clear, while the functions

$$(25) \quad K(x) := \int_0^{\pi/2} (1 - x^2 \sin^2 t)^{-1/2} dt, \quad E(x) := \int_0^{\pi/2} (1 - x^2 \sin^2 t)^{1/2} dt$$

do make a brief appearance in the proof of Lemma 5.1, the reader does not need to know that they are the complete elliptic integrals of the first and second kinds respectively. Haagerup had relied liberally on properties of K and E that

require substantial effort to establish [Haa]. We will only use trivialities that follow immediately from definitions.

Our point of departure from Haagerup's proof is the following lemma about two functions h_1 and h_2 , which we will see in Lemma 5.2 arise respectively from the real and imaginary parts of the analytic extension of the real Haagerup function $h : [-1, 1] \rightarrow [-1, 1]$ to the upper half plane.

Lemma 5.1. *Let $h_1, h_2 : [1, \infty) \rightarrow \mathbb{R}$ be defined by*

$$h_1(x) := \int_0^{\pi/2} \sqrt{1 - x^{-2} \sin^2 t} dt,$$

$$h_2(x) := (1 - x^{-2}) \int_0^{\pi/2} \frac{\sin^2 t}{\sqrt{1 - (1 - x^{-2}) \sin^2 t}} dt,$$

which are clearly strictly increasing functions on $[1, \infty)$ with

$$h_1(1) = 1, \quad \lim_{x \rightarrow \infty} h_1(x) = \pi/2, \quad h_2(1) = 0, \quad \lim_{x \rightarrow \infty} h_2(x) = \infty.$$

Then

$$(26) \quad \omega_1(x) := x(h_1(x)h'_2(x) - h'_1(x)h_2(x)) = \frac{\pi}{2} \quad \text{for } x \geq 1,$$

$$(27) \quad \omega_2(x) := x(h_1(x)h'_1(x) + h_2(x)h'_2(x)) \geq 2h_1(\sqrt{2})h_2(\sqrt{2}) > \frac{\pi}{4}$$

$$\quad \text{for } 1 \leq x \leq \sqrt{2}.$$

Proof. We start by observing some properties of h'_1 and h'_2 . As

$$h'_1(x) = \frac{1}{x^3} \int_0^{\pi/2} \frac{\sin^2 t}{\sqrt{1 - x^{-2} \sin^2 t}} dt = \frac{1}{x^2} \int_0^{\pi/2} \frac{\sin^2 t}{\sqrt{x^2 - \sin^2 t}} dt,$$

h'_1 is strictly decreasing on $(1, \infty)$. As $\int_0^{\pi/2} \cos^{-1} t dt = \infty$, $\lim_{x \rightarrow 1^+} h'_1(x) = \infty$. Clearly $\lim_{x \rightarrow \infty} h'_1(x) = 0$. Furthermore, when $x > 1$, since $\sqrt{x^2 - \sin^2 t} \geq \sqrt{x^2 - 1}$, we have

$$(28) \quad 0 < h'_1(x) \leq \frac{\pi}{4x^2 \sqrt{x^2 - 1}} \quad \text{for } x > 1.$$

It is straightforward to see that the functions E and K in (25) have derivatives given by

$$(29) \quad E'(y) = \frac{1}{y}(E(y) - K(y)), \quad K'(y) = \frac{1}{y(1 - y^2)}(E(y) - (1 - y^2)K(y)).$$

Clearly, $h_2(x) = K(y) - E(y)$, where $y = y(x) = \sqrt{1 - x^{-2}}$. So by chain rule,

$$h'_2(x) = y'(x) \frac{d}{dy}(K - E)(y(x)) = \frac{1}{x} \int_0^{\pi/2} (1 - (1 - x^{-2}) \sin^2 t)^{1/2} dt.$$

Hence h'_2 is strictly decreasing on $[1, \infty)$, $h'_2(1) = \pi/2$, and $\lim_{x \rightarrow \infty} h'_2(x) = 0$.

To show (26), observe that

$$\begin{aligned} h_1(x) &= E(1/x), \quad xh'_1(x) = K(1/x) - E(1/x), \\ h_2(x) &= K(y) - E(y), \quad xh'_2(x) = E(y), \end{aligned}$$

where again $y = \sqrt{1 - x^{-2}}$. Hence

$$\begin{aligned} \omega_1(x) &= E(1/x)E(y) - [K(1/x) - E(1/x)][K(y) - E(y)] \\ &= E(1/x)K(y) + K(1/x)E(y) - K(1/x)K(y). \end{aligned}$$

Computing ω'_1 , we see from (29) that $\omega'_1 \equiv 0$. So ω_1 is a constant function. By (28), $\lim_{x \searrow 1} h'_1(x)(1 - x^{-2}) = 0$, and so $\lim_{x \searrow 1} \omega_1(x) = \pi/2$. Thus $\omega_1(x) = \pi/2$ for all $x > 1$ and we may set $\omega_1(1) = \pi/2$.

We now show (27) following Haagerup's arguments. Note that

$$\omega_2(x) = E(1/x)(K(1/x) - E(1/x)) + E(y)(K(y) - E(y)).$$

Let $g(x) := E(\sqrt{x})(K(\sqrt{x}) - E(\sqrt{x}))$. A straightforward calculation using (29) shows that

$$g''(x) = \frac{1}{2} \left[\frac{E(\sqrt{x})}{1-x} - \frac{K(\sqrt{x}) - E(\sqrt{x})}{x} \right]^2 \geq 0, \quad x \in [0, 1].$$

So g is convex on $[0, 1]$. Hence $g(1-x)$ is also convex on $[0, 1]$. Let $f(x) := g(x) + g(1-x)$. Then f is convex on $[0, 1]$ and $f'(1/2) = 0$. Therefore $f(x) \geq f(1/2) \geq 2g(1/2)$. This yields the first inequality in (27): $\omega_2(x) \geq 2h_1(\sqrt{2})h_2(\sqrt{2})$ for $x \in [1, \sqrt{2}]$.

The Taylor expansions of h_1 and h_2 may be obtained as that in (11),

$$(30) \quad h_1(x) = \frac{\pi}{2} \sum_{k=0}^{\infty} \left[\frac{(2k)!}{2^{2k}(k!)^2} \right]^2 \frac{1}{1-2k} x^{-2k},$$

$$(31) \quad h_2(x) = \frac{\pi}{2} \sum_{k=0}^{\infty} \left[\frac{(2k)!}{2^{2k}(k!)^2} \right]^2 \frac{2k}{2k-1} (1-x^{-2})^k.$$

Approximate numerical values of h_1 and h_2 at $x = \sqrt{2}$ and 4 are calculated⁵ to be:

⁵ For example, using www.wolframalpha.com, which is freely available. Such numerical calculations cannot be completely avoided – Haagerup's proof implicitly contains them as he used tabulated values of elliptic integrals.

$$(32) \quad \begin{aligned} h_1(\sqrt{2}) &\approx 1.3506438, & h_2(\sqrt{2}) &\approx 0.5034307, \\ h_1(4) &\approx 1.5459572, & h_2(4) &\approx 1.7289033. \end{aligned}$$

The second inequality in (27) then follows from $2h_1(\sqrt{2})h_2(\sqrt{2}) \approx 2 \times 1.35064 \times 0.50343 > \pi/4$. \square

In the next two lemmas and their proofs, Arg will denote principal argument.

Lemma 5.2. *Let $h : [-1, 1] \rightarrow [-1, 1]$ be the real Haagerup function as defined in Section 3. Then h can be extended to a function $h_+ : \overline{\mathbb{H}} \rightarrow \mathbb{C}$ that is continuous on the closed upper half-plane $\overline{\mathbb{H}} = \{z \in \mathbb{C} : \operatorname{Im}(z) \geq 0\}$ and analytic on the upper half-plane $\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im}(z) > 0\}$. In addition, h_+ has the following properties:*

(i) $\operatorname{Im}(h_+(z)) \geq \operatorname{Im}(h_+(|z|))$ for all $z \in \overline{\mathbb{H}} \cap \{z \in \mathbb{C} : |z| \geq 1\}$ and $h_+(z) \neq 0$ for all $z \in \overline{\mathbb{H}} \setminus \{0\}$.

(ii) For $x \in [1, \infty)$,

$$\operatorname{Re}(h_+(x)) = h_1(x), \quad \operatorname{Im}(h_+(x)) = h_2(x),$$

where h_1, h_2 are as defined in Lemma 5.1.

(iii) For all $k \in \mathbb{N}$ and all real $\alpha > 1$,

$$(33) \quad b_{2k+1} = \frac{2}{\pi(2k+1)} \int_1^\alpha \operatorname{Im}\left(\left(h_+(x)\right)^{-(2k+1)}\right) dx + r_k(\alpha)$$

where

$$(34) \quad |r_k(\alpha)| \leq \frac{\alpha}{2k+1} \left(\operatorname{Im}(h_+(\alpha))\right)^{-(2k+1)}.$$

Proof. Integrating by parts, we obtain

$$h(x) = \int_0^{\pi/2} \cos t \cdot d(\arcsin(x \sin t)) = \int_0^{\pi/2} \sin t \arcsin(x \sin t) dt, \quad x \in [-1, 1].$$

The analytic function $\sin z$ is a bijection of $[-\pi/2, \pi/2] \times [0, \infty)$ onto $\overline{\mathbb{H}}$ and it maps the line segment $\{t + ia : -\pi/2 \leq t \leq \pi/2\}$ onto the half ellipsoid $\{z \in \overline{\mathbb{H}} : |z - 1| + |z + 1| = 2 \cosh a\}$. Let \arcsin_+ be the inverse of this mapping. Then \arcsin_+ is continuous in $\overline{\mathbb{H}}$ and analytic in \mathbb{H} . In addition, we have:

$$\arcsin_+ x = \begin{cases} \arcsin x & \text{if } x \in [-1, 1], \\ \frac{\pi}{2} \operatorname{sgn} x + i \operatorname{arccosh} |x| & \text{if } x \in (-\infty, -1) \cup (1, \infty), \end{cases}$$

$$\operatorname{Im}(\arcsin_+ z) = \operatorname{arccosh}\left(\frac{1}{2}(|z - 1| + |z + 1|)\right), \quad z \in \overline{\mathbb{H}}.$$

If we define

$$h_+(z) := \int_0^{\pi/2} \sin t \arcsin_+(z \sin t) dt, \quad z \in \overline{\mathbb{H}},$$

then h_+ is a continuous extension of h to $\overline{\mathbb{H}}$ and is analytic in \mathbb{H} .

(i) Since $\operatorname{arccosh}$ is increasing on $[1, \infty)$, we have

$$\operatorname{Im}(\arcsin_+ z) = \operatorname{arccosh}\left(\frac{1}{2}(|z-1| + |z+1|)\right) \geq \begin{cases} \operatorname{arccosh}|z| & \text{if } |z| \geq 1, \\ 0 & \text{if } |z| < 1. \end{cases}$$

Therefore for $z \in \overline{\mathbb{H}} \cap \{z \in \mathbb{C} : |z| \geq 1\}$,

$$\begin{aligned} \operatorname{Im}(h_+(z)) &= \int_0^{\pi/2} \sin t \cdot \operatorname{Im}(\arcsin_+(z \sin t)) dt \\ &\geq \int_{\arcsin(1/|z|)}^{\pi/2} \sin t \operatorname{arccosh}(|z| \sin t) dt = \operatorname{Im}(h_+(|z|)). \end{aligned}$$

As $\operatorname{Im}(\arcsin_+ z) > 0$ on \mathbb{H} , we have $\operatorname{Im}(h_+(z)) > 0$ on \mathbb{H} . For $x \in [-1, 1]$, $h_+(x) = h(x)$ is zero only at $x = 0$. For $x \in (-\infty, -1) \cup (1, \infty)$,

$$\operatorname{Im}(h_+(x)) = \int_{\arcsin(1/|x|)}^{\pi/2} \sin t \operatorname{arccosh}(|x| \sin t) dt > 0.$$

Hence h_+ has no zero in $\overline{\mathbb{H}} \setminus \{0\}$.

(ii) Let $x \in (1, \infty)$. Integrating by parts followed by a change-of-variables $\sin u = x \sin t$ in the next-to-last equality gives us:

$$\begin{aligned} \operatorname{Re}(h_+(x)) &= \int_0^{\arcsin(1/x)} \sin t \arcsin(x \sin t) dt + \frac{\pi}{2} \int_{\arcsin(1/x)}^{\pi/2} \sin t dt \\ &= x \int_0^{\arcsin(1/x)} \frac{\cos^2 t}{\sqrt{1 - x^2 \sin^2 t}} dt = \int_0^{\pi/2} \sqrt{1 - x^{-2} \sin^2 u} du \\ &= h_1(x). \end{aligned}$$

A change-of-variables $\sin v = (1 - x^{-2})^{-1/2} \cos t$ in the next-to-last equality gives us:

$$\begin{aligned} \operatorname{Im}(h_+(x)) &= \int_{\arcsin(1/x)}^{\pi/2} \sin t \operatorname{arccosh}(x \sin t) dt = x \int_{\arcsin(1/x)}^{\pi/2} \frac{\cos^2 t}{\sqrt{x^2 \sin^2 t - 1}} dt \\ &= (1 - x^{-2}) \int_0^{\pi/2} \frac{\sin^2 v}{\sqrt{1 - (1 - x^{-2}) \sin^2 v}} dv = h_2(x). \end{aligned}$$

(iii) The power series (11) shows that h defines an analytic function $h(z)$ in the open unit disk that is identically equal to $h_+(z)$ on $\{z \in \mathbb{C} : |z| < 1\} \cap \overline{\mathbb{H}}$. Since $h(0) = 0$ and $h'(0) \neq 0$, we can find some $\delta_0 \in (0, 1]$ such that $h(z)$ has an analytic inverse function (12) in $\{z \in \mathbb{C} : |z| < \delta_0\}$. For $0 < \delta < \delta_0$, let C_δ be a counterclockwise orientated circle with radius δ . It follows that $h(C_\delta)$ is a simple closed curve with winding number +1. Integrating by parts with a change-of-variables, we have

$$b_{2k+1} = \frac{1}{2\pi i} \int_{h(C_\delta)} \frac{h^{-1}(z)}{z^{2k+2}} dz = \frac{1}{2\pi i} \int_{C_\delta} \frac{z}{h(z)^{2k+2}} h'(z) dz.$$

Note that $b_{2k+1} \in \mathbb{R}$ and

$$-(2k+1) \int_{C_\delta} \frac{zh'(z)}{h(z)^{2k+2}} dz + \int_{C_\delta} \frac{1}{h(z)^{2k+1}} dz = \int_{C_\delta} \frac{d}{dz} \left[\frac{z}{h(z)^{2k+1}} \right] dz = 0.$$

Then we get

$$\begin{aligned} b_{2k+1} &= \frac{1}{2\pi(2k+1)} \int_{C_\delta} h(z)^{-(2k+1)} dz \\ &= \frac{1}{2\pi(2k+1)} \int_{C_\delta} \operatorname{Im}(h(z)^{-(2k+1)}) dz \\ &= \frac{2}{\pi(2k+1)} \int_{C'_\delta} \operatorname{Im}(h(z)^{-(2k+1)}) dz \end{aligned}$$

where C'_δ is the quarter circle $\{\delta e^{i\theta} : 0 \leq \theta \leq \pi/2\}$. Since $h(z)$ identically equals $h_+(z)$ on C'_δ and $h_+(z)$ has no zeros in the set $\{z \in \mathbb{C} : \delta \leq |z| \leq \alpha, 0 \leq \operatorname{Arg} z \leq \pi/2\}$ by (i), Cauchy's integral formula yields

$$\begin{aligned} b_{2k+1} &= \frac{2}{\pi(2k+1)} \operatorname{Im} \left[\int_\delta^\alpha h_+(z)^{-(2k+1)} dz + \int_{C'_\alpha} h_+(z)^{-(2k+1)} dz \right. \\ &\quad \left. + \int_{i\alpha}^{i\delta} h_+(z)^{-(2k+1)} dz \right]. \end{aligned}$$

Moreover, since $h_+(z)$ is real on $[\delta, 1]$ and its real part vanishes on the imaginary axis, we are left with

$$\begin{aligned} b_{2k+1} &= \frac{2}{\pi(2k+1)} \int_1^\alpha \operatorname{Im}(h_+(z)^{-(2k+1)}) dz \\ &\quad + \frac{2}{\pi(2k+1)} \operatorname{Im} \left[\int_{C'_\alpha} h_+(z)^{-(2k+1)} dz \right]. \end{aligned}$$

By (i), $h_+(z) \geq \operatorname{Im}(h_+(z)) \geq \operatorname{Im}(h_+(|z|))$. Thus

$$\left| \int_{C'_\alpha} h_+(z)^{-(2k+1)} dz \right| \leq \frac{\pi\alpha}{2} (\operatorname{Im}(h_+(\alpha)))^{-(2k+1)}. \quad \square$$

The integral expression of b_{2k+1} in (33) will be an important ingredient in the proof that $b_{2k+1} \leq 0$ for $k \geq 1$. We establish some further approximations for this integral in the next and final lemma.

Lemma 5.3. *Let $\alpha = 4$ throughout.⁶ Let $\theta(x) := \operatorname{Arg}(h_+(x))$ for $x \in [1, \infty)$. Then $\theta : [1, \infty) \rightarrow [0, 2\pi]$ is strictly increasing on for $x \geq 1$, $\theta(1) = 0$, and $\lim_{x \rightarrow \infty} \theta(x) = \pi/2$. In addition, we have the following:*

(i) *Let $p := \lfloor (2k+1)\theta(\alpha)/\pi \rfloor$. Let*

$$I_r := \frac{2}{\pi(2k+1)} \int_{\theta(x)=\pi(r-1)/(2k+1)}^{\theta(x)=\pi r/(2k+1)} |h_+(x)|^{-(2k+1)} |\sin((2k+1)\theta(x))| dx$$

for $r = 1, 2, \dots, p$, and

$$J := \frac{2}{\pi(2k+1)} \int_{\theta(x)=\pi p/(2k+1)}^{\alpha} |h_+(x)|^{-(2k+1)} |\sin((2k+1)\theta(x))| dx.$$

Then

$$\frac{2}{\pi(2k+1)} \int_1^{\alpha} \operatorname{Im}(h_+(x)^{-(2k+1)}) dx = -I_1 + I_2 - \dots + (-1)^p I_p + (-1)^{p+1} J.$$

(ii) *Let $k \geq 4$. Then $p \geq 2$ and $I_1 > I_2 > \dots > I_p > J$.*

(iii) *Let $k \geq 4$ and $c = |h_+(\sqrt{2})|e^{-\theta(\sqrt{2})/2}$. Then $I_1 > 0.57c^{-(2k+1)/(2k+1)^2}$ and $I_2 < 0.85I_1$.*

Proof. Since $\theta(x) = \arctan(h_2(x)/h_1(x))$, by (26), we get

$$(35) \quad \frac{d\theta(x)}{dx} = \frac{h_1(x)h_2'(x) - h_1'(x)h_2(x)}{|h_+(x)|^2} > 0, \quad x > 1.$$

So $\theta(x)$ is strictly increasing on for $x \geq 1$. It is clear that $\theta(1) = 0$. By Lemma 5.1, $\lim_{x \rightarrow \infty} h_1(x) = \pi/2$ and $\lim_{x \rightarrow \infty} h_2(x) = +\infty$, so $\lim_{x \rightarrow \infty} \theta(x) = \pi/2$.

(i) This follows from dividing the interval of the integral $[1, \alpha]$ into $p+1$ subsets:

$$\begin{aligned} \frac{2}{\pi(2k+1)} \int_1^{\alpha} \operatorname{Im}(h_+(x)^{-(2k+1)}) dx \\ = -\frac{2}{\pi(2k+1)} \int_1^{\alpha} |h_+(x)|^{-(2k+1)} \sin((2k+1)\theta(x)) dx \\ = -I_1 + I_2 - \dots + (-1)^p I_p + (-1)^{p+1} J. \end{aligned}$$

⁶To avoid confusion, we write ‘ α ’ for the upper limit of our integrals instead of ‘4’ as the same number will also appear in an unrelated context ‘ $k \geq 4$ ’.

(ii) We write $x = x(\theta)$, $\theta \in [0, \pi/2]$, for the inverse function of $\theta = \theta(x)$. By (35), we have

$$I_r = \frac{4}{\pi^2(2k+1)} \int_{\pi(r-1)/(2k+1)}^{\pi r/(2k+1)} x(\theta) |h_+(x(\theta))|^{-2k+1} |\sin((2k+1)\theta)| d\theta,$$

$$J = \frac{4}{\pi^2(2k+1)} \int_{\pi p/(2k+1)}^{\theta(\alpha)} x(\theta) |h_+(x(\theta))|^{-2k+1} |\sin((2k+1)\theta)| d\theta.$$

By Lemma 5.1, $h_1(x)$ and $h_2(x)$ are strictly increasing function of $x \in [1, \infty)$, therefore, so is $|h_+(x)|^2 = h_1(x)^2 + h_2(x)^2$. With this, we deduce that $x|h_+(x)|^{-2k+1}$ is strictly decreasing on $[1, \alpha]$ for $k \geq 4$ as

$$\begin{aligned} & \frac{d}{dx} (x|h_+(x)|^{-2k+1}) \\ &= |h_+(x)|^{-2k+1} + \frac{(-2k+1)x}{2} |h_+(x)|^{-2k-1} \frac{d}{dx} |h_+(x)|^2 \\ &= |h_+(x)|^{-2k-1} \left(|h_+(x)|^2 - (2k-1)x(h_1(x)h_1'(x) + h_2(x)h_2'(x)) \right) \\ &\leq |h_+(x)|^{-2k-1} \left(|h_+(x)|^2 - (2k-1)\frac{\pi}{4} \right) \\ &\leq |h_+(x)|^{-2k-1} \left(|h_+(\alpha)|^2 - \frac{7\pi}{4} \right) \approx -0.1187 < 0, \end{aligned}$$

where we have used the fact that $|h_+(x)|^2$ is increasing on $[1, \alpha]$ in the next-to-last inequality and the numerical value is calculated from those of $h_1(4)$ and $h_2(4)$ in (32). Since $|\sin((2k+1)\theta)|$ is periodic with period $\pi/(2k+1)$, we obtain $I_1 > I_2 > \dots > I_p$. In addition,

$$\begin{aligned} J &= \frac{4}{\pi^2(2k+1)} \int_{\pi p/(2k+1)}^{\theta(\alpha)} x(\theta) |h_+(x(\theta))|^{-2k+1} |\sin((2k+1)\theta)| d\theta \\ &\leq \frac{4}{\pi^2(2k+1)} \int_{(p-1)\pi/(2k+1)}^{\theta(\alpha)-\pi/(2k+1)} x(\theta) |h_+(x(\theta))|^{-2k+1} |\sin((2k+1)\theta)| d\theta \\ &< I_p. \end{aligned}$$

Finally, we have $\theta(\alpha) = \arctan(h_1(\alpha)/h_2(\alpha)) \approx 0.8412 > \pi/4 = \arctan(1)$, and so $p = \lfloor (2k+1)\theta(\alpha)/\pi \rfloor \geq \lfloor 9\theta(\alpha)/\pi \rfloor = 2$ for $k \geq 4$.

(iii) Since $x(\theta) \geq 1$ for $\theta \in [0, \pi/2]$, we have

$$I_1 \geq \frac{4}{\pi^2(2k+1)} \int_0^{\pi/(2k+1)} x(\theta) |h_+(x(\theta))|^{-2k+1} |\sin((2k+1)\theta)| d\theta.$$

Recall that $\theta = \theta(x)$ and $x = x(\theta)$ are inverse functions of one another. For $\theta \in [0, \theta(\sqrt{2})]$,

$$\begin{aligned}\frac{d}{d\theta} \log |h_+(x(\theta))| &= \frac{1}{2} \frac{d}{dx} \log |h_+(x)|^2 \cdot \left(\frac{d\theta}{dx}\right)^{-1} \\ &= \frac{h_1(x)h'_1(x) + h_2(x)h'_2(x)}{h_1(x)h'_2(x) - h'_1(x)h_2(x)} = \frac{\omega_2(x)}{\omega_1(x)} > \frac{1}{2},\end{aligned}$$

for $x \in [1, \sqrt{2}]$, where we have used (26), (27), and the fact that $\theta(x)$ is strictly increasing for $x \geq 1$. Hence $\log |h_+(x(\theta))| \leq \log |h_+(\sqrt{2})| - (\theta(\sqrt{2}) - \theta)/2$ which is equivalent to

$$|h_+(x(\theta))| \leq ce^{\theta/2}, \quad \theta \in [0, \theta(\sqrt{2})]$$

where $c = |h_+(\sqrt{2})|e^{-\theta(\sqrt{2})/2} \approx 1.2059$ and $\theta(\sqrt{2}) > \pi/9$, using values of $h_1(\sqrt{2})$ and $h_2(\sqrt{2})$ in (32).

It follows that for $k \geq 4$, we have

$$\begin{aligned}I_1 &\geq \frac{4}{\pi^2(2k+1)} \int_0^{\pi/(2k+1)} (ce^{\theta/2})^{-2k+1} \sin((2k+1)\theta) d\theta \\ &= \frac{4c^{-2k+1}}{\pi^2(2k+1)^2} \int_0^{\pi} e^{-(k-1/2)\theta/(2k+1)} \sin \theta d\theta \\ &\geq \frac{4c^{-2k+1}}{\pi^2(2k+1)^2} \int_0^{\pi} e^{-\theta/2} \sin \theta d\theta \\ &= \left(\frac{2c}{\pi}\right)^2 \frac{1+e^{-\pi/2}}{1+1/4} \cdot \frac{c^{-(2k+1)}}{(2k+1)^2} > \frac{0.57c^{-(2k+1)}}{(2k+1)^2}.\end{aligned}$$

Since $\frac{d}{d\theta} \log |h_+(x(\theta))| \geq 1/2$, we get

$$|h_+(x(\theta + \pi/(2k+1)))|^{-2k+1} \leq e^{-(k-1/2)\pi/(2k+1)} |h_+(x(\theta))|^{-2k+1}$$

Moreover, since $\theta(5/\sqrt{3}) > 2\pi/9$, we know that $x(\theta) \leq 5/\sqrt{3}$ on $[0, 2\pi/9]$. Hence for $k \geq 4$, it follows from the above results that

$$\begin{aligned}I_r &= \frac{4}{\pi^2(2k+1)} \frac{5}{\sqrt{3}} \int_{\pi/(2k+1)}^{2\pi/(2k+1)} x(\theta) |h_+(x(\theta))|^{-2k+1} |\sin((2k+1)\theta)| d\theta \\ &= \frac{4}{\pi^2(2k+1)} \frac{5}{\sqrt{3}} \int_0^{\pi/(2k+1)} x(\theta) |h_+(x(\theta + \pi/(2k+1)))|^{-2k+1} \\ &\quad \sin((2k+1)\theta) d\theta \\ &\leq \frac{4}{\pi^2(2k+1)} \frac{5}{\sqrt{3}} e^{-(k-1/2)\pi/(2k+1)} \int_0^{\pi/(2k+1)} x(\theta) |h_+(x(\theta))|^{-2k+1} \\ &\quad \sin((2k+1)\theta) d\theta \\ &\leq \frac{5}{\sqrt{3}} e^{-7\pi/18} I_1 < 0.85 I_1.\end{aligned}$$

□

The fact that $x|h_+(x)|^{-2k+1}$ is strictly decreasing on $[1, 4]$ for $k \geq 4$, established in the proof of (ii) above, is a crucial observation for establishing the nonpositivity of b_{2k+1} for $k \geq 4$. Observe that since $|h_+(x)|$ is strictly increasing for $x > 1$, it is enough to show that $x|h_+(x)|^{-7}$ is strictly decreasing on $[1, 4]$, which is what we did. Note that for a fixed $k \geq 1$, $x|h_+(x)|^{-2k+1}$ is *increasing* for large enough x , as $|h_+(x)|$ behaves like $C \log x$ for $x \gg 1$.

Theorem 5.4. *Let the Taylor expansion of $h^{-1}(x)$ be as in (12). Then $b_{2k+1} \leq 0$ for $k \geq 1$.*

Proof. Let $k \geq 4$ and let I_1, I_2, \dots, I_p, J be as defined in Lemma 5.3. By (33) with $\alpha = 4$ and Lemma 5.3(i) and (ii), we have

$$\begin{aligned} -b_{2k+1} &= I_1 - I_2 + \dots + (-1)^{p-1} I_p + (-1)^p J - r_{2k+1}(5\sqrt{2}) \\ &> I_1 - I_2 - r_{2k+1}(5\sqrt{2}). \end{aligned}$$

By (34) and Lemma 5.3(iii) with $c \approx 1.2059$ (established in its proof), we get

$$I_1 - I_2 > \frac{0.0855}{(2k+1)^2} (1.206)^{-(2k+1)}, \quad |r_{2k+1}(4)| \leq \frac{4}{2k+1} (1.728)^{-(2k+1)}.$$

Since $-b_{2k+1} > I_1 - I_2 - r_{2k+1}(4)$, we get $b_{2k+1} < 0$ for $k \geq 9$. Direct computation using the Lagrange inversion formula gives us $b_3, b_5, \dots, b_{17} \leq 0$, proving nonpositivity for $k \leq 8$. \square

Acknowledgment. We thank the anonymous referee for his careful reading and many helpful suggestions that greatly improved our exposition. We would also like to thank Małgorzata Stawiska for suggesting that we send this article to *L'Enseignement Mathématique*. This work is generously supported by DARPA D15AP00109 and NSF IIS 1546413. LHL acknowledges additional support from a DARPA Director's Fellowship and the Eckhardt Faculty Fund.

References

- [AK] F. ALBIAC and N.J. KALTON, *Topics in Banach Space Theory*, volume 233 of *Graduate Texts in Mathematics*. Springer, [Cham], second edition, 2016. With a foreword by Gilles Godefory. Zbl 1094.46002 MR 2192298
- [Ble] R. C. BLEI, An elementary proof of the Grothendieck inequality. *Proc. Amer. Math. Soc.* **100** (1987), 58–60. Zbl 0637.46017 MR 0883401

[Bra] M. BRAVERMAN, K. MAKARYCHEV, Y. MAKARYCHEV, and A. NAOR, The Grothendieck constant is strictly smaller than Krivine's bound. *Forum Math. Pi* 1:e4, **42** (2013). Zbl 1320.15016 MR 3141414

[Dav1] A. M. DAVIE, Lower bound for k_g . Unpublished note, 1984.

[Dav2] —— Matrix norms related to Grothendieck's inequality. In *Banach Spaces (Columbia, Mo., 1984)*, volume 1166 of *Lecture Notes in Math.*, pages 22–26. Springer, Berlin, 1985. Zbl 0611.46069 MR 0827755

[DFS] J. DIESTEL, J. H. FOURIE, and J. SWART, *The Metric Theory of Tensor Products*. American Mathematical Society, Providence, RI, 2008. Grothendieck's résumé revisited. Zbl 1186.46004 MR 2428264

[DJT] J. DIESTEL, H. JARCHOW, and A. TONGE, *Absolutely Summing Operators*, volume 43 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1995. Zbl 0855.47016 MR 1342297

[FA] S. FRIEDLAND and M. ALIABADI, *Linear Algebra and Matrices*. SIAM, Philadelphia, PA, 2018. MR 3767232

[FLZ] S. FRIEDLAND, L.-H. LIM, and J. ZHANG, Grothendieck constant is norm of Strassen matrix multiplication tensor. arXiv:1711.04427

[Gro] A. GROTHENDIECK, Résumé de la théorie métrique des produits tensoriels topologiques. *Bol. Soc. Mat. São Paulo* **8** (1953), 1–79. Zbl 0074.32303 MR 1466414

[Haa] U. HAAGERUP, A new upper bound for the complex Grothendieck constant. *Israel J. Math.* **60** (1987), 199–224. Zbl 0646.46019 MR 0931877

[Jam] G. J. O. JAMESON, *Summing and Nuclear Norms in Banach Space Theory*, volume 8 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1987. Zbl 0634.46007 MR 0902804

[Kai] S. KAIJSER, A simple-minded proof of the Pisier–Grothendieck inequality. In *Banach Spaces, Harmonic Analysis, and Probability Theory*, volume 995 of *Lecture Notes in Math.*, pages 33–44. Springer, Berlin, 1983. Zbl 0547.46038 MR 0717227

[Kon1] H. KÖNIG, On an extremal problem originating in questions of unconditional convergence. In *Recent Progress in Multivariate Approximation (Witten-Bommerholz, 2000)*, volume 137 of *Internat. Ser. Numer. Math.*, pages 185–192. Birkhäuser, Basel, 2001. Zbl 0994.47029 MR 1877506

[Kon2] —— Some remarks on the Grothendieck inequality. *Internat. Ser. Numer. Math.* **103** (1992), 201–206. Zbl 0779.47017 MR 1213007

[Kri] J.-L. KRIVINE, Constantes de Grothendieck et fonctions de type positif sur les sphères. *Adv. in Math.* **31** (1979), 16–30. Zbl 0413.46054 MR 0521464

[Lim] L.-H. LIM, Tensors and hypermatrices. Chapter 15, 30 pp., *Handbook of Linear Algebra*, 2nd Ed., CRC Press, Boca Raton, FL, 2013.

[LP] J. LINDENSTRAUSS and A. PEŁCZYŃSKI, Absolutely summing operators in L_p -spaces and their applications. *Studia Math.* **29** (1968), 275–326. Zbl 0183.40501 MR 0231188

- [LT] J. LINDENSTRAUSS and L. TZAFRIRI, *Classical Banach Spaces. I*. Springer-Verlag, Berlin-New York, 1977. Sequence spaces, *Ergebnisse der Mathematik und ihrer Grenzgebiete*, Vol. 92. Zbl 0362.46013 MR 0500056
- [Mau] B. MAUREY, Une nouvelle démonstration d'un théorème de Grothendieck. page 7, 1973. Zbl 0262.47014 MR 0399818
- [Pis1] G. PISIER, *Factorization of linear operators and geometry of Banach spaces*, volume 60 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1986. Zbl 0588.46010 MR 0829919
- [Pis2] —— Grothendieck's theorem, past and present. *Bull. Amer. Math. Soc. (N.S.)* **49** (2012), 237–323. Zbl 1244.46006 MR 2888168
- [Rie] R.E. RIETZ, A proof of the Grothendieck inequality. *Israel J. Math.* **19** (1974), 271–276. Zbl 0321.46018 MR 0367628

(Reçu le 19 avril 2018)

Shmuel FRIEDLAND, Department of Mathematics, Statistics and Computer Science, University of Illinois, Chicago, 851 South Morgan Street, Chicago, IL 60607-7045, USA

e-mail: friedlan@uic.edu

Lek-Heng LIM, Computational and Applied Mathematics Initiative, Department of Statistics, University of Chicago, 5747 South Ellis Avenue, Chicago, IL 60637-1514, USA

e-mail: lekheng@galton.uchicago.edu

Jinjie ZHANG, Department of Statistics, University of Chicago, 5747 South Ellis Avenue, Chicago, IL 60637-1514, USA

Current address:

Department of Mathematics, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0112, USA

e-mail: jiz003@ucsd.edu

