Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	66 (2020)
Heft:	1-2
Artikel:	Right Angled Artin Groups und partial commutation, old und new
Autor:	Bartholdi, Laurent / Härer, Henrika / Schick, Thomas
DOI:	https://doi.org/10.5169/seals-880822

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Right Angled Artin Groups and partial commutation, old and new

Laurent BARTHOLDI, Henrika HÄRER and Thomas SCHICK

Abstract. We systematically treat algebraic objects with free partially commuting generators and give short and modern proofs of the various relations between them. These objects include right angled Artin groups, polynomial rings, Lie algebras, and restricted Lie algebras in partially commuting free generators. In particular, we compute the p-central and exponent-p series of all right angled Artin groups, and compute the dimensions of their subquotients. We also describe their associated Lie algebras, and relate them to the cohomology ring of the group as well as to polynomial and power series rings in partially commuting variables. We finally show how the growth series of these various objects are related to each other.

Mathematics Subject Classification (2010). Primary: 20F36; Secondary: 20F65, 20F12, 20F14, 20F40, 17B50, 16S37.

Keywords. Right-angled Artin group, CAT(0) cubical group, partially commutative group, graph group, lower central series, dimension series, *p*-central series, growth, Lie algebras, Malcev algebra.

1. Introduction

Right angled Artin groups (RAAGs) are a prominent geometric/combinatorial class of groups. Originally introduced as "partially commuting free groups", they interpolate in an interesting way between free groups and free abelian groups. Of particular interest are several additional algebraic objects which are canonically coming along and are closely related to the structure of the RAAGs, in particular (graded) Lie algebras and polynomial rings, both in free partially commuting generators. The purpose of this article is to give a complete description of many relevant properties and relations, offering modern and accessible proofs. Many of the results quoted below appear already in other sources, though the computation

of the exponent-p and lower p-central series is new; throughout this text p denotes a prime number, fixed once and for all.

1.1. The actors. Let Γ be an undirected graph, with vertex set V and edge set E (consisting of 2-element subsets of V). The right angled Artin group (RAAG) A_{Γ} associated with Γ is the group defined in terms of generators and relations as

$$A_{\Gamma} = \langle V \mid vw = wv \text{ whenever } \{v, w\} \in E \rangle.$$

The purpose of this note is to describe classical subgroup series in A_{Γ} such as the lower-central and *p*-lower-central series, and relate them to other algebraic objects defined in terms of Γ as follows.

Let k be a commutative ring. We define unital associative k-algebras

$$R_{\Gamma} = \langle V \mid vw = wv \text{ whenever } \{v, w\} \in E \rangle,$$

$$S_{\Gamma} = \langle V \mid v^2 = 0, vw = -wv \forall v, w \in V, \text{ and } vw = 0 \text{ whenever } \{v, w\} \notin E \rangle.$$

Note that R_{Γ} is the familiar algebra of polynomials in partially commuting variables, and similarly S_{Γ} can be considered as an exterior algebra in partially commuting variables.

Observe that R_{Γ} and S_{Γ} are *graded* algebras with $\deg(v) = 1$ for all $v \in V$. Therefore, they admit a natural topology, in which basic eighbourhoods of 0 (say in R_{Γ}) are spans of the set of all monomials of degree $\geq n$. We define

 $\overline{R_{\Gamma}}$ = the completion of R_{Γ} in this topology.

Just as R_{Γ} is a non-commutative polynomial algebra, $\overline{R_{\Gamma}}$ is an algebra of power series in partially commuting variables.

We also define a Lie algebra over k,

$$L_{\Gamma} = \langle V \mid [v, w] = 0$$
 whenever $\{v, w\} \in E \rangle$,

and, if k is an algebra over \mathbb{F}_p , a *restricted* Lie algebra (see Section 2 for a review of restricted Lie algebras)

$$L_{\Gamma,p} = \langle V \mid [v,w] = 0$$
 whenever $\{v,w\} \in E\rangle_p$.

Let us have a look at the extreme cases.

- (1) If Γ is the complete graph on d vertices then $A_{\Gamma} \cong \mathbb{Z}^d$, R_{Γ} is the polynomial algebra in d variables $\Bbbk[X_1, \ldots, X_d]$, S_{Γ} is the Grassmann algebra $\bigwedge^*(\Bbbk^d)$, and $L_{\Gamma} \cong \Bbbk^d$ with trivial bracket.
- (2) If Γ is the empty graph on d vertices then A_{Γ} is the free group F_d , R_{Γ} is the free associative algebra on d generators, $S_{\Gamma} \cong \Bbbk \cdot 1 \oplus \Bbbk^d$ with trivial multiplication except $1 \cdot x = x$, and L_{Γ} is the free Lie algebra on d generators; for more details see Section 1.4.

1.2. Subgroup series. Let *G* be any discrete group, and let $\rho: G \to R^{\times}$ be a representation of *G* in an associative augmented \Bbbk -algebra *R* with augmentation ideal ϖ (namely, an algebra equipped with an epimorphism to \Bbbk with kernel ϖ). With this representation is associated a natural sequence of subgroups, called *generalized dimension subgroups*,

$$\delta_{n,\rho} := \rho^{-1}(1 + \varpi^n) = \ker(G \to (R/\varpi^n)^{\times}).$$

In case $R = \Bbbk G$ and ρ is the regular representation, we write $\delta_{n,\Bbbk G}$ for $\delta_{n,\rho}$. In addition, there are classical subgroup series, defined intrinsically within G:

- The lower central series (γ_n) given by $\gamma_1 = G$ and $\gamma_n = [\gamma_{n-1}, G]$;
- the rational lower central series $\gamma_{n,0} = \{g \in G \mid g^k \in \gamma_n \text{ for some } k \neq 0\};$
- for a prime *p* fixed throughout the discussion, the *exponent-p* central series $\lambda_{n,p}$ given by $\lambda_{1,p} = G$ and $\lambda_{n,p} = [\lambda_{n-1,p}, G]\lambda_{n-1,p}^p$, or more directly $\lambda_{n,p} = \prod_{m+i \ge n} \gamma_m^{p^i}$;
- again for a prime *p* fixed throughout the discussion, the *Brauer–Jennings–Lazard–Zassenhaus* series [Zas, Jen1, Laz], also called *p*-dimension or *p*-central series, given by $\gamma_{1,p} = G$ and $\gamma_{n,p} = [\gamma_{n-1,p}, G]\gamma^p_{\lceil n/p \rceil, p}$, or more directly $\gamma_{n,p} = \prod_{mp^i \ge n} \gamma^{p^i}_m$.

All these series are *central*, meaning that γ_n/γ_{n+1} belongs to the center of G/γ_{n+1} , etc. We moreover have $[\gamma_m, \gamma_n] \subseteq \gamma_{m+n}$, etc. A classical consequence [MKS, Section 5.3] is that $\bigoplus_{n\geq 1} \gamma_n/\gamma_{n+1}$, etc., are graded Lie algebras over \mathbb{Z} . The addition is induced by the group multiplication and the Lie bracket is induced by the commutator.

The groups $\gamma_{n,0}$ enjoys the extra property that $\gamma_{n,0}/\gamma_{n+1,0}$ is torsion-free (and it is the fastest descending central series with this property), so $\bigoplus_{n\geq 1} \gamma_{n,0}/\gamma_{n+1,0}$ is \mathbb{Z} -free. In particular, if γ_n/γ_{n+1} is torsion free for each n, then $\gamma_{n,0} = \gamma_n$ for each n.

We have $\lambda_{n,p}^p \subseteq \lambda_{n+1,p}$ so $\bigoplus_{n \ge 1} \lambda_{n,p}/\lambda_{n+1,p}$ is an elementary abelian *p*-group. Similarly, $\gamma_{n,p}^p \subseteq \gamma_{np,p}$. Furthermore, these series are fastest descending under these requirements. It is now classical [Zas] that $\bigoplus_{n \ge 1} \gamma_{n,p}/\gamma_{n+1,p}$ is a restricted Lie algebra over \mathbb{F}_p . The additional, "*p*-power" operation as part of the *restricted* Lie algebra structure is induced by the *p*-power operation in the group.

Classical results identify $\delta_{n,\Bbbk G}$ with some of the above series in case \Bbbk is a field: we have $\delta_{n,\Bbbk G} = \gamma_{n,p}$ where $p \ge 0$ is the characteristic of \Bbbk [Hal2, Jen1, Jen2]. However, for general G, the identification of $\delta_{n,\mathbb{Z}G}$ is a fundamental open problem of group theory.

1.3. Results. We consider the series defined above for the group A_{Γ} . The main purpose of this text is to exhibit numerous relations between these algebraic objects; detailed definitions and proofs will be given in subsequent sections. The main tool is an extension to A_{Γ} of Magnus's work on the free group [MKS, §5], embedding it into the units of the free non-commuting power series ring. This extension seems first considered in [Dro].

Definition 1.1 (Augmentation ideals). Recall that a commutative ring k is fixed. Denote by $\overline{\omega}$ the augmentation ideal of R_{Γ} (i.e. the ideal of polynomials in partially commuting variables with zero constant term), and by $\overline{\omega}(A_{\Gamma})$ the augmentation ideal of $\Bbbk A_{\Gamma}$. We also denote by $\overline{\omega}_p$ the ideal $\overline{\omega} + pR_{\Gamma}$.

Theorem 1.2 (Augmentation powers). For all *n* we have

$$\overline{\omega}(A_{\Gamma})^n/\overline{\omega}(A_{\Gamma})^{n+1} \cong \overline{\omega}^n/\overline{\omega}^{n+1}.$$

We remind the reader that Koszul algebras are a particular kind of associative algebras (see [Pri] or Section 4) for which a "small" projective resolution may easily be computed. Moreover, there is the important concept of Koszul duality. We obtain the following results, which for $\mathbb{k} = \mathbb{Q}$ already appear in [PS2]. Recall that a *clique* in a graph is a subset of vertices in which all vertices are connected to each other.

Theorem 1.3 (Group cohomology). Let S^1 be the circle with base point *. The following subspace of the torus $(S^1)^V$ is a classifying space for A_{Γ} :

(1.1)
$$X_{\Gamma} = \bigcup_{C \subseteq V \ a \ clique} (S^1)^C \times \{*\}^{V \setminus C}.$$

We have $H^*(A_{\Gamma}; \Bbbk) = S_{\Gamma}$.

The rings R_{Γ} and S_{Γ} are Koszul algebras, and Koszul duals to each other: $(S_{\Gamma})^! = R_{\Gamma}.$

Theorem 1.4 i (Central series and dimension subgroups). In the group A_{Γ} we have

(1.2)
$$\gamma_n = \gamma_{n,0} \quad and \quad \bigcap_{n \in \mathbb{Z}} \gamma_{n,0} = \bigcap_{n \in \mathbb{Z}} \gamma_{n,p} = \{1\}.$$

In particular, A_{Γ} is finitely generated and residually torsion-free nilpotent, so (by [Gru]*Theorem 2.1) A_{Γ} is also a residually finite p-group for every p.

Theorem 1.4ii (Central series and dimension subgroups). For arbitrary \Bbbk there is a faithful representation

(1.3)
$$\mu: A_{\Gamma} \to \overline{R_{\Gamma}}^{\times}; \quad v \mapsto 1 + v \quad for \ v \in V.$$

The corresponding generalized dimension subgroups satisfy

(1.4)
$$\delta_{n,\mu} = \begin{cases} \gamma_{n,0} & \text{if } \Bbbk \text{ has characteristic } 0, \\ \gamma_{n,p} & \text{if } \Bbbk \text{ has characteristic } p. \end{cases}$$

Together with Theorem 1.2 we obtain an isomorphism of filtered associative \Bbbk -algebras (but not Hopf algebras; see Theorem 1.7 below!)

$$\overline{\Bbbk A_{\Gamma}} := \lim(\Bbbk A_{\Gamma}/\varpi(A_{\Gamma})^n) \xrightarrow{\cong} \lim(R_{\Gamma}/\varpi^n) = \overline{R_{\Gamma}}.$$

In particular, the classical dimension subgroups coincide:

(1.5)
$$\delta_{n,\Bbbk A_{\Gamma}} = \delta_{n,\mu}$$

The Lie algebras L_{Γ} and $L_{\Gamma,p}$ are tightly connected to their associative counterparts:

Theorem 1.5i (Lie algebras). The algebra R_{Γ} is a Hopf algebra. If the ring \Bbbk is a \mathbb{Z} -free module then we have

(1.6)
$$L_{\Gamma} \cong Primitives(R_{\Gamma}) \quad and \quad R_{\Gamma} \cong U(L_{\Gamma}),$$

the universal enveloping algebra of L_{Γ} , while if \Bbbk is an \mathbb{F}_p -algebra then

(1.7)
$$L_{\Gamma,p} \cong Primitives(R_{\Gamma}) \quad and \quad R_{\Gamma} \cong U_p(L_{\Gamma,p}),$$

the *p*-universal enveloping algebra of $L_{\Gamma,p}$. The Lie algebra cohomology of L_{Γ} is

$$H^*(L_{\Gamma}; \Bbbk) \cong S_{\Gamma}.$$

All the above isomorphisms are natural, in the sense that they are induced by the identity map $V \rightarrow V$, and therefore compatible with homomorphisms induced by a map of graphs $V \rightarrow V'$.

The Lie algebra associated with the lower central series was already determined in [DK2] as L_{Γ} . We extend this result as follows:

Theorem 1.5ii (Lie algebras). For any ring \Bbbk , we have, as Lie algebras,

$$L_{\Gamma} \cong \bigoplus_{n \ge 1} (\gamma_n / \gamma_{n+1}) \otimes_{\mathbb{Z}} \mathbb{k}.$$

If \Bbbk has characteristic p then as restricted Lie algebras

$$L_{\Gamma,p} \cong \bigoplus_{n \ge 1} (\gamma_{n,p} / \gamma_{n+1,p}) \otimes_{\mathbb{Z}} \mathbb{k}.$$

If k has characteristic $p \ge 3$ then with $k[\pi]$ the polynomial ring in one degree-1 variable π

under that isomorphism, multiplication by π corresponds to the map induced by $\lambda_{n,p} \ni g \mapsto g^p \in \lambda_{n+1,p}$.

All the above isomorphisms are natural, in the sense that they are induced by the identity map $V \rightarrow V$, and therefore compatible with homomorphisms induced by a map of graphs $V \rightarrow V'$.

For a graded algebra $R = \bigoplus_{n \ge 0} R_n$ over k such that each R_n is a finitely generated free k-module, recall that its *Poincaré series* is the power series

$$\Phi_R(t) = \sum_{n \ge 0} \operatorname{rank}(R_n) t^n.$$

For a group $G = \langle X \rangle$, its growth series is $\Phi_G(t) = \sum_{g \in G} t^{|g|}$, with |g| denoting the word length of $g \in G$ (word length and growth series depend on the fixed generating set X). The first two claims of the following result appear in [DK1]:

Theorem 1.6 (Poincaré and growth series). The Poincaré series of S_{Γ} is

$$\Phi_{S_{\Gamma}}(t) = \sum_{n=0}^{|V|} c_n(\Gamma) t^n,$$

where $c_n(\Gamma)$ denotes the number of cliques of size n (i.e., complete subgraphs of Γ with n vertices).

The Poincaré series of R_{Γ} and S_{Γ} are connected by the relation

$$\Phi_{R_{\Gamma}}(t) \cdot \Phi_{S_{\Gamma}}(-t) = 1,$$

and the growth series of A_{Γ} is

$$\Phi_{A_{\Gamma}}(t) = \Phi_{R_{\Gamma}}\left(\frac{2t}{1+t}\right).$$

In our next result, originally appearing in [KM, Theorem 16.10], we determine the Malcev completion of A_{Γ} . We refer to [Mal, Qui2] and the more recent [PS1] for a review of this construction. **Theorem 1.7** (Malcev completions). Assume $\mathbb{k} = \mathbb{Q}$. There is then an isomorphism $\mu_{\exp}: \overline{R_{\Gamma}} \to \overline{\mathbb{Q}A_{\Gamma}}$ of filtered, complete Hopf algebras; via this isomorphism, $\overline{L_{\Gamma}}$ is the Malcev Lie algebra of A_{Γ} , and the Malcev completion of A_{Γ} is given on generators by

$$A_{\Gamma} \to \exp(L_{\Gamma}) \subset \overline{R_{\Gamma}}$$
 via the classical power series $v \mapsto \sum_{n \ge 0} \frac{v^n}{n!} \quad \forall v \in V.$

We also show the following related result on formality in the sense of rational homotopy theory; see Section 8 for a review of the notion.

Theorem 1.8 (Formality). The classifying space X_{Γ} of A_{Γ} of (1.1) is formal.

1.4. Examples and illustrations. Let us consider, as sketched in the Introduction, the two extreme cases of graphs Γ , the complete and empty graphs.

If Γ is the complete graph on V, then A_{Γ} is free Abelian with basis V, and R_{Γ} is a usual polynomial algebra in variables V. The standard Koszul complex is given by the exterior algebra $S_{\Gamma} = \bigwedge^{*}(V)$, and coincides with the cohomology ring of $A_{\Gamma} = \mathbb{Z}V$. The classifying space X_{Γ} is the usual torus $(S^{1})^{V}$. The exponent-p central series satisfies $\lambda_{n,p} = p^{n-1}\mathbb{Z}V$, and the p-dimension series satisfies $\gamma_{n,p} = p^{i}\mathbb{Z}V$ whenever $p^{i-1} < n \leq p^{i}$. The growth series are readily computed as

$$\Phi_{A_{\Gamma}}(t) = \left(\frac{1+t}{1-t}\right)^{|V|}, \quad \Phi_{R_{\Gamma}}(t) = \left(\frac{1}{1-t}\right)^{|V|}, \quad \Phi_{S_{\Gamma}}(t) = \left(1+t\right)^{|V|}.$$

If, on the other hand, Γ is the empty graph on V, then A_{Γ} is free with basis V, and R_{Γ} is a polynomial algebra in non-commuting variables V. The algebra S_{Γ} is reduced to $\mathbb{k} \oplus \mathbb{k} V$ with $V^2 = 0$, and coincides with the cohomology ring of A_{Γ} . The classifying space X_{Γ} is a wedge of |V| circles. The Lie algebras L_{Γ} and the restricted Lie algebra $L_{\Gamma,p}$ are free. The growth series are readily computed as

$$\Phi_{A_{\Gamma}}(t) = \frac{1+t}{1-(2|V|-1)t}, \quad \Phi_{R_{\Gamma}}(t) = \frac{1}{1-|V|t}, \quad \Phi_{S_{\Gamma}}(t) = 1+|V|t.$$

These results can be seen as special cases of the following constructions. If Γ is the disjoint union of two graphs $\Gamma_1 \sqcup \Gamma_2$, then $A_{\Gamma} = A_{\Gamma_1} * A_{\Gamma_2}$ is a free product of groups, and similarly L_{Γ} and R_{Γ} are free products in their respective categories, and $S_{\Gamma} = S_{\Gamma_1} \oplus S_{\Gamma_2}/(1 \oplus 0 - 0 \oplus 1)$. The space X_{Γ} is the wedge (one-point union) of X_{Γ_1} and X_{Γ_2} , and the growth series of $A_{\Gamma}, R_{\Gamma}, S_{\Gamma}$ may be easily be deduced from those of $A_{\Gamma_1}, A_{\Gamma_2}$, etc.:

L. BARTHOLDI, H. HÄRER and T. SCHICK

$$1 - \frac{1}{\Phi_{A_{\Gamma}}} = \left(1 - \frac{1}{\Phi_{A_{\Gamma_1}}}\right) + \left(1 - \frac{1}{\Phi_{A_{\Gamma_2}}}\right),$$

$$1 - \frac{1}{\Phi_{R_{\Gamma}}} = \left(1 - \frac{1}{\Phi_{R_{\Gamma_1}}}\right) + \left(1 - \frac{1}{\Phi_{R_{\Gamma_2}}}\right),$$

$$1 - \Phi_{S_{\Gamma}} = \left(1 - \Phi_{S_{\Gamma_1}}\right) + \left(1 - \Phi_{A_{\Gamma_2}}\right).$$

If Γ is the join of two graphs Γ_1 and Γ_2 , namely the graph obtained from $\Gamma_1 \sqcup \Gamma_2$ by adding all edges between Γ_1 and Γ_2 , then $A_{\Gamma} = A_{\Gamma_1} \times A_{\Gamma_2}$ is a direct product, and similarly $L_{\Gamma} = L_{\Gamma_1} \times L_{\Gamma_2}$ and $R_{\Gamma} = R_{\Gamma_1} \otimes R_{\Gamma_2}$, while S_{Γ} is $S_{\Gamma_1} \otimes S_{\Gamma_2}$ qua k-module, with product $(a \otimes b)(c \otimes d) = (-1)^{\deg(b)\deg(c)}(ac \otimes bd)$. The classifying space is $X_{\Gamma} = X_{\Gamma_1} \times X_{\Gamma_2}$, and the growth series $\Phi_{A_{\Gamma}}$, $\Phi_{R_{\Gamma}}$ and $\Phi_{S_{\Gamma}}$ behave multiplicatively:

$$\Phi_{A_{\Gamma}} = \Phi_{A_{\Gamma_1}} \cdot \Phi_{A_{\Gamma_2}}, \qquad \Phi_{R_{\Gamma}} = \Phi_{R_{\Gamma_1}} \cdot \Phi_{R_{\Gamma_2}}, \qquad \Phi_{S_{\Gamma}} = \Phi_{S_{\Gamma_1}} \cdot \Phi_{S_{\Gamma_2}}.$$

Finally, all the objects constructed are functorial, in the sense that graph morphisms induce maps between the corresponding objects: if Γ , Γ' are graphs and $f: \Gamma \to \Gamma'$ is a map from the vertex set of Γ to that of Γ' sending edges of Γ to edges of Γ' , then there is an induced group homomorphism $f_*: A_{\Gamma} \to A_{\Gamma'}$, ring homomorphism $R_{\Gamma} \to R_{\Gamma'}$ and $S_{\Gamma'} \to S_{\Gamma}$ (note the direction!), etc. Furthermore, if f is injective and full (meaning that $\{f(v), f(w)\}$ is an edge in Γ' precisely when $\{v, w\}$ is an edge in Γ) then the corresponding group and ring homomorphisms are injective.

1.5. Structure of the article. The article introduces and relies on quite a number of different concepts (Hopf algebras, the Magnus map, ...). These are introduced one after the other in the following sections. In particular, Section 2 collects some basic information about (restricted) Lie algebras and Hopf algebras which we use as technical tools; we prove the first part of Theorem 1.5i in it.

Section 3 introduces the Magnus map, which embeds the group A_{Γ} into the units of the partially commuting power series ring $\overline{R_{\Gamma}}$, proving (1.3). We show that this map is compatible with the central series filtrations (and dimension series filtrations). The explicit knowledge of the structure of the power series ring can be transferred to A_{Γ} to give the desired information about the latter. We also prove Theorem 1.2 in it.

We next introduce cohomological notions in Section 4, and use them to prove Theorem 1.3.

We study central series in more depth in Section 5, and prove there further parts of Theorem 1.4ii. The remaining statements about the central series require more knowledge on the Lie algebras L_{Γ} , which we describe in Section 6; we prove Theorems 1.4 i and 1.5 ii there. We also complete there the proof of Theorem 1.5 i that pertains to Lie algebra cohomology.

Finally Section 7 proves Theorem 1.6 and Section 8 proves Theorem 1.7. We apologize to the reader if the proofs are not given in strictly linear order; we found it preferable to prove individual statements of the main results where the appropriate tools were introduced.

2. Lie and Hopf algebras

We first recall from [Jac] that a restricted Lie algebra over \Bbbk , in characteristic p, is a Lie algebra equipped with an extra operation, written $x \mapsto x^{[p]}$, called the *p*-mapping and subject to the following axioms, where we use the standard multi-commutator convention [x, y, z] = [x, [y, z]], etc. For all x, y in the Lie algebra and $\alpha \in \Bbbk$,

$$[y, x^{[p]}] = [y, x, \dots, x] \quad (p \text{ factors } 'x'); \qquad (\alpha x)^{[p]} = \alpha^p x^{[p]};$$
$$(x + y)^{[p]} = x^{[p]} + y^{[p]} + \sum_{i=1}^{p-1} s_i(x, y)$$

for the Lie expressions $s_i(X, Y)$ defined by

$$\frac{d}{dt}[X, tX + Y, \dots, tX + Y] = \sum s_i(X, Y)t^i \quad \text{with } p - 1 \text{ factors } tX + Y'.$$

For example, if p = 2 then $s_1(X, Y) = [X, Y]$, and if p = 3 then $s_1(X, Y) = [Y, X, Y]$ and $s_2(X, Y) = [X, Y, X]$.

We adopt the convention that, in characteristic 0, every Lie algebra is restricted with trivial p-mapping. This way, from now on we can uniformly work with restricted Lie algebras.

Recall that every restricted Lie algebra L has a *restricted universal enveloping* algebra, a unital associative algebra $U_p(L)$ equipped with a map of restricted Lie algebras $L \to U_p(L)$, universal with respect to this property. The Lie bracket in L is identified with the commutator [x, y] = xy - yx, and the p-mapping in L is identified with the p-power operation in $U_p(L)$. The map $L \to U_p(L)$ is injective.

Recall next that a Hopf algebra is an associative algebra R equipped with additional structure, in particular an *augmentation* $\varepsilon \colon R \to \Bbbk$ and a *coproduct* $\Delta \colon R \to R \otimes R$ which are algebra homomorphisms, and an *antipode* $S \colon R \to R$ which is an algebra antihomomorphism, subject to some axioms that we shall not need; see [Swe].

We will use the following classical facts, see [Ser, Theorem III.5.4 and Exercise 2].

Proposition 2.1. The (restricted) universal enveloping algebra U(L), respectively $U_p(L)$, is a Hopf algebra. The augmentation, coproduct and antipode are given by

$$\varepsilon \colon U(L) \to U(L)/\langle L \rangle \xrightarrow{\cong} \mathbb{k}; \qquad \Delta(x) = x \otimes 1 + 1 \otimes x; \qquad S(x) = -x \ \forall x \in L.$$

In a Hopf algebra H, call $x \in H$ PRIMITIVE if $\Delta(x) = x \otimes 1 + 1 \otimes x$; the primitive elements of H form a Lie subalgebra P of H. If the ring \Bbbk is a \mathbb{Z} -free module, then the primitive elements in U(L) coincide with L, while if L is restricted and \Bbbk is p-torsion then the primitive elements in $U_p(L)$ coincide with L;

If a (restricted) Lie algebra over \Bbbk is given by a (restricted) Lie algebra presentation, then by the universal property the same presentation, now as a presentation of algebras over \Bbbk , defines its (restricted) universal enveloping algebra. In particular, R_{Γ} is the (restricted) enveloping algebra of L_{Γ} or $L_{\Gamma,p}$, respectively.

Proof of Theorem 1.5i. As a universal enveloping algebra, $R_{\Gamma} = U(L_{\Gamma})$ is by Proposition 2.1 a Hopf algebra (this also appears in [Sch]), and its Lie subalgebra of primitive elements P is equal to L_{Γ} or $L_{\Gamma,p}$, when considered as subset of R_{Γ} in the obvious way.

We note for later use the following standard constructions, see also [Quil].

Proposition 2.2. If G is a group then the group ring $\Bbbk G$ is a Hopf algebra with augmentation, coproduct and antipode given as follows:

 $\varepsilon \colon \Bbbk G \to \Bbbk \text{ induced by the map } G \to \{1\}; \quad \Delta(g) = g \otimes g; \quad S(g) = g^{-1} \; \forall g \in G.$

Furthermore, if H is a Hopf algebra and ϖ denotes its augmentation ideal $\ker(\varepsilon)$, then $\bigoplus_{n>0} \varpi^n / \varpi^{n+1}$ is naturally a graded Hopf algebra.

3. The Magnus map

3.1. Filtrations and gradings. We first recall that, since the relations of R_{Γ} and S_{Γ} are homogeneous, these rings are naturally graded by setting deg(v) = 1 for all $v \in V$. We view R_{Γ} as a ring of polynomials in partially commuting variables $v \in V$.

Let us consider the augmentation ideal $\varpi = \langle V \rangle$ in R_{Γ} . It consists of all polynomials without constant term. Note that ϖ^n then consists of all polynomials with no terms of degree $\langle n \rangle$. We define a topology on R_{Γ} by declaring the sets

 ϖ^n to form a basis of neighbourhoods of 0, and let $\overline{R_{\Gamma}}$ be the completion of R_{Γ} in this topology. We thus have

$$R_{\Gamma} \cong \bigoplus_{n \ge 0} \overline{\varpi}^n / \overline{\varpi}^{n+1}, \qquad \overline{R_{\Gamma}} \cong \prod_{n \ge 0} \overline{\varpi}^n / \overline{\varpi}^{n+1}.$$

We write $\overline{\varpi}$ for the closure of $\overline{\varpi}$ in $\overline{R_{\Gamma}}$. It consists of all power series with vanishing constant term, and similarly $\overline{\varpi}^n$ consists of the power series with no terms of degree < n.

For comparison, consider the group ring $\Bbbk A_{\Gamma}$, and let $\varpi(A_{\Gamma})$ denote the augmentation ideal of $\Bbbk A_{\Gamma}$; it is the ideal

$$\langle g-1 \mid g \in A_{\Gamma} \rangle = \langle v-1 \mid v \in V \rangle.$$

We topologize $\mathbb{k}A_{\Gamma}$ by declaring the $\varpi(A_{\Gamma})^n$ to form a basis of neighbourhoods of the identity, and let $\overline{\mathbb{k}A_{\Gamma}}$ denote the corresponding completion. Moreover, let $\operatorname{gr}(\mathbb{k}A_{\Gamma}) := \bigoplus_{n \ge 0} \varpi(A_{\Gamma})^n / \varpi(A_{\Gamma})^{n+1}$ be the associated graded algebra. We isolate the main ingredient of Theorem 1.2:

Lemma 3.1. We have $R_{\Gamma} \cong \operatorname{gr}(\Bbbk A_{\Gamma})$ as graded algebras via the natural map

$$\alpha \colon R_{\Gamma} \to \operatorname{gr}(\Bbbk A_{\Gamma}); \quad v_j \mapsto [(v_j - 1)] \text{ for } v_j \in V.$$

Proof. The isomorphism between the degree-*n* subspace of R_{Γ} and $\varpi(A_{\Gamma})^n/\varpi(A_{\Gamma})^{n+1}$ can be proven by elementary considerations, since $\varpi(A_{\Gamma})^n/\varpi(A_{\Gamma})^{n+1}$ is generated by expressions $(v_1 - 1) \cdots (v_n - 1)$.

However, here is a somewhat more elegant shortcut: As we noted in Propositions 2.1 and 2.2, $\mathbb{k}A_{\Gamma}$, $gr(\mathbb{k}A_{\Gamma})$, and R_{Γ} are all cocommutative Hopf algebras, with coproduct induced respectively by $\Delta(g) = g \otimes g$, by $\Delta([g-1]) = [(g-1) \otimes 1 + 1 \otimes (g-1)]$ for $g \in A_{\Gamma}$ and by $\Delta(v) = v \otimes 1 + 1 \otimes v$ for $v \in V$.

The map $\alpha \colon R_{\Gamma} \to \operatorname{gr}(\Bbbk A_{\Gamma})$ is a well defined map of unital graded algebras because the defining commutation property for the v_j in R_{Γ} is satisfied for their images, and all these elements are of degree 1. Moreover, we see that this map is a map of Hopf algebras.

Finally, α is an isomorphism when restricted to the degree 1 subspaces, since

$$\varpi/\varpi^2 \cong \Bbbk V \cong (A_{\Gamma}/[A_{\Gamma}, A_{\Gamma}]) \otimes \Bbbk \cong \varpi(A_{\Gamma})/\varpi(A_{\Gamma})^2.$$

Here, the last isomorphism is the standard isomorphism of the first group homology $H^1(A_{\Gamma}; \mathbb{k}) = A_{\Gamma}/[A_{\Gamma}, A_{\Gamma}] \otimes \mathbb{k}$ as $\varpi(A_{\Gamma})/\varpi(A_{\Gamma})^2$. We conclude by [MM, Theorems 5.18 and 6.11] that α is an isomorphism: it is a map between cocommutative Hopf algebras both generated as algebras in degree 1 and the map is an isomorphism in degree 1. This shortcut already appears in [Qui1]. \Box **Remark 3.2.** An alternative proof of Lemma 3.1 was kindly suggested to us by Jacques Darné: there are natural maps

$$A_{\Gamma} \supset V \to \overline{R_{\Gamma}}$$
 and $R_{\Gamma} \supset V \to \overline{\Bbbk A_{\Gamma}}$,

which induce isomorphisms $\overline{\Bbbk A_{\Gamma}} \leftrightarrow \overline{R_{\Gamma}}$ by universal properties. Since gr $R_{\Gamma} \cong R_{\Gamma}$, the result (and the last statement of Proposition 3.6) follow.

Proof of Theorem 1.2. Lemma 3.1 gives an isomorphism between the degree n part of R_{Γ} and $\varpi(A_{\Gamma})^n/\varpi(A_{\Gamma})^{n+1}$. Since R_{Γ} is graded and not only filtered, its degree-n part is ϖ^n/ϖ^{n+1} , so we get the desired isomorphism $\varpi^n/\varpi^{n+1} \cong \varpi(A_{\Gamma})^n/\varpi(A_{\Gamma})^{n+1}$ for each $n \in \mathbb{N}$.

3.2. The Magnus map. We turn to the fundamental tool we use in relating the group A_{Γ} with the algebra R_{Γ} : it is the "Magnus map"

(3.1)
$$\mu \colon \begin{cases} A_{\Gamma} \to 1 + \overline{\varpi} \subseteq \overline{R_{\Gamma}}^{\times} \subseteq \overline{R_{\Gamma}}, \\ v \mapsto 1 + v \text{ for } v \in V. \end{cases}$$

Here, $\overline{R_{\Gamma}}^{\times}$ is the group of multiplicative units of $\overline{R_{\Gamma}}$. We have to map to the completion because we have to map v^{-1} to $\mu(v)^{-1} = 1 - v + v^2 - v^3 + \cdots$ which is an infinite sum. It is immediate that the commutation relations between the $v \in V$ defining A_{Γ} also hold between the $\mu(v)$, therefore μ is well defined.

It is easy to describe quite explicitly a basis of the polynomial ring in partially commuting variables R_{Γ} . This comes hand-in-hand with a kind of normal form for elements of A_{Γ} :

Definition 3.3. A word $v_1^{e_1} \cdots v_n^{e_n}$ with $v_i \in V$ and $e^i \in \mathbb{Z}$ is called Γ -reduced if the number *n* of factors $v_i^{e_i}$ cannot be reduced by application of any sequence of moves which are either

- (M1) remove v_i^0 ,
- (M2) replace the piece $v_i^{e_i}v_{i+1}^{e_{i+1}}$ by $v_i^{e_i+e_{i+1}}$ (if $v_i = v_{i+1}$), or
- (M3) replace $v_i^{e_i} v_{i+1}^{e_{i+1}}$ by $v_{i+1}^{e_{i+1}} v_i^{e_i}$ (if $\{v_i, v_{i+1}\} \in E$).

Note that none of these moves increases the number of factors.

As we shall see in Proposition 3.5, the set of (M3)-equivalence classes of Γ -reduced words is a basis of R_{Γ} ; more precisely, any set of representatives of (M3)-equivalence classes of reduced words of length *n* forms a basis of the degree-*n* component of R_{Γ} . Indeed R_{Γ} is a monoid ring, so a family of words forms a linearly independent set if the words represent distinct elements of the monoid, or *a fortiori* of the group A_{Γ} .

In case $\mathbb{k} = \mathbb{Z}$, or more generally if \mathbb{k} has characteristic 0, it is known that the Magnus map μ is injective, see [Wad, Corollary 4.8]. We adapt this argument to \mathbb{k} of non-zero characteristic, arriving at some of the original results of this note:

Lemma 3.4. Let \Bbbk be a ring of characteristic p > 0.

Consider $g \in A_{\Gamma}$. There exists a maximal $k \in \mathbb{N}$, and minimal $s_1, \ldots, s_k \in \mathbb{N}$, such that there is a Γ -reduced monomial $m = w_1^{p^{s_1}} \cdots w_k^{p^{s_k}}$ with non-zero coefficient in $\mu(g)$. This monomial is unique. Furthermore, if $v_1^{e_1} \cdots v_n^{e_n}$ is a reduced representative of g then n = k and $v_1 \cdots v_n = w_1 \cdots w_k$ and $p^{s_i}|_{e_i}$ and the coefficient of m in $\mu(g)$ is $(e_1 p^{-s_1}) \cdots (e_n p^{-s_n})$.

Proof. Consider a Γ -reduced representative $v_1^{e_1} \cdots v_n^{e_n}$ of g. By definition,

$$\mu(v_1^{e_1}\cdots v_n^{e_n}) = (1+v_1)^{e_1}\cdots (1+v_n)^{e_n}$$

which is a possibly infinite (if one of the e_i is negative) \mathbb{F}_p -linear combination of words over V. Write $e_j = p^{s_j} \ell_j$ so that p does not divide ℓ_j . Because we are in characteristic p, we have $(1 + v_j)^{e_j} = (1 + v_j^{p^{s_j}})^{\ell_j}$.

We may now apply a variant of Magnus's original argument [Mag, Satz I]: Multiplying out (using the power series for the inverse), we obtain a multiple of $v_1^{p^{s_1}} \cdots v_n^{p^{s_n}}$ precisely once, with coefficient $\ell_1 \cdots \ell_n \neq 0 \in \mathbb{F}_p$. Other terms either have fewer syllables or larger exponents. The monomial $v_1 \cdots v_n$ and all other monomials with the same number of syllables and possibly larger exponents are Γ -reduced, because any sequence of moves which would reduce one of them could be applied in the same way to the original $v_1^{e_1} \cdots v_n^{e_n}$ and would reduce its number of factors, as well. Therefore the term $v_1^{p^{s_1}} \cdots v_n^{p^{s_n}}$ indeed is uniquely determined as the Γ -reduced monomial in $\mu(g)$ with non-zero coefficient with maximal number of syllables and minimal exponents.

Since $\mu(g)$ is independent of the choice of representative of g, every other Γ -reduced representative $(v'_1)^{e'_1} \cdots (v'_{n'})^{e'_{n'}}$ must satisfy n = n' and $v_1 \cdots v_n = v'_1 \cdots v'_{n'}$.

From this (and we note it for further use) we may deduce that every element of A_{Γ} has an essentially unique reduced representative:

Proposition 3.5 ([Wad, Theorem 4.14]). If $v_1^{e_1} \cdots v_n^{e_n}$ and $w_1^{f_1} \cdots w_m^{f_m}$ are two reduced words representing the same element of A_{Γ} , then one can be obtained from the other by a finite number of applications of (M3). In particular, n = m.

Proof. We note first by Lemma 3.4 that m = n. We then proceed by induction on m. Consider the equal elements $v_2^{e_2} \cdots v_m^{e_m}$ and $v_1^{-e_1} w_1^{f_1} \cdots w_m^{f_m}$. The latter

is not Γ -reduced, again by Lemma 3.4, so there must exist $k \in \mathbb{N}$ with $w_k = v_1$ and $\{v_1, w_i\} \in E$ for all $i \leq k$. If $f_k \neq e_1$ then $w_1^{f_1} \cdots w_k^{f_k - e_1} \cdots w_m^{f_m}$ is Γ reduced, yet again contradicting Lemma 3.4, so $f_k = e_1$ and we apply induction to $v_2^{e_2} \cdots v_m^{e_m}$ and $w_1^{f_1} \cdots \widehat{w_k^{f_k}} \cdots w_m^{f_m}$, where the factor with hat is left out. \Box

We express the first claim of Theorem 1.4ii as the following

Proposition 3.6. For arbitrary \Bbbk , the Magnus map $\mu: A_{\Gamma} \to \overline{R_{\Gamma}}$ is injective.

It maps $\gamma_n(A_{\Gamma})$ into the subgroup $1 + \overline{\varpi}^n$ of $1 + \overline{\varpi} \subset \overline{R_{\Gamma}}$. We get an induced map of graded Lie algebras

$$\mu_L \colon \bigoplus_{n \ge 1} \gamma_n(A_{\Gamma}) / \gamma_{n+1}(A_{\Gamma}) \to \bigoplus_{n \ge 1} (1 + \overline{\varpi}^n) / (1 + \overline{\varpi}^{n+1}) \cong \bigoplus_{n \ge 1} \overline{\varpi}^n / \overline{\varpi}^{n+1} \subset R_{\Gamma},$$

where the Lie algebra structure of R_{Γ} is the one induced from the algebra structure.

The algebra map induced by μ on the group algebra $\mathbb{k}A_{\Gamma}$ extends continuously to an isomorphism of filtered associative \mathbb{k} -algebras

$$\overline{\mu} \colon \overline{\Bbbk A_{\Gamma}} \xrightarrow{\cong} \overline{R_{\Gamma}}.$$

In particular,

 $\mathbb{k}A_{\Gamma}/\varpi^{n}(A_{\Gamma}) \cong \overline{\mathbb{k}A_{\Gamma}}/\overline{\varpi^{n}(A_{\Gamma})} \cong \overline{R_{\Gamma}}/\overline{\varpi^{n}} = R_{\Gamma}/\overline{\varpi^{n}} \cong \operatorname{gr}(\mathbb{k}A_{\Gamma})/\operatorname{gr}(\mathbb{k}A_{\Gamma})_{\geq n},$

using Lemma 3.1 for the last isomorphism. As \Bbbk -modules, these are of course also isomorphic to $(R_{\Gamma})_{\leq n} \cong \operatorname{gr}(\Bbbk A_{\Gamma})_{\leq n}$.

Proof. Let \Bbbk' be the image of \mathbb{Z} in \Bbbk ; it is either \mathbb{Z} or \mathbb{Z}/N for some integer N. The case \mathbb{Z} is already covered; if $\Bbbk' = \mathbb{Z}/N$, let p be a prime number dividing N. We prove the stronger statement that the composition $A_{\Gamma} \xrightarrow{\mu} \overline{R_{\Gamma}} \to \overline{R_{\Gamma} \otimes_{\mathbb{Z}} \mathbb{F}_p}$ is injective, i.e., we assume without loss of generality that $\Bbbk = \mathbb{F}_p$. Injectivity of μ for $\Bbbk = \mathbb{F}_p$ directly follows from Lemma 3.4.

It is an elementary calculation in non-commutative power series that the $1+\overline{\varpi}^n$ form a central series of subgroups of $1+\overline{\varpi}$. By the minimality and functoriality of the lower central series,

$$\gamma_n(1+\overline{\omega}) \subseteq 1+\overline{\omega}^n$$
 and then $\mu(\gamma_n(A_{\Gamma})) \subseteq 1+\overline{\omega}^n$.

Elementary calculations in the non-commutative power series ring also show that we have an isomorphism of associated graded Lie algebras

$$\bigoplus_{n\geq 1} (1+\overline{\varpi}^n)/(1+\overline{\varpi})^{n+1} \xrightarrow{\cong} \bigoplus_{n\geq 1} \overline{\varpi}^n/\overline{\varpi}^{n+1}; [1+w] \mapsto [w]$$

where the right hand side is the graded Lie algebra structure underlying the associated graded algebra R_{Γ} (with only the central summand ϖ/ϖ^1 of R_{Γ} missing). As R_{Γ} is already a graded algebra, it coincides with its associated graded. For details of these computations, compare, e.g., [Wad, Lemma 4.10].

Finally, the induced algebra map $\mathbb{k}A_{\Gamma} \to \overline{R_{\Gamma}}$ is compatible with the augmentation homomorphisms as the same is true for the initial map $\mu: A_{\Gamma} \to 1 + \overline{\varpi}$ (all elements on the left and on the right have augmentation 1). Consequently, it preserves the filtrations by powers of the augmentation ideals and induces a homomorphism $gr(\mu)$ on the associated graded algebra. On the generating set V this homomorphism is evidently the inverse of the map α of Lemma 3.1.

We learn that our homomorphism of complete filtered algebras $\overline{\mu} : \overline{\Bbbk A_{\Gamma}} \to \overline{R_{\Gamma}}$ induces an isomorphism of the associated graded algebras. By general theory therefore $\overline{\mu}$ itself is an isomorphism. In more detail, $\overline{\Bbbk A_{\Gamma}}$ is the inverse limit of the $\overline{\Bbbk A_{\Gamma}}/\overline{\varpi^n(A_{\Gamma})}$, and correspondingly for $\overline{R_{\Gamma}}$. Inductively and using the 5-lemma, $\overline{\mu}/\overline{\varpi^n} : \overline{\Bbbk A_{\Gamma}}/\overline{\varpi^n(A_{\Gamma})} \to \overline{R_{\Gamma}}/\overline{\varpi^n}$ is an isomorphism (as $\overline{\mu}/\overline{\varpi^n}$ is the extension of $\overline{\mu}/\overline{\varpi^{n-1}}$ by the isomorphism $\operatorname{gr}(\mu)_n$). Finally, $\overline{\mu}$ is an isomorphism as limit of isomorphisms.

4. Cohomology

A (topological) way to define and compute the cohomology of a discrete group G is via a *classifying space* X_G . By definition, this is a connected CWcell complex with $\pi_1(X_G) = G$ whose universal covering is contractible. We then have $H^*(G; \Bbbk) = H^*(X_G; \Bbbk)$.

Proof of Theorem 1.3, first claims. To compute the structure of the cohomology ring $H^*(A_{\Gamma}; \mathbb{k})$, we first show that X_{Γ} of (1.1) is a (particularly nice) classifying space for A_{Γ} . The space X_{Γ} inherits a CW-cell structure (indeed a cube complex structure) from the product cell structure of $(S^1)^V$, where S^1 has just one 0-cell {*} consisting of the base point and one 1-cell. Then X_{Γ} has a single vertex $*^V$ and precisely one loop $(S^1)^{\{v\}} \times \{*\}^{V \setminus \{v\}}$ for each generator $v \in V$. The 2-cells in X_{Γ} give the commutation relations. By the standard computation of the fundamental group of CW-complexes (based on the van Kampen theorem) we then have $\pi_1(X_{\Gamma}, *^V) = A_{\Gamma}$.

Furthermore, the link of the single vertex in X_{Γ} is a flag complex, since every subset of a clique is a clique. Therefore, X_{Γ} is a cube complex whose link is a flag complex, so X_{Γ} is a locally CAT(0) space [Gro], see [BH, Theorem 5.18], so its universal cover is contractible.

The cells given in the expression of X_{Γ} above form a basis of the homology of X_{Γ} : the differentials in the cellular chain complex vanish identically, because every cell sits in a subcomplex which is the cellular chain complex of a torus with precisely this property. Note that we get a basis of $H_*(X_{\Gamma}; \Bbbk)$ as free \Bbbk module by the images of the fundamental classes of all subtori T^{C} where C runs through the cliques in Γ . As the homology is finitely generated free, the cohomology is canonically the dual of the homology. We see that $H^*(X_{\Gamma}; \Bbbk)$ is precisely the quotient of the exterior algebra $H^*(T^V; \Bbbk) = \bigwedge^*(\Bbbk V)$, the cohomology of the ambient torus T^V , by the submodule generated by all products $v_1 \dots v_r$ such that v_1, \dots, v_r do not span a clique in Γ . The comparison map is induced by the inclusion $X_{\Gamma} \hookrightarrow T^{V}$. That this map is surjective with the claimed kernel follows by naturality and the know (co)homology of T^V , together with the information about the rank of $H^*(X_{\Gamma}; \mathbb{k})$ we obtained from the cellular complex. Now the quotient algebra is precisely the algebra S_{Γ} and we have proven $H^*(A_{\Gamma}; \mathbb{k}) = H^*(X_{\Gamma}; \mathbb{k}) = S_{\Gamma}$ as algebras.

We note that $H^*(X_{\Gamma}; \mathbb{k}) = S_{\Gamma}$ has a natural \mathbb{k} -basis indexed by cliques Cin Γ : a degree-k basis element corresponding to a clique $C = \{v_0, \ldots, v_{k-1}\}$ is given by the product $v_C := v_{k-1} \cdots v_0$ – to make this definite, we pick a total ordering of the vertices and write the factors in decreasing order.

4.1. Koszul algebras. Returning to general theory, consider a graded associative algebra R presented as T(W)/I for a finitely generated free k-module W, its tensor algebra T(W) and an ideal $I \leq T(W)$. In case I is generated by a subspace I_2 of $W^{\otimes 2}$, the algebra is called *quadratic*; and it then admits a *quadratic* dual $R^! := T(W^*)/(I_2^{\perp})$; here by I_2^{\perp} we mean the subset of $(W^*)^{\otimes 2} \cong (W^{\otimes 2})^*$ annihilating I_2 . Clearly $R^{!!} \cong R$. Now, with kV the free k-module with basis V, setting

$$G_R := \langle v \otimes w - w \otimes v \text{ for } \{v, w\} \in E \rangle \subset \mathbb{k} V^{\otimes 2},$$

$$G_S := \langle v \otimes w \text{ for } \{v, w\} \notin E, v \otimes w + w \otimes v \text{ for } \{v, w\} \in E \rangle \subset \mathbb{k} V^{\otimes 2},$$

we have as algebras

$$R_{\Gamma} = T(\Bbbk V)/\langle G_R \rangle$$
 and $S_{\Gamma} = \bigwedge^*(\Bbbk V)/\langle v \wedge w \text{ for } \{v, w\} \notin E \rangle = T(\Bbbk V)/\langle G_S \rangle.$

Let us identify $\mathbb{k}V^{\otimes 2}$ with $(\mathbb{k}V^{\otimes 2})^*$ via the basis $\{v \otimes w \mid v, w \in V\}$ and its dual basis. Then G_S is the annihilator of G_R (they clearly annihilate each other, and the ranks add up to the total dimension $|V|^2$), and therefore R_{Γ} and S_{Γ} are quadratic duals of each other.

Recall that a quadratic algebra R is called *Koszul* if its *Koszul complex* is acyclic, see [LV, §3.4.3]. We recall the Koszul complex (in our concrete

situation) below and we mention that this is only one of a number of different equivalent characterizations of the Koszul property. It implies that the Yoneda algebra $\text{Ext}_R(\Bbbk, \Bbbk)$ is isomorphic to $R^!$, compare [Pri, Theorem 2.5].

Proof of Theorem 1.3, second claim. We now show that R_{Γ} and S_{Γ} are Koszul. We use the fact [LV, Proposition 3.4.5] that a quadratic algebra is Koszul if and only if its quadratic dual is. Therefore it suffices to prove the Koszul property for S_{Γ} , and for supercommutative $(ab = (-1)^{\deg a \deg b} ba)$ algebras there is a simple sufficient (but not necessary) condition, the existence of a quadratic Gröbner basis [Yuz, Theorem 6.16]. Recall that a *Gröbner basis* for an ideal $I \leq \bigwedge^*(\mathbb{k}V)$ is a set G of generators for I such that the leading terms (with respect to a compatible order of monomials) of elements of G generate the same ideal as the leading terms of all elements of I. Now $G := \{v \land w \mid \{v, w\} \notin E\}$ is a Gröbner basis, as follows from *Buchberger's criterion*: "for all $f, f' \in G$ whose respective leading terms g, g' have least common multiple ℓ , the syzygy $(\ell/g) f - (\ell/g') f'$ must vanish".

Alternatively and without using Gröbner basis, the work of Fröberg [Fro, in particular Section 3] also implies that R_{Γ} (and S_{Γ}) are Koszul. His proof runs essentially as follows and uses directly the *Koszul complex* of R_{Γ} which we now construct. Consider the right R_{Γ} -module $P_* = \text{Hom}_{\Bbbk}(S_{\Gamma}, R_{\Gamma})$. Recall that, qua \Bbbk -module, S_{Γ} is finitely generated free with basis indexed by cliques in Γ . Consequently, this basis induces an isomorphism $P_* \xrightarrow{\cong} \bigoplus_C v_C R_{\Gamma}$, where the sum is over the cliques in Γ . It is bigraded by S_{Γ} - and R_{Γ} -degree. Consider the map $d: P_* \to P_*$ with

$$d(f)(p) = \sum_{v \in V} v f(vp) \quad \text{ for } f \in P_*, \ p \in S_{\Gamma}.$$

In our basis, $d((v_{k-1}\cdots v_0)\cdot r) = \sum_{j=1}^{j}(-1)^j(v_{k-1}\cdots \hat{v_j}\cdots v_0)\cdot v_jr$. A direct computation shows that $d^2 = 0$. Note that d increases the R_{Γ} -degree by 1, and decreases the S_{Γ} -degree by 1, so (P_*, d) becomes a chain complex of finitely generated free R_{Γ} -modules, graded by S_{Γ} -degree.

To prove acyclicity of the Koszul complex (P_*, d) we define a chain contraction map $s: P_* \to P_{*+1}$ of k-modules as follows. Recall that we have a k-basis of P_* given by elements $v_C \cdot w$ for a clique C of Γ and a basis element w of R_{Γ} given as a Γ -reduced monomial over V according to Definition 3.3. To define $s(v_C \cdot w)$ we consider two cases. If we can write w = vw' in reduced form with $v \in V$ and with w' a word in letters from V in such a manner that $v < \min C$ (for the total ordering on V picked above) and such that $C \cup \{v\}$ is a clique of Γ , then we choose v minimal with this property, and we set $s(v_C \cdot vw') := v_{C \cup \{v\}} \cdot w'$. Otherwise, we set $s(v_C \cdot w) := 0$. We now carry out the elementary calculation to see that s is a chain contraction, meaning $sd + ds = 1 - \epsilon$, where $\epsilon \colon P_* \to \Bbbk$ is the augmentation map, projecting onto the summand of bidegree (0,0). For this, consider $x = v_C \cdot w$. The calculation splits into three cases.

- (1) If $C = \emptyset$ and w = 1, then $(sd + ds)(x) = 0 = (1 \epsilon)(x)$.
- (2) Assume that $C = \{v_0, \dots, v_k\} \neq \emptyset$ and w cannot be written in the form w = vw' as above. Then

$$ds(x) = 0$$
 while $sd(x) = \sum (-1)^j s((v_{k-1} \cdots \widehat{v_j} \cdots v_0) \cdot v_j w).$

By hypothesis, no letter in w can be swapped with v_j and added to $C \setminus \{v_j\}$, so all summands vanish except the 0th which is x.

(3) Assume that $C = \{v_0, \ldots, v_k\}$ and w can be written in the form $v_{-1}w'$ such that $C \cup \{v_{-1}\}$ is a clique in Γ , with $v_{-1} < \min C$, chosen minimal among all such possibilities. Then v_{-1} commutes with all v_j , so

$$sd(x) = \sum (-1)^{j} s(v_{C \setminus \{v_{j}\}} \cdot v_{j} v_{-1} w') = \sum_{j=0}^{k-1} (-1)^{j} v_{C \setminus \{v_{j}\} \cup \{v_{-1}\}} \cdot v_{j} w',$$
$$ds(x) = d(v_{C \cup \{v\}} \cdot w') = \sum_{j=-1}^{k-1} (-1)^{j+1} v_{C \setminus \{v_{j}\} \cup \{v_{-1}\}} \cdot v_{j} w',$$

and the terms cancel pairwise except the one with j = -1, giving again (ds + sd)(x) = x.

It follows that P_* is a free R_{Γ} -resolution of \Bbbk .

We note that the usual definition of Koszul algebras is given over fields of characteristic 0; however, in our case, we need not impose any restriction on the commutative ring \Bbbk (other than interpreting $(\Bbbk V)^*$ as naturally isomorphic to $\Bbbk V$), since the rings R_{Γ} and S_{Γ} are \Bbbk -free.

5. Central series

5.1. Labute's general theory. Labute gave in [Lab] a condition under which a presentation $\langle V | \mathcal{R} \rangle$ of a group G determines a presentation of the associated Lie algebra $L(G) := \bigoplus_{n \ge 1} \gamma_n(G)/\gamma_{n+1}(G)$. Such a group presentation is now called "mild", and Anick gave in [Ani] a valuable criterion for this to happen: view all $r \in \mathcal{R}$ as elements of the free associative algebra $T(\mathbb{Z}V)$, under the Magnus embedding $F_V \to T(\mathbb{Z}V)$. Let n be such that $r - 1 \in \varpi^n \setminus \varpi^{n+1}$, and let r' denote the image of r in the quotient ϖ^n/ϖ^{n+1} . Then $\langle V | \mathcal{R} \rangle$

is mild if and only if $\{r' \mid r \in \mathcal{R}\}$ is "inert". We need not define here the meaning of "inert" (a.k.a. "strongly free", see, e.g., [HL]), but merely note that there are powerful sufficient conditions guaranteeing that a set is inert in the free associative algebra, one of them being that it forms a Gröbner basis. It follows then quite generally that the Lie algebra L(G) admits as presentation $\langle V \mid r' \forall r \in \mathcal{R} \rangle$, see [Lab, Theorem 1]; and a similar statement holds for the restricted Lie algebra $\bigoplus_{n\geq 1} \lambda_{n,p}(G)/\lambda_{n+1,p}(G)$, see [Lab, Theorem 3]. Labute's conditions are non-trivial to check, so we shall in fact recover his results rather than use them.

5.2. First easy results for RAAGs. By Proposition 3.6 the rings R_{Γ}/ϖ^n and $\Bbbk A_{\Gamma}/\varpi(G)^n$ are isomorphic, so the dimension subgroups $\delta_{n,\mu}$ and $\delta_{n,\Bbbk A_{\Gamma}}$ are equal. This establishes the corresponding part of Theorem 1.4ii, in particular (1.5). Furthermore, since the Magnus map μ has image in the subring of R_{Γ} generated by 1 and V, the groups $\delta_{n,\mu}$ depend on \Bbbk only via the image \Bbbk' of \mathbb{Z} in \Bbbk .

We consider two cases: if k has characteristic 0 then the dimension subgroups associated with the rings k and Q agree. If, on the other hand, k has characteristic p, then the dimension subgroups associated with the rings k and \mathbb{F}_p agree. In all cases, we reduce to the case $k \in \{\mathbb{F}_p, \mathbb{Z}\}$.

Proof of Equation (1.4) *of Theorem* 1.4ii. We apply the classical results of Jennings and Hall. For $\mathbb{k} = \mathbb{Q}$ we have $\gamma_{n,0} = \delta_{n,\mathbb{k}A_{\Gamma}}$; compare [Jen2, Hal2] which treat the case of torsion-free nilpotent groups to which the general case easily reduces. For $\mathbb{k} = \mathbb{F}_p$ we have $\gamma_{n,p} = \delta_{n,\mathbb{k}A_{\Gamma}}$; compare [Jen1] which treats the case of finite *p*-groups to which the general case easily reduces. Since we already established (1.5), Equation (1.4) follows.

Proof of second part of (1.2) in Theorem 1.4 i. In $\overline{R_{\Gamma}}$, clearly $\bigcap_{n\geq 0} \overline{\varpi}^n = \{0\}$. As μ is injective by the already established (1.3), it follows that $\bigcap_{n\geq 0} \delta_{n,\mu} = \{1\}$ and therefore by (1.4) also $\bigcap_{n\geq 0} \gamma_{n,0} = \{1\} = \bigcap_{n\geq 0} \gamma_{n,p}$.

6. Lie algebras associated with Γ

Recall that the cohomology of a Lie algebra L, defined as $\operatorname{Ext}_{U(L)}(\Bbbk, \Bbbk)$, may be computed using its *Chevalley complex* ($\bigwedge^*(L^\circ), d$), with L° the "small dual" of L, namely

 $L^{\circ} = \{ \phi \in L^* \mid \ker \phi \text{ contains a finite-codimensional ideal} \},\$

and the differential $d: L^{\circ} \to \bigwedge^2 L^{\circ}$ is the dual of the Lie bracket map $\bigwedge^2 L \to L$ (extended to all degrees by requiring d to be a graded derivation). Note that

 L° is just so defined that the image of d belongs to $\bigwedge^2 L^{\circ} \subset (\bigwedge^2 L)^*$. Since $\bigwedge^*(L^{\circ})$ is a graded commutative algebra and d is a derivation, the homology $(\bigwedge^*(L^{\circ}), d)$ is naturally a graded commutative algebra.

Proof of Theorem 1.5i, Lie algebra cohomology of L_{Γ} . The enveloping algebra of L_{Γ} is R_{Γ} , which is Koszul with Koszul dual S_{Γ} , so we have

$$H^*(L_{\Gamma}; \mathbb{k}) = H^*\left(\bigwedge^* (L^\circ), d\right) = \operatorname{Ext}_{R_{\Gamma}}(\mathbb{k}, \mathbb{k}) = S_{\Gamma}.$$

Note that $\bigwedge^* L_{\Gamma}^{\circ}$ admits two gradings, one as an exterior algebra and one inherited from the grading of L_{Γ} . In $H^*(\bigwedge^*(L^{\circ}), d)$, these two gradings coincide – this is precisely the content of S_{Γ} being a Koszul algebra.

In the following, we write L for L_{Γ} if the characteristic of \Bbbk is 0, and for $L_{\Gamma,p}$ if the characteristic of \Bbbk is p, and view L as a subset of $R_{\Gamma} = U(L)$. Following Magnus' method [MKS, Theorem 5.12], consider $x \in L_n$, i.e. homogeneous of degree n. Then x is a linear combination (with coefficients in \Bbbk) of a collection of bracket arrangements $\phi_i = \phi_i(v_1, \ldots, v_n)$. The assignment

$$L_n \ni \phi_i \mapsto \phi_i(v_1,\ldots,v_n) \in \gamma_n \subseteq A_{\Gamma}$$

is well defined on the subset of bracket arrangements, since $[v, w] = 1 \in A_{\Gamma}$ for each $\{v, w\} \in E$. It extends k-linearly to a map

$$\nu \colon L_n \to \gamma_n / \gamma_{n+1} \otimes_{\mathbb{Z}} \Bbbk$$

of k-modules. This map is clearly surjective, since γ_n/γ_{n+1} is spanned by *n*-fold bracket arrangements, for an arbitrary group. Furthermore, the composition $\mu_L \circ \nu \colon L \to R_{\Gamma}$ with μ_L given in Proposition 3.6 is a Lie algebra map sending v to v. Therefore this composition is the inclusion of L into R_{Γ} and is in particular injective. This implies that v is an isomorphism with inverse the Magnus map μ_L .

Proof of Theorem 1.4 i. Consider $\mathbb{k} = \mathbb{Z}$. Since L_{Γ} is \mathbb{Z} -free, it follows in particular that $\gamma_n(A_{\Gamma})/\gamma_{n+1}(A_{\Gamma})$ is torsion-free for each n, and therefore $\gamma_{n,0}(A_{\Gamma}) = \gamma_n(A_{\Gamma})$ for all n.

Proof of Theorem 1.5ii, *first two claims*. The isomorphism ν identifies L_{Γ} and $\bigoplus_{n\geq 1}(\gamma_n/\gamma_{n+1})\otimes_{\mathbb{Z}} \Bbbk$.

Proposition 6.1. Consider \Bbbk a ring in which p is regular (namely $px = 0 \Rightarrow x = 0$), and define the ideal $\varpi_p = \langle p, V \rangle$ of R_{Γ} .

The associated graded ring $\bigoplus_{n\geq 0} \overline{\varpi}_p^n / \overline{\varpi}_p^{n+1}$ is isomorphic to $R_{\Gamma} \otimes_{\mathbb{Z}} \mathbb{F}_p[\pi]$, with π of degree 1 mapped to $[p] \in \overline{\varpi}_p / \overline{\varpi}_p^2$ under the isomorphism.

Proof. Powers of ϖ_p define a new filtration on R_{Γ} , in which $v \in V$ still has degree 1, but in addition p also has degree 1; thus for instance p^2v^3 belongs to the fifth term of the filtration. The ring R_{Γ} is k-free. When passing to the associated graded ring for the new grading, we get on the one hand $\bigoplus \varpi_p^n / \varpi_p^{n+1}$. On the other hand, this graded ring is obtained from the old associated graded (which is the graded algebra R_{Γ}) by replacing each copy of k by its own associated graded under the filtration (p^n) , namely by $\bigoplus_{n\geq 0} p^n k / p^{n+1} k \cong (k / p k)[\pi]$; it is here that we require p to be regular. This replacement amounts to tensoring over \mathbb{Z} with $\mathbb{F}_p[\pi]$.

In case $p \ge 3$, we are now ready to identify the non-restricted Lie algebra $\bigoplus_{n\ge 1} \lambda_{n,p}/\lambda_{n+1,p}$ with $L_{\Gamma} \otimes_{\mathbb{Z}} \mathbb{F}_p[\pi]$. Let us temporarily write $\beta_n := \mu^{-1}(1 + \overline{\varpi}_p^n)$. We make the following claim.

Lemma 6.2. For $p \ge 3$ prime, the Magnus map μ induces a composition of (non-restricted) Lie algebra isomorphisms over $\mathbb{F}_p[\pi]$, still written μ_L ,

$$\mu_L \colon \bigoplus_{n \ge 1} \lambda_{n,p} / \lambda_{n+1,p} \to \bigoplus_{n \ge 1} \beta_n / \beta_{n+1} \to L_{\Gamma} \otimes_{\mathbb{Z}} \mathbb{F}_p[\pi],$$

with the first map induced by inclusion $\lambda_{n,p} \leq \beta_n$ and the second map induced by $\beta_n/\beta_{n+1} \ni [1+a] \mapsto a \in \overline{\omega}_p^n/\overline{\omega}_p^{n+1}$.

In particular, we have $\beta_n = \lambda_{n,p}$.

Proof. To check that the first map is well-defined, it suffices to show $\lambda_{n,p} \leq \beta_n$. We have $\varpi_p^n = \sum_{m+i\geq n} p^i \varpi^m$. Consider $g \in \gamma_m$, so by definition $\mu(g) = 1 + x$ for some $x \in \varpi^m$. We then have $\mu(g^{p^i}) = (1+x)^{p^i} = 1 + p^i x + \cdots \in 1 + \varpi_p^{m+i}$, so $\mu(\gamma_m^{p^i}) \subseteq 1 + \varpi_p^{m+i}$. Since $\lambda_{n,p} = \prod_{m+i\geq n} \gamma_m^{p^i}$, we have shown $\lambda_{n,p} \leq \beta_n$.

Because the Magnus map $\mu: A_{\Gamma} \to 1 + \overline{\varpi} \subset R_{\Gamma}$ is injective by Proposition 3.6, so is the induced map $\beta_n/\beta_{n+1} \to (1 + \overline{\varpi}_p^n)/(1 + \overline{\varpi}_p^{n+1}) = \overline{\varpi}_p^n/\overline{\varpi}_p^{n+1}$, which is our second map.

Since $p \ge 3$, the assignment $\pi \cdot [g] := [g^p]$ for $g \in \lambda_{n,p}$ (with $g^p \in \lambda_{n+1,p}$) gives $\bigoplus \lambda_{n,p}/\lambda_{n+1,p}$ the structure of an $\mathbb{F}_p[\pi]$ -module. For this we use the Hall–Petrescu identities [Hall, Theorems 3.1, 3.2]: If g,h belong to an arbitrary group G, then $(gh)^p \equiv g^p h^p [h, g]^n \alpha(g, h)$ with $n = \binom{p}{2}$ and $\alpha(\cdot, \cdot)$ a universal expression in γ_3 . This implies $(gh)^p = g^p h^p \mod \gamma_{n+2,p}$ for $g,h \in \gamma_{n,p}$ if either $n \ge 2$ or n = 1 and $p \ge 3$. However, beware that if n = 1 and p = 2then this does not hold in general, so the p-power operation is not linear. We see that μ_L maps this p-power operation to multiplication by π on $L_{\Gamma} \otimes_{\mathbb{Z}} \mathbb{F}_p[\pi]$. It follows that μ_L is an $\mathbb{F}_p[\pi]$ -Lie algebra homomorphism. Its image contains V which generates L_{Γ} , so μ_L is surjective. Finally, $L_{\Gamma} \otimes_{\mathbb{Z}} \mathbb{F}_p[\pi]$ is the free Lie algebra over $\mathbb{F}_p[\pi]$ modulo the relations [v, w] = 0 for $\{v, w\} \in E$. Those relations are clearly satisfied in the $\mathbb{F}_p[\pi]$ -Lie algebra $\bigoplus_{n\geq 1} \lambda_{n,p}/\lambda_{n+1,p}$, so the map μ_L is an isomorphism.

It then follows that the second map is surjective and therefore an isomorphism, so the first is also bijective, from which we deduce $\beta_n = \lambda_{n,p}$.

Proof of Theorem 1.4ii, $\delta_{n,\mu} = \gamma_{n,p}$ in characteristic p. Let \Bbbk be an algebra over \mathbb{F}_p . By [Quil], the Lie algebra $\bigoplus_{n\geq 1} (\gamma_{n,p}/\gamma_{n+1,p}) \otimes_{\mathbb{Z}} \Bbbk$ is isomorphic to the primitive subalgebra of $\bigoplus_{n\geq 0} \overline{\omega} (A_{\Gamma})^n / \overline{\omega} (A_{\Gamma})^{n+1} \cong R_{\Gamma}$, namely to $L_{\Gamma,p}$.

Proof of Theorem 1.5ii, *last claim*. This is precisely Lemma 6.2 and Proposition 6.1. \Box

7. Growth series

We derive now some relations between the Poincaré series of S_{Γ} , R_{Γ} , L_{Γ} and $L_{\Gamma,p}$ from general considerations. We recall that, for a graded algebra $R = \bigoplus_{n \ge 0} R_n$, its Poincaré series is $\Phi_R(t) = \sum_{n \ge 0} \operatorname{rank}(R_n)t^n$.

Proof of Theorem 1.6. First, we use Koszul duality between R_{Γ} and S_{Γ} to deduce $\Phi_{R_{\Gamma}}(t) \cdot \Phi_{S_{\Gamma}}(-t) = 1$, see [LV, Theorem 3.5.1]. This relationship between the Poincaré series of R_{Γ} and S_{Γ} was already noted in [CF, SY].

We have $\Phi_{S_{\Gamma}}(t) = \sum_{n \ge 0} \operatorname{rank} H^n(A_{\Gamma}, \mathbb{k})t^n = \sum_{n \ge 0} c_n(\Gamma)t^n$, with $c_n(\Gamma)$ the number of *n*-cliques in Γ , from our explicit basis of S_{Γ} given in Section 4.

The relation between $\Phi_{R_{\Gamma}}$ and $\Phi_{L_{\Gamma}}$ is given by the Poincaré–Birkhoff–Witt theorem, namely the fact that R_{Γ} and the symmetric algebra over L_{Γ} , respectively the degree- *p* truncated symmetric algebra over $L_{\Gamma,p}$, are isomorphic as graded \Bbbk -modules. It is expressed by the relation

$$\sum_{n \ge 0} a_n t^n = \prod_{n \ge 1} \left(\frac{1}{1 - t^n} \right)^{b_n} = \prod_{n \ge 1} \left(\frac{1 - t^{p_n}}{1 - t^n} \right)^{c_n}$$

if $\Phi_{R_{\Gamma}}(t) = \sum_{n \ge 0} a_n t^n$, $\Phi_{L_{\Gamma}}(t) = \sum_{n \ge 1} b_n t^n$, and $\Phi_{L_{\Gamma, p}}(t) = \sum_{n \ge 1} c_n t^n$.

Finally, we consider the growth series of the group A_{Γ} . It is the function $\Phi_{A_{\Gamma}}(t) = \sum_{g \in A_{\Gamma}} t^{\|g\|}$, with $\|g\|$ the minimal number of terms of $V \cup V^{-1}$ required to write g as a product. We cite [AP]:

$$\Phi_{A_{\Gamma}}(t) = \Phi_{R_{\Gamma}}\left(\frac{2t}{1+t}\right).$$

Indeed, as we saw in Proposition 3.5, every element $g \in A_{\Gamma}$ can be written in the form $g = v_1^{e_1} \cdots v_n^{e_n}$ for some $e_i \in \mathbb{Z} \setminus \{0\}$ as a word of minimal length; and

this expression is unique up to permuting some terms according to rule (M3). Let *I* be the set of (M3)-equivalence classes (v_1, \ldots, v_n) of minimal-length sequences. For an element $[v_1, \ldots, v_n]$ of *I*, the collection of all such terms $v_1^{e_1} \cdots v_n^{e_n}$ contributes $(t + t^2 + t^3 + \cdots)^n = (t/(1-t))^n$ to the growth series of R_{Γ} because each e_i can be an arbitrary positive natural number; and it contributes $(2t/(1-t))^n$ to the growth of A_{Γ} , taking into account the signs of the e_i . Since we obtain all elements of A_{Γ} and all basis elements of R_{Γ} that way, we have

$$\Phi_{A_{\Gamma}}(t) = \sum_{I} \left(\frac{2t}{1-t}\right)^{n} = \sum_{I} \left(\frac{2t/(1+t)}{1-2t/(1+t)}\right)^{n} = \Phi_{R_{\Gamma}}\left(\frac{2t}{1+t}\right),$$

using 2t/(1-t) = (2t/(1+t))/(1-2t/(1+t)). We have finished the proof of Theorem 1.6.

8. Malcev completions

In this section we fix $\mathbb{k} = \mathbb{Q}$. Recall from [PS1] that a *Malcev Lie algebra* is a Lie algebra L over \mathbb{Q} , given with a descending filtration $(L_n)_{n\geq 1}$ of ideals such that L is complete with respect to the associated topology, and satisfying $L_1 = L$ and $[L_m, L_n] \subseteq L_{m+n}$ and such that $\bigoplus_{n\geq 1} L_n/L_{n+1}$ is generated in degree 1. Every Malcev Lie algebra admits an associated *exponential group* $\exp(L)$, which is L as a set, with product given by the Baker–Campbell–Hausdorff formula $x \cdot y = x + y + [x, y]/2 + \cdots$.

Lazard proved in [Laz] that every group homomorphism $\rho: G \to \exp(L)$ induces a morphism of graded Lie algebras $\bigoplus_{n\geq 1} \gamma_n/\gamma_{n+1} \otimes \mathbb{Q} \to \bigoplus_{n\geq 1} L_n/L_{n+1}$.

A Malcev completion of a group G is a homomorphism $\rho: G \to \exp(L)$ for a Malcev Lie algebra L, universal in the sense that every representation $G/\gamma_n \to \exp(L')$ for a (nilpotent) Malcev Lie algebra L' factors uniquely through $\exp(L/L_n)$; see [PS1, Definition 2.3].

Quillen gave a direct construction of the Malcev completion of a group in [Quil, Qui2]: let $\overline{\mathbb{Q}G} = \operatorname{proj}\lim \mathbb{Q}G/\varpi^n$ be the completion of the group ring; then $\overline{\mathbb{Q}G}$ is a complete Hopf algebra. Let *L* be its Lie subalgebra of primitive elements; it is a Malcev Lie algebra for the filtration $L_n = L \cap \overline{\varpi}^n$. Let exp: $L \to \overline{\mathbb{Q}G}$ be the usual power series map $\exp(x) = 1 + x + x^2/2 + \cdots$ which makes sense in $\overline{\mathbb{Q}G}$. Then its image $\overline{G} := \exp(L)$ is a subgroup of the group of multiplicative units. It identifies with the Lie group associated to the Malcev Lie algebra *L*, and it consists precisely of the grouplike elements in $\overline{\mathbb{Q}G}$, namely the $g \in 1 + \overline{\varpi}$ satisfying $\Delta(g) = g \otimes g$. The representation $\rho: G \to \exp(L); g \mapsto g$ is the Malcev completion of *G*. The Magnus map $\mu: A_{\Gamma} \to \overline{R_{\Gamma}}^{\times}$ yields an isomorphism of associative algebras $\overline{\mathbb{Q}A_{\Gamma}} \cong \overline{R_{\Gamma}}$. Both algebras are actually complete Hopf algebras, but the Magnus isomorphism does not preserve the Hopf algebra structure: $v \in V \subset \overline{\mathbb{Q}A_{\Gamma}}$ is group-like, meaning $\Delta(v) = v \otimes v$ while $v \in V \subset \overline{R_{\Gamma}}$ is primitive, meaning $\Delta(v) = v \otimes 1 + 1 \otimes v$; so $\Delta(\mu(v)) = \Delta(1 + v) = 1 \otimes 1 + v \otimes 1 + 1 \otimes v$ while $(\mu \otimes \mu)(\Delta(v)) = (1 + v) \otimes (1 + v)$.

The Magnus map μ is, in fact, the truncation to order 1 of a Hopf algebra isomorphism $\mu_{\exp}: \overline{\mathbb{Q}A_{\Gamma}} \to \overline{R_{\Gamma}}$, given on $v \in V$ by the classical exponential series

$$\mu_{\exp}(v) = \sum_{n \ge 0} \frac{v^n}{n!} = 1 + v + \mathcal{O}(v^2).$$

Proof of Theorem 1.7. The proof that μ_{exp} is an isomorphism of filtered associative algebras is exactly the same as that of Theorem 1.4ii, and will not be repeated. On the other hand, the fact that μ_{exp} is a coalgebra map follows formally from the fact that the power series exp maps primitive elements to group-like elements:

$$\Delta(\mu_{\exp}(v)) = \Delta\left(\sum_{n\geq 0} v^n/n!\right) = \sum_{n\geq 0} \Delta(v)^n/n!$$
$$= \sum_{n\geq 0} \frac{(v\otimes 1+1\otimes v)^n}{n!} = \sum_{\ell,m\geq 0} \frac{(v\otimes 1)^\ell (1\otimes v)^m}{\ell!m!}$$
$$= (\exp v \otimes 1)(1\otimes \exp v) = (\mu_{\exp}\otimes \mu_{\exp})(\Delta(v)).$$

We have proven the first claim.

It now suffices to use this isomorphism μ_{exp} to make even more concrete the construction of Quillen sketched above: in $\overline{\mathbb{Q}A_{\Gamma}}$ the space of primitive elements is slightly mysterious, for example, it contains

$$\log(g) = \log(1 - (1 - g)) = -\sum_{n \ge 1} (1 - g)^n / n \quad \forall g \in A_{\Gamma}.$$

In contrast to this, its exponential is the Malcev completion naturally containing A_{Γ} . In $\overline{R_{\Gamma}}$ the opposite holds: the space of primitive elements is the Lie subalgebra L_{Γ} while its exponential cannot be better defined than as the exponential of L_{Γ} .

In all cases, the Hopf algebra isomorphism μ_{exp} directly yields the remaining claims of Theorem 1.7.

We now turn to formality in the sense of Sullivan in rational homotopy theory. A finite CW-complex X is called *formal* if its algebraic minimal model is quasiisomorphic to $(H^*(X; \mathbb{Q}), 0)$. This implies that the rational homotopy type of X is determined in a precise way by its rational cohomology ring. For details on rational homotopy theory compare [Sul] or the more recent [FHT]. We finally prove that the space X_{Γ} constructed in Section 4 is formal. Recall that we defined X_{Γ} as a (cubical) subspace of the smooth manifold $(\mathbb{R}/\mathbb{Z})^V$. It makes perfect sense to restrict smooth differential forms on $(\mathbb{R}/\mathbb{Z})^V$ to X_{Γ} . We define $A^*(X_{\Gamma})$ to be the algebra of all such restrictions; it is a commutative differential graded algebra (cdga). It is an easy exercise that this cdga is quasiisomorphic to the standard cdga over \mathbb{R} of rational homotopy theory associated to X_{Γ} . There are basic one-forms dx_v on $(\mathbb{R}/\mathbb{Z})^V$ coming from the obvious coordinate functions, for $v \in V$. Their images in $A^*(X_{\Gamma})$ generate a sub-cdga with trivial differential, whose homology is $H^*(X_{\Gamma}; \mathbb{R})$ by Theorem 1.3. The inclusion of this sub-cdga in $A^*(X_{\Gamma})$ is a quasi-isomorphism, showing that X_{Γ} is formal.

We now explicitly exhibit a minimal model for X_{Γ} . Recall from Section 6 the Chevalley complex $(\bigwedge^*(L_{\Gamma}^\circ), d)$ of L_{Γ} . Note that L_{Γ} is graded, and L_{Γ}° may be identified with the graded dual of L_{Γ} . Consequently, there is a natural map $L_{\Gamma}^\circ \to \mathbb{Q}V$ given by restricting to the degree-1 part. This map induces a map of graded algebras $\bigwedge^*(L_{\Gamma}^\circ) \to \bigwedge^*(\mathbb{Q}V)/\langle v \land w$ for $\{v, w\} \notin E \rangle = S_{\Gamma}$. Even better, this is a map of cdgas from the Chevalley complex to S_{Γ} , the latter equipped with zero differential, and indeed is a quasi-isomorphism. These are manifestations of the Koszul duality of S_{Γ} and $R_{\Gamma} = U(L_{\Gamma})$. As X_{Γ} is formal and $S_{\Gamma} = H^*(X_{\Gamma}; \mathbb{Q})$ we conclude that $\bigwedge^*(L_{\Gamma}^\circ, d)$ is a minimal model of X_{Γ} .

Here is yet an alternative proof: a group is called 1-*formal* if its Malcev Lie algebra is quadratic. It therefore follows from Theorem 1.7 that A_{Γ} is 1-formal. The cohomology ring $H^*(X_{\Gamma}; \mathbb{Q}) \cong S_{\Gamma}$ is Koszul by Theorem 1.3, so X_{Γ} is formal by [PS2, Proposition 2.1].

9. Outlook

9.1. Subgroup growth. Baik, Petri, and Raimbault determined the subgroup growth of A_{Γ} in terms of the graph Γ . Define $s_n(A_{\Gamma})$ as the number of subgroups of A_{Γ} of index precisely *n*. Then [BPR, Theorem A] establishes

$$\lim_{n \to \infty} \frac{\log(s_n(A_{\Gamma}))}{n \log(n)} = \alpha(\Gamma) - 1,$$

i.e., $s_n(A_{\Gamma})$ grows like $(n!)^{\alpha(\Gamma)-1}$. Here, $\alpha(\Gamma)$ is the *independence number* of Γ , the largest number of vertices such that the full subgraph of Γ spanned by them is discrete.

We do not discuss the rather complicated proof here. We leave it an open question to find a corresponding result for the growth of the number of finite index Lie subalgebras of L_{Γ} . Indeed, we expect that these two series are closely related and that the latter is slightly easier to control than $(s_n(A_{\Gamma}))_{n \in \mathbb{N}}$.

We have identified $\gamma_{n,p}(A_{\Gamma})$ with $\delta_{n,\mathbb{F}_{p_p}A_{\Gamma}}$ in Theorem 1.4ii. For a group G, we could define γ_{n,p^e} as the subgroup generated by γ_n and all $\gamma_i^{p^j}$ with $ip^j \ge np^{e-1}$. When G is free, it was shown by Lazard that $\gamma_{n,p^e}(G)$ coincides with the dimension subgroup $\delta_{n,\mathbb{Z}/p^e\mathbb{Z}[G]}$ while this does not hold for general G, see [Mor].

We leave it as an exercise to extend Lazard's result to A_{Γ} .

9.2. Homology gradients. Given a group G and a nested sequence of finite index normal subgroups $G_n \triangleleft G$ with $\bigcap_n G_n = \{1\}$, one defines for a field k the k-homology gradients

$$b_i^{(2)}(G; \mathbb{k}) := \limsup_n \frac{b_i(G_n; \mathbb{k})}{[G:G_n]} \text{ for } i \in \mathbb{N}.$$

For general groups G, it is unclear whether this quantity depends on the particular chain $\{G_n\}$. Until recently, it was also unclear in which manner this quantity depends on the coefficients k. Avradmidi, Okun, and Schreve in [AOS] use the classifying space X_{Γ} and induced cell structures for coverings to explicitly compute these homology gradients. Let F_{Γ} be the flag complex generated by Γ , i.e., the largest simplicial complex with vertex set V and edge set E. Then

$$b_i^{(2)}(A_{\Gamma}; \Bbbk) = \overline{b_{i-1}}(F_{\Gamma}; \Bbbk)$$

where $\overline{b_*}(F_{\Gamma}; \Bbbk)$ denotes the dimension of the reduced homology of F_{Γ} . In particular, for RAAGs the homology gradient is independent of the chain of normal subgroups, even though in many examples it does depend on the field of coefficients \Bbbk .

Acknowledgments. We are deeply grateful to Jacques Darné, Pierre de la Harpe and the anonymous referee for their well-thought comments on preliminary versions of our text.

References

- [Ani] D. J. ANICK, Inert sets and the Lie algebra associated to a group. J. Algebra 111 (1987), 154–165. MR 0050280 MR 0913201
- [AOS] G. AVRAMIDI, B. OKUN and K. SCHREVE, Mod *p* and torsion homology growth in nonpositive curvature (2020). Preprint. arXiv:2003.01020
- [AP] J. S. ATHREYA and A. PRASAD, Growth in right-angled groups and monoids (2014). Preprint. arXiv:1409.4142
- [BH] M. R. BRIDSON and A. HAEFLIGER, *Metric Spaces of Non-Positive Curvature*. Springer-Verlag, Berlin (1999). Zbl 0988.53001 MR 1744486

- [BPR] H. BAIK, B. PETRI and J. RAIMBAULT, Subgroup growth of right-angled Artin and Coxeter groups. J. London Math. Soc. (2) 101 (2020), 556–588. Zbl 07216611
- [CF] P. CARTIER and D. FOATA, Problèmes combinatoires de commutation et réarrangements. Lecture Notes in Mathematics, No. 85. Springer-Verlag, Berlin, New York (1969). Zbl 0186.30101 MR 0239978
- [DK1] G. H. E. DUCHAMP and D. KROB, The free partially commutative Lie algebra: Bases and ranks. *Adv. Math.* **95** (1992), 92–126. Zb10763.17003 MR1176154
- [DK2] The lower central series of the free partially commutative group. *Semigroup Forum* **45** (1992), 385–394. Zbl 0814.20025 MR 1179860
- [Dro] C. G. A. DROMS, *Graph Groups (Algebra, Kim, Roush, Magnus)*. ProQuest LLC, Ann Arbor, MI (1983). MR 2633165
- [FHT] Y. FÉLIX, S. HALPERIN and J.-C. THOMAS, Rational Homotopy Theory, vol. 205 of Graduate Texts in Mathematics. Springer-Verlag, New York (2001). Zbl 0961.55002 MR 1802847
- [Fro] R. FRÖBERG, Determination of a class of Poincaré series. Math. Scand. 37 (1975), 29–39. Zbl 0318.13027 MR 0404254
- [Gro] M. L. GROMOV, Hyperbolic groups (1987). pp. 75–263. Zbl 0634.20015 MR 919829
- [Gru] K. W. GRUENBERG, Residual properties of infinite soluble groups. Proc. London Math. Soc. (3) 7 (1957), 29–62. Zbl 0077.02901 MR 0087652
- [Hal1] P. HALL, A contribution to the theory of groups of prime-power order. Proc. London Math. Soc. (2) 36 (1934), 29–95. Zbl 0007.29102 MR 1575964
- [Hal2] The Edmonton Notes on Nilpotent Groups. Queen Mary College Mathematics Notes. Mathematics Department, Queen Mary College, London (1969). Zbl 0211.34201 MR 0283083
- [HL] S. HALPERIN and J.-M. LEMAIRE, Suites inertes dans les algèbres de Lie graduées ("autopsie d'un meurtre. II"). *Math. Scand.* 61 (1987), 39–67. Zbl 0655.55004 MR 929396
- [Jac] N. JACOBSON, Restricted Lie algebras of characteristic p. Trans. Amer. Math. Soc. 50 (1941), 15–25. Zb10025.30301 MR 0005118
- [Jen1] S. A. JENNINGS, The structure of the group ring of a *p*-group over a modular field. *Trans. Amer. Math. Soc.* **50** (1941), 175–185. Zbl 0025.24401 MR 0004626
- [Jen2] The group ring of a class of infinite nilpotent groups. *Canad. J. Math.* 7 (1955), 169–187. Zb10066.01302 MR 0068540
- [KM] M. KAPOVICH and J. J. MILLSON, On representation varieties of Artin groups, projective arrangements and the fundamental groups of smooth complex algebraic varieties. *Inst. Hautes Études Sci. Publ. Math.* 88 (1998), 5–95 (1999). Zbl 0982.20023 MR 1733326
- [Lab] J. P. LABUTE, The determination of the Lie algebra associated to the lower central series of a group. *Trans. Amer. Math. Soc.* 288 (1985), 51–57. Zbl 0576.20022 MR 773046
- [Laz] M. LAZARD, Sur les groupes nilpotents et les anneaux de Lie. Ann. École Norm. Sup. (3) 71 (1954), 101–190. Zbl 0055.25103 MR 0088496

- [LV] J.-L. LODAY and B. VALLETTE, Algebraic operads, vol. 346 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer, Heidelberg (2012). Zbl 1260.18001 MR 2954392
- [Mag] W. MAGNUS, Beziehungen zwischen Gruppen und Idealen in einem speziellen Ring. Math. Ann. 111 (1935), 259–280. Zbl 0011.15201 MR 1512992
- [Mal] A. I. MAL'CEV, Nilpotent torsion-free groups. Izvestiya Akad. Nauk. SSSR. Ser. Mat. 13 (1949), 201–212. MR 0028843
- [MKS] W. MAGNUS, A. KARRASS and D. SOLITAR, Combinatorial group theory. Dover Publications, Inc., New York, second revised edition ed. (1976). Zbl 1130.20307 MR 0422434
- [MM] J. W. MILNOR and J. C. MOORE, On the structure of Hopf algebras. *Ann. of Math.* (2) **81** (1965), 211–264. Zbl 0163.28202 MR 0174052
- [Mor] S. MORAN, Dimension subgroups modulo *n*. *Proc. Cambridge Philos. Soc.* **68** (1970), 579–582. Zbl 0215.10802 MR 0267003
- [Pri] S. B. PRIDDY, Koszul resolutions. Trans. Amer. Math. Soc. 152 (1970), 39–60. Zbl 0261.18016 MR 0265437
- [PS1] S. PAPADIMA and A. I. SUCIU, Chen Lie algebras. Int. Math. Res. Not. 21 (2004), 1057–1086. Zbl 1076.17007 MR 2037049
- [PS2] Algebraic invariants for right-angled Artin groups. Math. Ann. 334 (2006), 533–555. Zbl1165.20032 MR 2207874
- [Qui1] D.G. QUILLEN, On the associated graded ring of a group ring. J. Algebra 10 (1968), 411–418. Zbl 0192.35803 MR 0231919
- [Qui2] Rational homotopy theory. Ann. of Math. (2) **90** (1969), 205–295. Zbl 0191.53702 MR 0258031
- [Sch] W. SCHMITT, Hopf algebras and identities in free partially commutative monoids. *Theoretical Computer Science* **73** (1990), 335–340. Zbl 0694.68056 MR 1073385
- [Ser] J.-P. SERRE, Lie Algebras and Lie Groups, vol. 1500 of Lecture Notes in Mathematics. Springer-Verlag, Berlin (2006). Zbl 0742.17008 MR 2179691
- [Sul] D. SULLIVAN, Infinitesimal computations in topology. Inst. Hautes Études Sci. Publ. Math. 47 (1977), 269–331 (1978). Zb10374.57002 MR 0646078
- [Swe] M.E. Sweedler, *Hopf Algebras*. Mathematics Lecture Note Series. W. A. Benjamin, Inc., New York (1969). Zbl 0194.32901 MR 0252485
- [SY] B. SHELTON and S. YUZVINSKY, Koszul algebras from graphs and hyperplane arrangements. J. London Math. Soc. (2) 56 (1997), 477–490. Zbl 0959.16026 MR 1610447
- [Wad] R. D. WADE, The lower central series of a right-angled Artin group. *Enseign. Math.* 61 (2015), 343–371. Zb11355.20023 MR 3539842
- [Yuz] S. YUZVINSKIĬ, Orlik–Solomon algebras in algebra and topology. Uspekhi Mat. Nauk 56 (2001), 87–166. Zb11033.52019 MR 1859708

60

 [Zas] H. ZASSENHAUS, Ein Verfahren, jeder endlichen p-Gruppe einen Lie-Ring mit der Charakteristik p zuzuordnen. Abh. Math. Sem. Univ. Hamburg 13 (1939), 200–207. JFM 65.0091.01 MR 3069703

(Reçu le 27 avril 2019)

Laurent BARTHOLDI, Mathematisches Institut, Georg-August Universität Göttingen, Bunsenstraße 3–5, 37073 Göttingen, Germany École Normale Supérieure, Lyon, France

e-mail: laurent.bartholdi@math.uni-goettingen.de

Henrika Härer, Mathematisches Institut, Georg-August Universität Göttingen, Bunsenstraße 3–5, 37073 Göttingen, Germany

e-mail: henrikamaria.runde@stud.uni-goettingen.de

Thomas Schick, Mathematisches Institut, Georg-August Universität Göttingen, Bunsenstraße 3–5, 37073 Göttingen, Germany

e-mail: thomas.schick@math.uni-goettingen.de