Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	66 (2020)
Heft:	1-2
Artikel:	A linking invariant for algebraic curves
Autor:	Guerville-Ballé, Benoît / Meilhan, Jean-Baptiste
DOI:	https://doi.org/10.5169/seals-880823

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

A linking invariant for algebraic curves

Benoît GUERVILLE-BALLÉ and Jean-Baptiste MEILHAN

Abstract. We construct a topological invariant of algebraic plane curves, which is in some sense an adaptation of the linking number of knot theory. This invariant is shown to be a generalization of the \mathcal{I} -invariant of line arrangements developed by the first author with Artal and Florens. We give two practical tools for computing this invariant, using a modification of the usual braid monodromy or using the connected numbers introduced by Shirane. As an application, we show that this invariant distinguishes several Zariski pairs, i.e., pairs of curves having same combinatorics, yet different topologies. The former is the well known Zariski pair found by Artal, composed of a smooth cubic with 3 tangent lines at its inflexion points. The latter is formed by a smooth quartic and 3 bitangents.

Mathematics Subject Classification (2010). Primary: 14H50; Secondary: 32Q55, 54F65, 57M20.

Keywords. Zariski pair, algebraic plane curves, embedded topology, topological invariant.

Introduction

The topological study of algebraic plane curves was initiated at the beginning of the 20th century by Klein and Poincaré. One of the main questions is to understand the relationship between the combinatorics and the topology of a curve. It is known, since the seminal work of Zariski [Zar1, Zar2, Zar3], that the topological type of the embedding of an algebraic curve in the complex projective plane is not determined by the combinatorics. Indeed, Zariski constructed two sextics with 6 cusps having same combinatorics, and proved that the fundamental group of their complements are not isomorphic. Geometrically, these two curves are distinguished by the fact that the cusps in the first curve lie on a conic, while they do not in the second curve. Since this historical example, using various methods, numerous examples of pairs of algebraic curves having same combinatorics but different topologies have been found, see for example Artal, Cogolludo and Tokunaga [ACT], Cassou-Noguès, Eyral and Oka [CNEO], Degtyarev [Deg], Oka [Oka], Shimada [Shim1], or the first author [GB]. Artal suggests in [Art] to call such examples *Zariski pairs*.

The topology of curves in \mathbb{CP}^2 is intimately connected to the topology of knots and links in S^3 . Several tools are indeed shared by these two domains, such as the homology or the fundamental group of the complement, the Alexander polynomial or module, although they usually have rather different behaviours.

Recently, Artal, Florens and the first author defined a topological invariant of line arrangements (i.e. algebraic plane curves with only irreducible components of degree 1) which is in some sense modelled on the linking number of knot theory [AFGB]. This invariant was then successfully used in [GB] to distinguish a new Zariski pair of line arrangements. In the present paper, we construct another invariant adapting the linking number to the more general case of algebraic plane curves. In the case of a line arrangement, this invariant is shown to be equivalent to the invariant of [AFGB], thus providing a generalization of this earlier work through a different adaptation of the linking number.

The construction of our linking invariant can be roughly outlined as follows. Consider a reducible algebraic curve decomposed in two nonempty subcurves Cand D, and pick a topological cycle γ in the subcurve C. The basic idea is to consider the image of a certain coset of γ in the first homology group of $\mathbb{CP}^2 \setminus D$. More precisely, this set is regarded in the quotient of $H_1(\mathbb{CP}^2 \setminus D)$ by an appropriate *indeterminacy subgroup* \mathcal{J}_C , which controls the topological differences among the various cycles in the considered coset of γ . This define the *linking invariant of* C *with* D *along* γ , which is an invariant of the pair $(\mathbb{CP}^2, C \cup D)$.

Our construction thus builds on a rather elementary idea, and is not technically involved. Remarkable is rather the fact that it reveals quite efficient in practice, despite its apparent simplicity. We mention below several applications of the linking invariant on concrete examples of Zariski pairs of various natures.

This linking invariant has a nice behaviour for some particular choices of curve or cycle. In the case of line arrangements, for example, the linking invariant is indeed a single homology class rather than a coset. This allows us to prove the equivalence with the \mathcal{I} -invariant of [AFGB] in this case.

From a practical viewpoint, we provide two methods of computation of this linking invariant. The first one is based on a topological construction using an adaptation of the braid monodromy. This makes a concrete connection between our invariant and the usual linking number of knot theory. The second method is algebraic and comes from the relation, observed in [GBS], between our linking invariant, the connected numbers and the splitting numbers introduced by Shirane in [Shir1, Shir2].

To illustrate the efficiency of this adaptation of the linking number to algebraic curves, we use it to distinguish two examples of Zariski pairs. The first example is formed by the well known 3-Artal curves introduced by Artal in [Art]. They are composed of a smooth cubic and three inflexional tangent; in the first curve the considered inflexion points are collinear, while they are not in the second one. The computation of the linking invariant of the three lines with the cubic is made using the above mentioned algebraic method. The second example of Zariski pair is formed by a smooth quartic and three bitangents. These curves have been very recently studied in [BTY]. For that example, we use the topological method based on the linking number of knot theory.

After an earlier version of this paper was circulated, our linking invariant (then called *linking set*) has been further studied, and being used to distinguish other examples of Zariski pairs.

In [Shir1] Shirane introduces the *splitting numbers* and detects the π_1 -equivalent Zariski *k*-plets suggested by Shimada [Shim2]. By proving that the splitting numbers and the linking invariant are equivalent (in some particular cases), the first author and Shirane obtain in [GBS] that the linking invariant distinguishes the Shirane-Shimada π_1 -equivalent Zariski *k*-plets. This implies that the linking invariant is not determined by the fundamental group of the complement.

The linking invariant is also used in [BGBS] to classify the topology of the k-Artal curves (i.e., a smooth cubic and k inflectional tangent lines). Furthermore, Shirane constructed recently in [Shir2] an adaptation of the splitting number, called the *connected numbers*, which allows to classify the topology of the Artal curves of degree b (i.e., smooth curves of degree b and with three total inflectional tangent lines). Here again, the proofs of [GBS] imply that the linking invariant can distinguish the Artal curves of degree b.

In the particular case of line arrangements, the linking invariant (in the form of the \mathcal{I} -invariant) has been successfully used in [GB] to detect a Zariski pair of 12 lines. Recently, the first author and Viu-Sos gave an effective diagrammatic reformulation of this invariant in the particular case of real line arrangements, see [GBVS]. Using this reformulation, they provide 10 examples of complexified real Zariski pairs.

Convention. All homology groups are to be understood with integral coefficients, and this will be omitted in the notation.

1. The linking invariant

1.1. Preliminaries. Let C be an algebraic plane curves, possibly non-reduced. Following [ACT], we define the combinatorics of C as the data:

 $\left(\operatorname{Irr}(\mathcal{C}), \operatorname{deg}, \operatorname{Sing}(\mathcal{C}), \Sigma_{\operatorname{top}}, \sigma_{\operatorname{top}}, \{\mathcal{C}(P)\}_{P \in \operatorname{Sing}(\mathcal{C})}, \{\beta_P\}_{P \in \operatorname{Sing}(\mathcal{C})}\right),$

where:

- $Irr(\mathcal{C})$ is the set of all irreducible components of \mathcal{C} ,
- deg assigns to each irreducible component its degree,
- Sing(C) is the set of all singular points of C,
- Σ_{top} is the set of topological types of singular points of C, and σ_{top} assigns to each singular point its topological type,
- for each singular point P of C, C(P) is the set of local branches of C at P, and β_P assigns to each local branch at P the global irreducible component containing it.

Two curves have the same combinatorics if there exist bijections between their sets Irr and Sing of irreducible components and singular points, which are compatible with the sets C(P) and Σ_{top} of local branches and topological types, and with the assignments deg, σ_{top} and $\{\beta_P\}$ in the natural way; see [ACT, Rem. 3] for details.

We also associate to the curve C the *intersection graph* Γ_C of its irreducible components. This is a bipartite graph whose first set of vertices, called *component-vertices*, corresponds to the irreducible components of C, while the second set of vertices, called *point-vertices*, corresponds to the singular points of C contained in at least two distinct irreducible components. An edge of Γ_C joins a point-vertex and a component-vertex if and only if the corresponding singular point is contained in the corresponding irreducible component. Note that the information encoded in Γ_C are contained (but not equivalent) to the combinatorics of C. For example, the information given by σ_{top} is not contained in Γ_C .

A cycle of Γ_{C} is a (non necessarily connected) closed oriented walk without repeated edges which contain at least one component-vertex. Note that this includes the case of a single vertex. A (combinatorial) cycle of Γ_{C} can be lifted to a (topological) cycle on the curve C, i.e., an oriented closed loop in C, although it is not uniquely determined in general. Such a topological lift has a natural induced orientation only if the combinatorial cycle is not simply connected. If the combinatorial cycle is simply connected (i.e., it consists of a single componentvertex) then a topological lift is any (possibly non-contractible) closed loop in the corresponding irreduccible component of C, with an arbitrary choice of orientation.

In what follows, we will be mainly interested in reducible algebraic curves $C \cup D$, which decompose into two subcurves C and D (without common irreducible component). In this context, a cycle of Γ_C is simply a cycle lying in the subgraph Γ_C of $\Gamma_{C\cup D}$. On one hand, such a cycle is called *maximal* if it contains all component-vertices of Γ_C ; in other words, a maximal cycle in Γ_C lifts to a topological cycle in $C \cup D$ that intersects the smooth part of all irreducible components of C. On the other hand, a cycle of $\Gamma_{C\cup D}$ is said to *avoid* D if it lies in Γ_C (and is thus disjoint from all component-vertices of D) and avoids all point-vertices of $C \cap D$.

In this paper, by a homeomorphism ϕ between two such reducible algebraic curves $C_1 \cup D_1$ and $C_2 \cup D_2$, we will always mean an ambient homeomorphism of \mathbb{CP}^2 which sends C_1 to C_2 and D_1 to D_2 . Furthermore, we will denote by $\phi_{\Gamma} : \Gamma_{C_1 \cup D_1} \to \Gamma_{C_2 \cup D_2}$ the induced map at the combinatorial level. Note that, if ϕ is orientation preserving, then ϕ_{Γ} preserves the cycle orientation.

1.2. The linking invariant. Let $C \cup D$ be a reducible algebraic curve, decomposed into two nonempty subcurves C and D.

Consider the inclusion maps $i : C \setminus D \hookrightarrow C$ and $j : C \setminus D \hookrightarrow \mathbb{CP}^2 \setminus D$, and denote respectively by i_* and j_* the induced map on the first homology groups. Note that ker (i_*) identifies with $H_1(\partial \bigcup_{C \in Irr(C)} C \setminus D) \simeq \bigoplus_{C \in Irr(C)} H_1(\partial(C \setminus D))$ in $H_1(C \setminus D)$.

Definition 1.1. The *indeterminacy subgroup* with respect to C, denoted by \mathcal{J}_C , is the subgroup of $H_1(\mathbb{CP}^2 \setminus D)$ defined as the image of $\bigoplus_{C \in Irr(C)} H_1(\partial(C \setminus D))$ by j_* .

Now, let γ be a maximal cycle in $\Gamma_{\mathcal{C}}$ avoiding \mathcal{D} . Pick a topological lift $\tilde{\gamma}$ of γ on the curve $\mathcal{C} \subset \mathcal{C} \cup \mathcal{D}$. By assumption, $\tilde{\gamma}$ lies in $\mathcal{C} \setminus \mathcal{D}$, and intersects the smooth part of all irreducible components of \mathcal{C} .

For brevity, we simply denote by $[\gamma]$ the image of $\widetilde{\gamma}$ in $H_1(\mathbb{CP}^2 \setminus \mathcal{D})/\mathcal{J}_c$. We also denote by \mathcal{I}_c the image of $\bigoplus_{C \in Irr(\mathcal{C})} H_1(C \setminus \mathcal{D})$ by j_* , composed with the projection map $H_1(\mathbb{CP}^2 \setminus \mathcal{D}) \to H_1(\mathbb{CP}^2 \setminus \mathcal{D})/\mathcal{J}_c$.

Definition 1.2. The oriented linking of C with D along γ , denoted by $lk_{\gamma}(C, D)$, is the coset of \mathcal{I}_{C} in $H_1(\mathbb{CP}^2 \setminus D)/\mathcal{J}_{C}$ with respect to $[\gamma]$. In other words,

$$\mathrm{lk}_{\gamma}(\mathcal{C},\mathcal{D}) = [\gamma]\mathcal{I}_{\mathcal{C}} \subset \mathrm{H}_{1}(\mathbb{C}\mathbb{P}^{2} \setminus \mathcal{D})/\mathcal{J}_{\mathcal{C}}.$$

Theorem 1.3. The above formula is well-defined, i.e., $lk_{\gamma}(\mathcal{C}, \mathcal{D})$ does not depend on the choice of topological lift of γ . *Proof.* Let $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ be two topological lifts of γ , and let $[\gamma]_1$ and $[\gamma]_2$ denote their homology classes in $H_1(\mathbb{CP}^2 \setminus \mathcal{D})$. There are essentially two ways in which $[\gamma]_1$ and $[\gamma]_2$ may differ. If $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ have same homology class in $H_1(\mathcal{C})$, then they differ by elements of $\bigcup_{C \in Irr(\mathcal{C})} \partial(C \setminus \mathcal{D})$, so that $[\gamma]_1$ and $[\gamma]_2$ differ by an element of the indeterminacy subgroup \mathcal{J}_C . Now, if $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ have different homology classes in $H_1(\mathcal{C})$, then the difference is mapped in \mathcal{I}_C by j_* , so that $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ yield the same coset of \mathcal{I}_C in $H_1(\mathbb{CP}^2 \setminus \mathcal{D})/\mathcal{J}_C$. \Box

Remark 1.4. We stress that the neither of the two assumptions made here, that γ is maximal and that it avoids \mathcal{D} , is necessary to define our invariant – this is discussed in Remark 1.7 and in Section 1.3.2 below. But, on one hand, these assumptions turn out to greatly simplify the exposition and, on the other hand, all the relevant topological information on $\mathcal{C} \cup \mathcal{D}$ are already essentially detected by this simple version of our invariant. As a matter of fact, all the examples of this paper will involve the above assumptions.

We have the following description of $\mathcal{J}_{\mathcal{C}}$.

Proposition 1.5. The indeterminacy subgroup $\mathcal{J}_{\mathcal{C}}$ is spanned by the elements of *the form:*

$$\sum_{d \in \mathcal{D}(P)} I_P(b,d).m_{\beta_P(d)}, \text{ for all } P \in \mathcal{C} \cap \mathcal{D} \text{ and all } b \in \mathcal{C}(P).$$

where $I_P(b,d)$ denotes the intersection multiplicity of the local branches b and d at P, and $m_{\beta_P(d)}$ is given by a meridian of the irreducible component $\beta_P(d)$ of \mathcal{D} containing d.

Proof. The indeterminacy subgroup is the image of $\bigoplus_{C \in Irr(\mathcal{C})} H_1(\partial(C \setminus \mathcal{D}))$ in $H_1(\mathbb{CP}^2 \setminus \mathcal{D})$. It is thus generated by the class of the cycles in $C \in Irr(\mathcal{C})$ around the points $P \in \mathcal{C} \cap \mathcal{D}$. Pick such a singular point P, and consider a small sphere S around P. Each local branch b of C at P intersects S along a knot K_b , and it is well-known that, for each local branch d of \mathcal{D} at P, the intersection of $b \cup d$ with S is an oriented two-component link whose linking number is precisely I(b,d) (see [BKS, pp. 439]). Hence the homology class of the knot K_b in $H_1(\mathbb{CP}^2 \setminus \mathcal{D})$ is given by $\sum_{d \in \mathcal{D}(P)} I(b,d).m_{\beta_P(d)}$, and the result follows. \Box

As a consequence of Proposition 1.5, the group $H_1(\mathbb{CP}^2 \setminus \mathcal{D})/\mathcal{J}_{\mathcal{C}}$ is determined by the combinatorics of the curve $\mathcal{C} \cup \mathcal{D}$. So we can use the linking invariant to compare the topology of curves with the same combinatorics. Indeed, we have the following theorem, which implies that the linking invariant is an invariant of the oriented topology of $(\mathbb{CP}^2, \mathcal{C} \cup \mathcal{D})$. **Theorem 1.6.** Let ϕ be an orientation-preserving homeomorphism between two algebraic curves $C_1 \cup D_1$ and $C_2 \cup D_2$. Then ϕ induces an isomorphism ϕ_* between $H_1(\mathbb{CP}^2 \setminus D_1)$ and $H_1(\mathbb{CP}^2 \setminus D_2)$ mapping \mathcal{J}_{C_1} to \mathcal{J}_{C_2} , and for any cycle $\gamma_1 \in \Gamma_{C_1}$ avoiding \mathcal{D}_1 , we have

$$\overline{\phi_*}\big(\mathrm{lk}_{\gamma_1}(\mathcal{C}_1,\mathcal{D}_1)\big) = \mathrm{lk}_{\phi_{\Gamma}(\gamma_1)}(\mathcal{C}_2,\mathcal{D}_2),$$

where $\overline{\phi_*}$ is map induced by ϕ_* on the quotients by the indeterminacy subgroups.

Proof. By definition, the homeomorphism $\phi : \mathbb{CP}^2 \to \mathbb{CP}^2$ maps \mathcal{D}_1 to \mathcal{D}_2 , so it induces an isomorphism ϕ_* between $H_1(\mathbb{CP}^2 \setminus \mathcal{D}_1)$ and $H_1(\mathbb{CP}^2 \setminus \mathcal{D}_2)$. Furthermore, for each $C_1 \in \operatorname{Irr}(\mathcal{C}_1)$ with image $C_2 = \phi(C_1) \in \operatorname{Irr}(\mathcal{C}_2)$, we have that ϕ maps $C_1 \cap \mathcal{D}_1$ to $C_2 \cap \mathcal{D}_2$, and maps $\partial(C_1 \setminus \mathcal{D}_1)$ to $\partial(C_2 \setminus \mathcal{D}_2)$; this implies that ϕ_* maps \mathcal{J}_{C_1} to \mathcal{J}_{C_2} . Now, ϕ maps any (oriented) lift of γ_1 to a cycle on \mathcal{C}_2 which is a lift of $\phi_{\Gamma}(\gamma_1)$, respecting the orientation. Since the linking invariant does not depend on the choice of lift, the result follows. \Box

Remark 1.7. As mentioned in Remark 1.4, the cycle γ in Definition 1.2 does not need be maximal. Indeed, if Γ does not contain all component-vertices of $\Gamma_{\mathcal{C}}$, then the coset $[\gamma]\mathcal{I}_{\mathcal{C}}$ still yields an invariant of the oriented topology of $(\mathbb{CP}^2, \mathcal{C} \cup \mathcal{D})$. But in this case a finer invariant is given by regarding the curve $\mathcal{C} \cup \mathcal{D}$ as decomposed into the union of \mathcal{C}_{γ} and $(\mathcal{C} \cup D) \setminus \mathcal{C}_{\gamma}$, where \mathcal{C}_{γ} denotes the union of all irreducible components of \mathcal{C} intersecting γ .

Remark 1.8. The linking invariant of Definition 1.2 is an invariant of the oriented topology of $(\mathbb{CP}^2, \mathcal{C} \cup \mathcal{D})$. If γ is simply connected, however, the linking $lk_{\gamma}(\mathcal{C}, \mathcal{D})$ of \mathcal{C} with \mathcal{D} along γ is a topological invariant of $(\mathbb{CP}^2, \mathcal{C} \cup \mathcal{D})$, since any choice of orientation of a topological lift yields the same coset. In general, we can easily remove the condition of orientation, simply by considering

$$-[\gamma]\mathcal{I}_{\mathcal{C}} \cup [\gamma]\mathcal{I}_{\mathcal{C}} \subset \mathrm{H}_{1}(\mathbb{C}\mathbb{P}^{2} \setminus \mathcal{D})/\mathcal{J}_{\mathcal{C}},$$

which is clearly an invariant that doesn't depend on the orientation of γ , but only on its combinatorics. As a corollary to Theorem 1.6, this *non-oriented linking* is a topological invariant of the pair ($\mathbb{CP}^2, \mathcal{C} \cup \mathcal{D}$).

1.3. Two variants. We now discuss two variants of our linking invariant. The first one is a 'global' version which doesn't rely upon the choice of a cycle; the second one is a generalization, where we allow arbitrary cycles.

As in the previous section, $C \cup D$ will denote here a reducible algebraic curve decomposed into two subcurves C and D.

1.3.1. Global linking. We can define the following coarser invariant, which is a 'global' version of the linking invariant, in the sense that it doesn't involve the choice of a cycle. Recall that j_* is the map induced in homology by the inclusion of $\mathcal{C} \setminus \mathcal{D}$ in $\mathbb{CP}^2 \setminus \mathcal{D}$.

Definition 1.9. The *global linking* of C with D, denoted by $\mathcal{L}(C, D)$, is the class of Im j_* in $H_1(\mathbb{CP}^2 \setminus D)/\mathcal{J}_C$.

Remark that the global linking of C with D can also be defined as the union, over all cycles γ in Γ_{C} avoiding D, of the linking invariants of C with D along γ .

The invariance of the global linking is a direct consequence of the proof of Theorem 1.6:

Theorem 1.10. Let ϕ be a homeomorphism between two curves $C_1 \cup D_1$ and $C_2 \cup D_2$. We have

$$\overline{\phi_*}(\mathcal{L}(\mathcal{C}_1,\mathcal{D}_1))=\mathcal{L}(\mathcal{C}_2,\mathcal{D}_2),$$

where $\overline{\phi_*}$ is the map induced by ϕ_* on the quotients by the indeterminacy subgroups.

Hence the global linking of C with D is a topological invariant of the pair $(\mathbb{CP}^2, C \cup D)$.

Remark 1.11. Notice that if C is an irreducible curve, then $lk_*(C, D) = \mathcal{L}(C, D)$, where * denotes the unique component-vertex of Γ_C .

1.3.2. Linking along an arbitrary cycle. In the above definition of the linking invariant, we assumed throughout that the cycle γ avoids all point-vertices of $\Gamma_{\mathcal{C}}$ corresponding to singularities in $\mathcal{C} \cap \mathcal{D}$. Although this will not be used in the main examples of this paper, we outline here how the construction can be easily generalized to arbitrary cycles.

Let γ be any cycle in $\Gamma_{\mathcal{C}}$. Denote by S_{γ} the set of singularities in $\mathcal{C} \cap \mathcal{D}$ whose corresponding point-vertices are contained in γ . We define the subgroup \mathcal{J}_{γ} of $H_1(\mathbb{CP}^2 \setminus \mathcal{D})$ as

 $\mathcal{J}_{\gamma} = \langle m_{\beta_P(d)}, \text{ for all } P \in \mathcal{S}_{\gamma} \text{ and all } d \in \mathcal{D}(P) \rangle.$

(here we make use of the same notation as in Proposition 1.5.)

Definition 1.12. The γ -indeterminacy subgroup is the subgroup $\mathcal{J}_{(\mathcal{C},\gamma)}$ of $H_1(\mathbb{CP}^2 \setminus \mathcal{D})$ generated by $\mathcal{J}_{\mathcal{C}} \cup \mathcal{J}_{\gamma}$.

Now, we need a slightly generalized notion of topological lift for the cycle γ . Specifically, for each singular point P in S_{γ} , pick a small closed 4-ball \mathbb{B}_P centered at P. A \mathcal{D} -avoiding lift of γ is a cycle in $\mathbb{CP}^2 \setminus \mathcal{D}$ which coincides with a topological lift outside $\cup_{P \in S_{\gamma}} \mathbb{B}_P$ (and in particular lies in \mathcal{C}), and whose intersection with each 4-ball \mathbb{B}_P is an arc lying in the boundary of \mathbb{B}_P . So, roughly speaking, such a cycle differs from a topological lift of γ by locally pushing it away from the curve $\mathcal{C} \cup \mathcal{D}$, so that it avoids the singularities in S_{γ} .

Now, using this refined indeterminacy subgroup and generalized notion of lift, the exact same construction yields an invariant: the (oriented) linking of C with D along an arbitrary cycle γ is the coset

$$\mathrm{lk}_{\gamma}(\mathcal{C},\mathcal{D}) = [\gamma]\mathcal{I}_{\mathcal{C}} \subset \mathrm{H}_{1}(\mathbb{CP}^{2} \setminus \mathcal{D})/\mathcal{J}_{(\mathcal{C},\gamma)},$$

where $[\gamma]$ denotes the image of a \mathcal{D} -avoiding lift of γ in $H_1(\mathbb{CP}^2 \setminus \mathcal{D})/\mathcal{J}_{(\mathcal{C},\gamma)}$, and where $\mathcal{I}_{\mathcal{C}}$ is the image of $\bigoplus_{C \in Irr(\mathcal{C})} H_1(C \setminus \mathcal{D})$ by j_* , seen in $H_1(\mathbb{CP}^2 \setminus \mathcal{D})/\mathcal{J}_{(\mathcal{C},\gamma)}$.

The proof that this is well-defined, i.e., does not depend on the choice of the \mathcal{D} -avoiding lift of γ , is completely similar to the proof of Theorem 1.3, and readily follows from the definition of $\mathcal{J}_{(\mathcal{C},\gamma)}$. The only difference here is that, if γ passes through a point-vertex P in $\mathcal{C} \cap \mathcal{D}$, then two \mathcal{D} -avoiding lifts of γ may only differ by a copy of a meridian $m_{\beta_P(d)}$, for any local branch d in $\mathcal{D}(P)$. Considering these cycles in $H_1(\mathbb{CP}^2 \setminus \mathcal{D})$, we have by definition that the difference lies precisely in \mathcal{J}_{γ} , hence in the γ -indeterminacy subgroup $\mathcal{J}_{(\mathcal{C},\gamma)}$.

Remark 1.13. In the case where γ avoids all point-vertices in $C \cap D$, then the γ -indeterminacy subgroup $\mathcal{J}_{(C,\gamma)}$ coincides with the original indeterminacy subgroup \mathcal{J}_C , and we recover the invariant of Definition 1.2.

1.4. Line arrangements and the \mathcal{I}-invariant. In this section, we restrict ourselves to the case where $\mathcal{C} \cup \mathcal{D}$ is a line arrangement, i.e., when all the irreducible components of \mathcal{C} and \mathcal{D} are of degree 1. In this particular case, another linking invariant, called the \mathcal{I} -invariant, has been defined by Artal, Florens and the first author in [AFGB]. We will prove here that the present linking invariant generalizes the \mathcal{I} -invariant.

First, let us recall some terminologies introduced in [AFGB] to define the \mathcal{I} -invariant. Let \mathcal{A} be a line arrangement. Recall that $H_1(\mathbb{CP}^2 \setminus \mathcal{A})$ is free of rank $|\operatorname{Irr}(\mathcal{A})| - 1$ and is generated by the set of all meridiens m_L for $L \in \operatorname{Irr}(\mathcal{A})$.

Definition 1.14. Let \mathcal{A} be a line arrangement, $\xi : H_1(\mathbb{CP}^2 \setminus \mathcal{A}) \to \mathbb{C}^*$ be a non-trivial character and γ be a cycle of the intersection graph $\Gamma_{\mathcal{A}}$. The triple $(\mathcal{A}, \xi, \gamma)$ is called an inner-cyclic arrangement if

(1) For each singular point P of A with associated point-vertex in γ ,

 $\xi(m_{\ell}) = 1$, for any line ℓ of \mathcal{A} containing P,

- (2) For each line L of A with associated component-vertex in γ ,
 - (i) $\xi(m_L) = 1$,
 - (ii) $\prod_{\ell \ni S} \xi(m_\ell) = 1$, for any singular point S in L, where the product runs over all the lines ℓ in \mathcal{A} containing S.

In [AFGB], the \mathcal{I} -invariant is then defined as

$$\mathcal{I}(\mathcal{A},\xi,\gamma)=\xi\circ\iota_*(\bar{\gamma}),$$

where ι_* is the map induced by the inclusion ι of the boundary $B_{\mathcal{A}}$ of a tubular neighbourhood Tub(\mathcal{A}) of \mathcal{A} in $\mathbb{CP}^2 \setminus \mathcal{A}$, and where $\bar{\gamma}$ is a suitably chosen lift of the cycle γ in $B_{\mathcal{A}}$. More precisely, this lift is a 'nearby cycle' in the terminology of [AFGB], which roughly means that this cycle is contained in $B_{\mathcal{C}} \setminus \text{Tub}(\mathcal{D}) \subset \mathbb{CP}^2 \setminus \mathcal{D}$, where $\mathcal{A} = \mathcal{C} \cup \mathcal{D}$ with $\text{Irr}(\mathcal{C}) = \{L \in \text{Irr}(\mathcal{A}) | L \cap \gamma \neq \emptyset\}$ and $\text{Irr}(\mathcal{D}) = \{L \in \text{Irr}(\mathcal{A}) | L \cap \gamma = \emptyset\} - \text{see}$ [AFGB, Def. 2.11] for a precise definition.

Now, if $\mathcal{A} = \mathcal{C} \cup \mathcal{D}$ is a line arrangement, the set $\mathcal{I}_{\mathcal{C}}$ is always trivial. The linking of \mathcal{C} and \mathcal{D} along a cycle γ is thus the class $[\gamma]$ of a lift of γ in $H_1(\mathbb{CP}^2 \setminus \mathcal{D})/\mathcal{J}_{\mathcal{C}}$. The relationship to the \mathcal{I} -invariant is as follows:

Theorem 1.15. Let $\mathcal{A} = \mathcal{C} \cup \mathcal{D}$ be a line arrangement, and let γ be a maximal cycle in $\Gamma_{\mathcal{C}}$. Let ξ be a character on $H_1(\mathbb{CP}^2 \setminus \mathcal{A})$ such that $(\mathcal{A}, \xi, \gamma)$ is an innercyclic arrangement. Then there is a nontrivial character ξ_* on $H_1(\mathbb{CP}^2 \setminus \mathcal{D})/\mathcal{J}_{(\mathcal{C},\gamma)}$ induced by ξ such that

$$\mathcal{I}(\mathcal{A},\xi,\gamma) = \xi_* \big(\mathrm{lk}_{\gamma}(\mathcal{C},\mathcal{D}) \big).$$

Proof. Since $(\mathcal{A}, \xi, \gamma)$ is an inner-cyclic arrangement, we have that $\xi(m_L) = 1$ for each line L of \mathcal{A} with corresponding component-vertex in γ – these correspond to the lines of \mathcal{C} since γ is maximal. This shows that ξ factors through the projection map $H_1(\mathbb{CP}^2 \setminus \mathcal{A}) \to H_1(\mathbb{CP}^2 \setminus \mathcal{D})$. Furthermore, conditions (1) and (2-ii) of Definition 1.14 ensures that ξ further factors to $H_1(\mathbb{CP}^2 \setminus \mathcal{D})/\mathcal{J}_{(\mathcal{C},\gamma)}$, thus providing the desired nontrivial character ξ_* .¹ So we have

$$\mathcal{I}(\mathcal{A},\xi,\gamma) = \xi \circ \iota_*(\bar{\gamma}) = \xi_*([\bar{\gamma}]),$$

where $\bar{\gamma}$ is any lift of γ which is a nearby cycle, and $[\bar{\gamma}]$ denotes its image in $H_1(\mathbb{CP}^2 \setminus D)/\mathcal{J}_{(\mathcal{C},\gamma)}$. On the other hand, we have by definition that

$$\mathrm{lk}_{\gamma}(\mathcal{C},\mathcal{D}) = [\widetilde{\gamma}] \in \mathrm{H}_{1}(\mathbb{CP}^{2} \setminus \mathcal{D}) / \mathcal{J}_{(\mathcal{C},\gamma)},$$

¹ This shows, in particular, that the quotient $H_1(\mathbb{CP}^2 \setminus \mathcal{D})/\mathcal{J}_{(\mathcal{C},\gamma)}$ is nontrivial.

where $\tilde{\gamma}$ is a \mathcal{D} -avoiding lift of γ . The result then follows from the fact that the homology classes of the cycles $\bar{\gamma}$ and $\tilde{\gamma}$ in $H_1(\mathbb{CP}^2 \setminus \mathcal{D})$ can only differ by elements of $\mathcal{J}_{(\mathcal{C},\gamma)}$, as follows from the definition of a nearby cycle given in [AFGB].

2. Computations

In this section, we describe two concrete methods for computing our invariant. The first one is topological and is based on a modification of the braid monodromy, and uses the usual linking number of links in the 3-sphere. The second one is an algebraic method using the connected numbers introduced by Shirane in [Shir2], and its relations with the linking invariant, observed in [GBS].

2.1. Topological method. For simplicity, we consider an algebraic curve $C \cup D$, such that all the irreductible components of C have topological genus zero (see however Remark 2.3), together with a cycle γ in the intersection graph of C. For this class of curves, we have that the set \mathcal{I}_C is trivial, so that the linking invariant $lk_{\gamma}(C, D)$ is the class of a lift of γ in $H_1(\mathbb{CP}^2 \setminus D)/\mathcal{J}_C$, rather than a coset (this was already observed in the case of line arrangements in Section 1.4).

Definition 2.1. A path $\widetilde{\gamma}$ in C is γ -admissible if it is a lift of γ and if there is a generic projection $\pi : \mathbb{CP}^2 \setminus \{*\} \to \mathbb{CP}^1$ such that $\pi(\widetilde{\gamma})$ has no self-intersection, and $(\pi^{-1} \circ \pi(\widetilde{\gamma})) \cap \operatorname{Sing}(\mathcal{D}) = \emptyset$.

Note that the latter condition can always be fulfilled, up to a small modification of π .

Let $\tilde{\gamma}$ be a γ -admissible path in C. For any point p of $\pi(\tilde{\gamma})$, we consider the fiber F_p over p. By the definition of γ -admissibility, the number of intersection points of D with F_p equals the degree of D for all points $p \in \pi(\tilde{\gamma})$. We denote by L_D the oriented link

(1)
$$L_{\mathcal{D}} = \left(\pi^{-1} \circ \pi(\widetilde{\gamma})\right) \cap \mathcal{D} \subset \mathbb{CP}^2.$$

Noting that $\tilde{\gamma}$ and $L_{\mathcal{D}}$ do not intersect, we define $\tilde{\gamma} \cup L_{\mathcal{D}}$. This link naturally sits in a copy of S^3 , as follows. Let D be the disc bounded by $\pi(\tilde{\gamma})$ in \mathbb{CP}^1 . Pick a polydisc \mathcal{P} of \mathbb{CP}^2 such that $\pi(\mathcal{P}) = D$ and $\pi^{-1}(D) \cap (\mathcal{C} \cup \mathcal{D}) \subset \mathcal{P}$. By construction, the link $\tilde{\gamma} \cup L_{\mathcal{D}}$ lies in the boundary of \mathcal{P} , which is homeomorphic to S^3 . This construction yields the following.

Theorem 2.2. Under the above assumptions, the homology class of the path $\tilde{\gamma}$ in $H_1(\mathbb{CP}^2 \setminus D)$ is given by:

$$\rho\left(\sum_{c\in L_{\mathcal{D}}}\ell(c,\,\widetilde{\gamma}\,).m_c\right),\,$$

where ℓ is the usual linking number in S^3 and ρ is the map induced by the inclusion of $S^3 \setminus L_D$ in $\mathbb{CP}^2 \setminus D$, which maps the meridian m_c of each component c of L_D to the meridian of the irreducible component of D containing c.

Remark 2.3. If the irreducible components of the curve C are allowed to have nonzero topological genus, we can also use the present method to compute the value of the generators of \mathcal{I}_C in $H_1(\mathbb{CP}^2 \setminus \mathcal{D})$, and thus to compute the coset $[\gamma]\mathcal{I}_C$.

This provides a computational formula for the linking invariant of C with D along γ in terms of the usual linking number. See Section 3.2 for an application on a concrete example.

2.2. Algebraic method. The second method of computation comes from the connected numbers introduced by Shirane in [Shir2]. This method applies when C is a nodal curve with $\operatorname{Sing}(C) \cap D = \emptyset$ (this implies that $C \setminus D$ is connected), and if $\operatorname{H}_1(\mathbb{CP}^2 \setminus D)/\mathcal{J}_C \simeq \mathbb{Z}/m\mathbb{Z}$.

Let $\psi : X \to \mathbb{CP}^2$ be a cyclic cover of degree *m* branched over \mathcal{D} . The *connected number* of \mathcal{C} for ψ is the number of connected components of $\psi^{-1}(\mathcal{C} \setminus \mathcal{D})$) in *X*. Based on a previous version of the present paper, it has been proved in [GBS] that if \mathcal{C} and \mathcal{D} are smooth curves then the connected number and the global linking of \mathcal{C} and \mathcal{D} are essentially equivalent. For the purpose of this paper, however, we will rather give the following statement.

Theorem 2.4. Suppose that C is a nodal curve such that $\operatorname{Sing}(\mathcal{C}) \cap \mathcal{D} = \emptyset$, and that $\operatorname{H}_1(\mathbb{CP}^2 \setminus \mathcal{D})/\mathcal{J}_C \simeq \mathbb{Z}/m\mathbb{Z}$. Then the global linking of C and \mathcal{D} is the unique subgroup of $\operatorname{H}_1(\mathbb{CP}^2 \setminus \mathcal{D})/\mathcal{J}_C$ of index $\operatorname{deg}(\mathcal{D})/\mu$, where μ is the minimal degree of a plane curve E such that $C \cap \mathcal{D} = C \cap E$ and for each point $P \in C \cap \mathcal{D}$, we have $(\operatorname{deg}(\mathcal{D})/\mu) \times I_P(C, E) = I_P(C, \mathcal{D})$.

Sketch of proof. By the proof of [GBS, Theorem 2.5], we know that the index of $\mathcal{L}(\mathcal{C},\mathcal{D})$ in $H_1(\mathbb{CP}^2 \setminus \mathcal{D})/\mathcal{J}_{\mathcal{C}}$ is equal to the connected number of \mathcal{C} for ψ . Since $H_1(\mathbb{CP}^2 \setminus \mathcal{D})/\mathcal{J}_{\mathcal{C}}$ is cyclic then each of its subgroup is determined by its index. The result is thus a consequence of [Shir2, Corollary 2.5].

3. Applications

In this section, we use our linking invariants (both the oriented and global versions) to distinguish two types of Zariski pairs.

3.1. 3-Artal curves. As an application of the algebraic method of computation of the linking invariant, we propose to distinguish the Zariski pair found by Artal in [Art]. These curves are formed by a smooth cubic C and three inflexional tangent lines. The geometry of the 9 inflexion points of a cubic is well known; the collinearity relations are the same as in \mathbb{F}_3^2 , the plane over the finite field of 3 elements. We consider P_1, \ldots, P_4 four inflexion points of C such that P_1, P_2, P_3 are collinear and P_1, P_2, P_4 are not. Set $\mathcal{A}_1 = L_1 \cup L_2 \cup L_3$ and $\mathcal{A}_2 = L_1 \cup L_2 \cup L_4$, where L_i denotes the inflexional tangent line at P_i .

Theorem 3.1. The global linking of the line arrangement A_i with the cubic C is:

$$\mathcal{L}(\mathcal{A}_i, \mathcal{C}) = \begin{cases} \{0\} & \text{if } i = 1, \\ \{0, 1, 2\} & \text{if } i = 2. \end{cases}$$

Proof. Since the cubic C is smooth, we have $H_1(\mathbb{CP}^2 \setminus C) \simeq \mathbb{Z}/3\mathbb{Z}$. Furthermore, Proposition 1.5 implies that $\mathcal{J}_{\mathcal{A}_i} = \langle 3m \rangle$, where *m* is a meridian of the cubic. So the quotient $H_1(\mathbb{CP}^2 \setminus C)/\mathcal{J}_{\mathcal{A}_i}$ is also isomorphic to $\mathbb{Z}/3\mathbb{Z}$.

We first compute the global linking $\mathcal{L}(\mathcal{A}_1, \mathcal{C})$. By construction, there is a line E (i.e., an algebraic curve of degree 1) passing through P_1, P_2 and P_3 . By Bezout theorem, we have $I_{P_j}(\mathcal{A}_1, E) = 1$ for $j \in \{1, 2, 3\}$, and the following equality holds for $j \in \{1, 2, 3\}$

$$\frac{\deg(\mathcal{C})}{\deg(E)} \times I_{P_j}(\mathcal{A}_1, E) = I_{P_j}(\mathcal{A}_1, \mathcal{C}).$$

Thus by Theorem 2.4 the index of $\mathcal{L}(\mathcal{A}_1, \mathcal{C})$ in $H_1(\mathbb{CP}^2 \setminus \mathcal{C})/\mathcal{J}_{\mathcal{A}_1}$ is 3.

Let us now turn to $\mathcal{L}(\mathcal{A}_2, \mathcal{C})$, and look for the minimal degree curve E passing through the points P_1, P_2 and P_4 and satisfying $\frac{\deg(\mathcal{C})}{\deg(E)} \times I_{P_j}(\mathcal{A}_2, E) = I_{P_j}(\mathcal{A}_2, \mathcal{C})$ for $j \in \{1, 2, 4\}$. By construction, no line E satisfies these conditions. Similar considerations as above show that no conic can verify these conditions either.² But taking E to be the cubic \mathcal{C} obviously works, and it follows by Theorem 2.4 that the index of $\mathcal{L}(\mathcal{A}_2, \mathcal{C})$ in $H_1(\mathbb{CP}^2 \setminus \mathcal{C})/\mathcal{J}_{\mathcal{A}_2}$ is 1.

Corollary 3.2. The curves $C \cup A_1$ and $C \cup A_2$ form a Zariski pair.

² This also follows from Theorem 2.4, since the existence of such a conic would imply that $\mathcal{L}(\mathcal{A}_2, \mathcal{C})$ is an index 2 subgroup of $\mathbb{Z}/3\mathbb{Z}$.

3.2. The quartic and its bitangents. As an application of the topological method, we will distinguish a Zariski pair formed by a quartic and 3 bitangents. Let Q be the Klein quartic defined by $x^3y + y^3z + z^3x = 0$. The full list of its 28 bitangents is given in [Shi]. We will consider here only four of them. Let ζ be a primitive 7th root of unity, and define the real numbers $\varepsilon_i = \zeta^i + \zeta^{-i}$, for $i \in \{1, 2, 3\}$. We consider the following bitangents:

$$L_1: x + y + z = 0, \qquad L_2: x + \varepsilon_3^{-2}y + \varepsilon_1^2 z = 0, L_3: x + \zeta^3 \varepsilon_3^{-2}y + \zeta \varepsilon_1^2 z = 0, \qquad L_4: x + \zeta^2 \varepsilon_2^{-2}y + \zeta^3 \varepsilon_3^2 z = 0.$$

Let $\mathcal{A}_i = L_1 \cup L_2 \cup L_{2+i}$, for $i \in \{1,2\}$. We will compute the linking of \mathcal{A}_i with \mathcal{Q} along γ_i , where γ_i is a cycle generating $H_1(\mathcal{A}_i) \simeq \mathbb{Z}$. Since $H_1(L_j) = 0$ for all j, we have $\mathcal{I}_{\mathcal{A}_i} = 0$ for i = 1, 2. Furthermore, \mathcal{Q} is smooth so $H_1(\mathbb{CP}^2 \setminus \mathcal{Q}) \simeq \mathbb{Z}/4\mathbb{Z}$. By Proposition 1.5, we have that $\mathcal{J}_{\mathcal{A}_i} = \langle 2m \rangle$, where m is a meridian of \mathcal{Q} . So we have

$$\mathrm{H}_1(\mathbb{CP}^2 \setminus \mathcal{Q}) / \mathcal{J}_{\mathcal{A}_i} \simeq \mathbb{Z}/2\mathbb{Z}.$$

In order to simplify the computations, we apply on $Q \cup A_1$ and $Q \cup A_2$ the linear change of variables given respectively by the following matrices:

$$P_{1} = \begin{pmatrix} 4\xi^{5} + 4\zeta^{4} - \zeta^{2} + \zeta + 6 & -\zeta^{5} - 5\zeta^{4} + 2\zeta^{3} + 6\zeta^{2} + 5 \\ -6\zeta^{5} - 6\zeta^{4} - 2\zeta^{2} - 5\zeta - 2 & 5\zeta^{5} + 4\zeta^{4} - 3\zeta^{3} - 2\zeta^{2} + 7\zeta + 3 \\ 2\zeta^{5} + 2\zeta^{4} + 3\zeta^{2} + 4\zeta + 3 & -4\zeta^{5} + \zeta^{4} + \zeta^{3} - 4\zeta^{2} - 7\zeta - 1 \\ & -3\zeta^{5} + \zeta^{4} - 2\zeta^{3} - 5\zeta^{2} - \zeta + 3 \\ \zeta^{5} + 2\zeta^{4} + 3\zeta^{3} + 4\zeta^{2} - 2\zeta - 1 \\ 2\zeta^{5} - 3\zeta^{4} - \zeta^{3} + \zeta^{2} + 3\zeta - 2 \end{pmatrix},$$

$$P_{2} = \begin{pmatrix} -3\zeta^{5} - 2\zeta^{4} - 4\zeta^{3} - 2\zeta^{2} - 3\zeta + 7 & 8\zeta^{5} - 2\zeta^{4} - 9\zeta^{3} + \zeta^{2} + 9 \\ -6\zeta^{5} + 3\zeta^{4} - \zeta^{3} - 4\zeta^{2} + \zeta & -12\zeta^{5} - 11\zeta^{4} - 4\zeta^{3} - 5\zeta^{2} - 7\zeta - 3 \\ 9\zeta^{5} - \zeta^{4} + 5\zeta^{3} + 6\zeta^{2} + 2\zeta + 7 & 4\zeta^{5} + 13\zeta^{4} + 13\zeta^{3} + 4\zeta^{2} + 7\zeta + 8 \\ & -5\zeta^{5} + 4\zeta^{4} + 13\zeta^{3} + \zeta^{2} + 3\zeta + 12 \\ 18\zeta^{5} + 8\zeta^{4} + 5\zeta^{3} + 9\zeta^{2} + 6\zeta + 3 \\ -13\zeta^{5} - 12\zeta^{4} - 18\zeta^{3} - 10\zeta^{2} - 9\zeta - 15 \end{pmatrix}$$

The change of variables P_i sends L_1 , L_2 and L_{2+i} to x + y = 0, x + z = 0 and y + z = 0 respectively (we denote by $\overline{\mathcal{A}}_i$ the arrangement (x + y)(x + z)(y + z) = 0). The images of \mathcal{Q} by the changes of variables P_i are denoted by $\overline{\mathcal{Q}}^i$, and their equations are given in Appendix A. Since P_i is a linear change of variables, we have

$$\operatorname{lk}_{\gamma_i}(\mathcal{A}_i, \mathcal{Q}) = \operatorname{lk}_{\gamma_i}(\overline{\mathcal{A}}_i, \overline{\mathcal{Q}}^\iota).$$

By Theorem 2.2, the linking invariant $lk_{\gamma_i}(\overline{\mathcal{A}}_i, \overline{\mathcal{Q}}^i)$ is given by the linking numbers of the cycle γ_i with each component of the link $L_{\overline{\mathcal{Q}}^i}$ defined in (1). We can take as topological cycle $\tilde{\gamma}_1$ (resp. $\tilde{\gamma}_2$) the triangle formed by the lines x + y = 0, x + z = 0 and y + z = 0 in the chart z = 1 (resp. y = 1) of \mathbb{RP}^2 . These charts together with the cycles $\tilde{\gamma}_i$ are represented in Figure 1.

The link $L_{\overline{Q}^i}$ lives in the intersection of \overline{Q}^i with the boundary of the polydisk $\mathcal{P} = \mathcal{T} \times D^2$, where \mathcal{T} is the disk of the real plane bounded by $\tilde{\gamma}_i$ and where D^2 is a 2-disk intersecting all the components of \overline{Q}^i (see Section 2.1). The desired linking number can be computed as the algebraic intersection number of $L_{\overline{Q}^i}$ with this disk \mathcal{T} . Since we are working here with $(\mathbb{Z}/2\mathbb{Z})$ -coefficients, this amounts to counting the parity of the number of intersection points.

In order to compute these numbers of intersection points between the interior of the triangle \mathcal{T} and the quartics $\overline{\mathcal{Q}}^i$, we proceed as follows. First of all, we choose an embedding of ζ in the field of complex number. In the following, we take $\zeta = \exp\left(\frac{2I\pi}{7}\right) \simeq 0.62349 + 0.78183i$. Then, we decompose $\overline{\mathcal{Q}}^i$ into two quartics $\overline{\mathcal{Q}}_{Re}^i$ and $\overline{\mathcal{Q}}_{Im}^i$, obtained from $\overline{\mathcal{Q}}^i$ by taking only the real part (resp. the imaginary part) of its coefficients. Using the previous approximation of ζ , we have the following approximations

$$\begin{aligned} \overline{\mathcal{Q}}_{Re}^{1} \simeq &-838.65x^{4} + 1903.1x^{3}y + 2540.3x^{2}y^{2} + 2686.2xy^{3} - 4017.2y^{4} + \\ & 1074.6x^{3}z - 2073.7x^{2}yz - 10166.xy^{2}z + 28829.y^{3}z + 7382.2x^{2}z^{2} - \\ & 13500.xyz^{2} - 14908.y^{2}z^{2} + 5379.5xz^{3} - 3835.5yz^{3} - 456.29z^{4}, \end{aligned}$$

$$\overline{\mathcal{Q}}_{Im}^{1} \simeq &-2525.0x^{3}y + 5338.2x^{2}y^{2} - 9450.5xy^{3} + 12939.y^{4} - 2525.0x^{3}z + \\ & 5824.0x^{2}yz - 4948.5xy^{2}z - 5115.5y^{3}z + 485.83x^{2}z^{2} + 8936.1xyz^{2} - \\ & 17722.y^{2}z^{2} + 4434.2xz^{3} + 2330.9yz^{3} + 1999.1z^{4}, \end{aligned}$$

$$\begin{split} \overline{\mathcal{Q}}_{Re}^2 \simeq& 2327.4x^4 + 28631.x^3y + 105760.x^2y^2 + 115460.xy^3 + 28853.y^4 - \\ & 22196.x^3z - 114530.x^2yz - 79537.xy^2z + 22480.y^3z + 9417.8x^2z^2 - \\ & 107220.xyz^2 - 116350.y^2z^2 + 64365.xz^3 + 62192.yz^3 + 348.69z^4, \\ \overline{\mathcal{Q}}_{Im}^2 \simeq& 518.62x^4 - 5686.3x^3y - 56227.x^2y^2 - 101110.xy^3 - 35790.y^4 + \\ & 10085.x^3z + 133160.x^2yz + 275020.xy^2z + 106780.y^3z - 71462.x^2z^2 - \\ & 187780.xyz^2 - 76906.y^2z^2 + 9523.7xz^3 - 961.76yz^3 + 838.07z^4. \end{split}$$

Thus, the real points of $\overline{\mathcal{Q}}^i$ correspond to those points on which both $\overline{\mathcal{Q}}_{Re}^i$ and $\overline{\mathcal{Q}}_{Im}^i$ vanish; in other words, they correspond to the intersection points of these two quartics considered in the real plane. Therefore, we count how many of these real points lie in the disk \mathcal{T} , since by definition, they are the intersection points of $L_{\overline{\mathcal{Q}}^i}$ with \mathcal{T} . Using the previous equations, for each i, we depicts (see



FIGURE 1 Real part of $\overline{Q}^i \cup \overline{A}_i$ (\overline{A}_i solid line, \overline{Q}^i_{Re} dashed line, and \overline{Q}^i_{Im} dotted line)

Figure 1) in the real plan the arrangement $\overline{\mathcal{A}}_i$ (solid line) and the quartics $\overline{\mathcal{Q}}_{Re}^i$ (dashed line) and $\overline{\mathcal{Q}}_{Im}^i$ (dotted line). In this way, we get that the value of $\tilde{\gamma}_i$ in $H_1(\mathbb{CP}^2 \setminus \overline{\mathcal{Q}}^i)/\mathcal{J}_{\overline{\mathcal{A}}_i}$ is

$$\rho(\tilde{\gamma}_i) = \begin{cases} 0 & \text{if } i = 1, \\ 1 & \text{if } i = 2. \end{cases}$$

From these computations and using Remark 1.8, we have the following theorem and its corollary.

Theorem 3.3. The linking of the line arrangement A_i with the quartic Q along γ_i is

$$lk_{\gamma_i}(\mathcal{A}_i, \mathcal{Q}) = \begin{cases} \{0\} & \text{if } i = 1, \\ \{1\} & \text{if } i = 2. \end{cases}$$

Corollary 3.4. The curves $Q \cup A_1$ and $Q \cup A_2$ form a Zariski pair.

Remark 3.5. Since γ_i generates $H_1(A_i)$, we can also compute the global linking of A_i with Q. It is given by:

$$\mathcal{L}(\mathcal{A}_i, \mathcal{Q}) = \begin{cases} \{0\} & \text{if } i = 1, \\ \{0, 1\} & \text{if } i = 2. \end{cases}$$

Remark 3.6. This result is in adequation with the computation of the connected numbers of these curves made in [BTY].

Acknowledgments. The authors would like to thanks E. Artal for seminal discussions at an early stage of this work. They also thank S. Bannai, V. Florens, P. Popescu-Pampu, T. Shirane, H. Tokunaga, as well as the anonymous referee, for their questions and remarks.

A. Explicit equations of the quartics \overline{Q}^i

The equations of the quartics \overline{Q}^i introduced in Section 3.2 are given by:

$$\begin{split} \overline{\mathcal{Q}}^1 : (343\xi^5 + 343\xi^2 - 686)x^4 + \\ (-2058\xi^5 + 4802\xi^4 + 2058\xi^3 - 4802\xi^2 + 1715\xi + 5488)x^3y + \\ (-1029\xi^5 - 26754\xi^4 - 14406\xi^3 + 7203\xi^2 - 10290\xi - 26754)x^2y^2 + \\ (-1029\xi^5 + 37044\xi^4 + 17493\xi^3 - 17150\xi^2 + 18865\xi + 36015)xy^3 + \\ (-2401\xi^5 - 19551\xi^4 - 4802\xi^3 + 10633\xi^2 - 7889\xi - 19208)y^4 + \\ (-2058\xi^5 + 7546\xi^4 + 4802\xi^3 - 4802\xi^2 + 1715\xi + 9604)x^3z + \\ (-2058\xi^5 - 20580\xi^4 - 6174\xi^3 + 8232\xi^2 - 13377\xi - 16464)x^2yz + \\ (27783\xi^5 + 14406\xi^4 - 3087\xi^3 + 16464\xi^2 + 17493\xi - 1029)xy^2z + \\ (-18865\xi^5 - 1372\xi^4 - 343\xi^3 - 19894\xi^2 - 5831\xi + 22295)y^3z + \\ (-11319\xi^5 + 2058\xi^4 + 4116\xi^3 - 9261\xi^2 - 3087\xi + 10290)x^2z^2 + \\ (3087\xi^5 - 10290\xi^4 - 3087\xi^3 + 12348\xi^2 - 4116\xi - 19551)xyz^2 + \\ (14406\xi^5 + 14406\xi^4 - 1029\xi^3 - 1029\xi^2 + 5145\xi - 3087)y^2z^2 + \\ (-6517\xi^5 - 4116\xi^4 + 1029\xi^3 - 2058\xi^2 - 2744\xi + 2401)xz^3 + \\ (1715\xi^5 + 4116\xi^4 + 3773\xi^3 + 2058\xi^2 + 2744\xi + 2401)yz^3 + \\ (-1715\xi^4 - 343\xi^3 + 1715\xi^2 - 343\xi - 1715)z^4 = 0. \\ \overline{\mathcal{Q}}^2 : (13720\xi^5 - 6174\xi^4 + 16121\xi^3 + 1029\xi^2 + 4116\xi + 12005)x^4 + \\ (34300\xi^5 - 19208\xi^4 - 51450\xi^3 - 129654\xi^2 - 30870\xi - 32928)x^2y^2 + \\ (-94668\xi^5 - 111132\xi^4 - 107016\xi^3 - 127596\xi^2 - 90552\xi - 74088)xy^3 + \\ (686\xi^5 - 49735\xi^4 - 56595\xi^3 - 15092\xi^2 - 21295\xi - 56252)y^4 + \\ (-94668\xi^5 + 10976\xi^4 - 26068\xi^3 - 42532\xi^2 - 31556\xi - 46648)x^3z + \\ (-49392\xi^5 + 49392\xi^4 - 78204\xi^3 + 176988\xi^2 - 41160\xi - 86436)x^2yz + \\ (139944\xi^5 + 209916\xi^4 + 10290\xi^3 + 2582\xi^2 + 187964\xi + 238728)y^3z + \\ (5896\xi^5 + 214032\xi^4 + 127596\xi^3 + 56252\xi^2 + 187964\xi + 238728)y^3z + \\ (5896\xi^5 + 214032\xi^4 + 127596\xi^3 + 56252\xi^2 + 187964\xi + 238728)y^3z + \\ (5896\xi^5 + 214032\xi^4 + 127596\xi^3 + 56252\xi^2 + 187964\xi + 238728)y^3z + \\ (5896\xi^5 + 214032\xi^4 + 127596\xi^3 + 56252\xi^2 + 187964\xi + 238728)y^3z + \\ (5896\xi^5 + 214032\xi^4 + 127596\xi^3 + 56252\xi^2 + 187964\xi + 238728)y^3z + \\ (5896\xi^5 + 214032\xi^4 + 127596\xi^3 + 56252\xi^2 + 187964\xi + 238728)y^3z + \\ (5896\xi^5 + 214032\xi^4 + 127596\xi^3 + 56252\xi^2 + 187964\xi + 238728)y^3z + \\ (5896\xi^5 + 214032\xi^4 + 127596\xi^3 + 56252\xi^2 + 187964\xi + 238728)y^3z + \\ (58$$

$$(104958\xi^{5} + 98784\xi^{4} + 69972\xi^{3} + 32928\xi^{2} + 14406\xi + 183162)x^{2}z^{2} + (-8232\xi^{5} - 176988\xi^{4} + 107016\xi^{3} - 152292\xi^{2} - 218148\xi - 69972)xyz^{2} + (-80262\xi^{5} - 183162\xi^{4} + 12348\xi^{3} - 67914\xi^{2} - 222264\xi - 164640)y^{2}z^{2} + (23324\xi^{5} + 37044\xi^{4} - 32928\xi^{3} - 58996\xi^{2} + 153664\xi - 35672)xz^{3} + (-78204\xi^{5} - 117992\xi^{4} - 93296\xi^{3} - 16464\xi^{2} - 91924\xi - 91924)yz^{3} + (117992\xi^{5} + 114219\xi^{4} - 4116\xi^{3} + 79919\xi^{2} + 114219\xi + 72373)z^{4} = 0.$$

References

- [ACT] E. ARTAL, J. I. COGOLLUDO and H.-O. TOKUNAGA, A survey on Zariski pairs. In Algebraic Geometry in East Asia–Hanoi 2005, vol. 50 of Adv. Stud. Pure Math. Math. Soc. Japan, Tokyo (2008). pp. 1–100. Zbl1141.14015 MR 2409555
- [AFGB] E. ARTAL, V. FLORENS and B. GUERVILLE-BALLÉ, A new topological invariant of line arrangements. Ann. Sc. Norm. Super. Pisa Cl. Sci. XVII (2017), 1–20.
- [Art] E. ARTAL, Sur les couples de Zariski. J. Algebraic Geom. 3 (1994), 223–247.
 Zbl 0823.14013 MR 1257321
- [BGBS] S. BANNAI, B. GUERVILLE-BALLÉ, T. SHIRANE and H.-O. TOKUNAGA, On the topology of arrangements of a cubic and its inflectional tangents. *Proc. Japan Acad. Ser. A Math. Sci.* 93 (2017), 50–53. Zbl1429.14020 MR 3659944
- [BKS] E. BRIESKORN, H. KNÖRRER and J. STILLWELL, Plane Algebraic Curves: Translated by John Stillwell. Modern Birkhäuser Classics. Springer Basel (2012). Zbl 1254.14002 MR 2975988
- [BTY] S. BANNAI, H. о ТокиNAGA and M. YAMAMOTO, A note on the topology of arrangements for a smooth plane quartic and its bitangent lines.*Hiroshima Math. J.* 49 (2019), 289–302. Zbl1423.14227 MR 3984996
- [CNEO] P. CASSOU-NOGUÈS, C. EYRAL and M. OKA, Topology of septics with the set of singularities $\mathbf{B}_{4,4} \oplus 2\mathbf{A}_3 \oplus 5\mathbf{A}_1$ and π_1 -equivalent weak Zariski pairs. *Topology Appl.* **159** (2012), 2592–2608. Zbl 1248.14035 MR 2923428
- [Deg] A. DEGTYAREV, On deformations of singular plane sextics. J. Algebraic Geom. 17 (2008), 101–135. Zbl 1131.14040 MR 2357681
- [GB] B. GUERVILLE-BALLÉ, An arithmetic Zariski 4-tuple of twelve lines. Geom. Topol. 20 (2016), 537–553. Zbl 1337.32042 MR 3470721
- [GBS] B. GUERVILLE-BALLÉ and T. SHIRANE, Non-homotopicity of the linking set of algebraic plane curves. J. Knot Theory Ramifications 26 (2017), 1750089, 13. Zbl 1375.14103 MR 3730557

[GBVS]	B. GUERVILLE-BALLÉ and J. VIU-Sos, Configurations of points and topolog
	of real line arrangements. Math. Ann. 374 (2019), 1-35. Zbl 1417.5203
	MR 3961303

- [Oka] M. OкA, Two transforms of plane curves and their fundamental groups. J. Math. Sci. Univ. Tokyo **3** (1996), 399–443. Zbl 0935.14013 MR 1424436
- [Shi] T. Shioda, Plane quartics and Mordell–Weil lattices of type E₇. Comment. Math. Univ. St. Pauli 42 (1993), 61–79. Zbl 0790.14025 MR 1223188
- [Shim1] I. SHIMADA, Fundamental groups of complements to singular plane curves. Amer. J. Math. 119 (1997), 127–157. Zbl 0956.14012 MR 1428061
- [Shim2] Equisingular families of plane curves with many connected components. *Vietnam J. Math.* **31** (2003), 193–205. Zbl1069.14036 MR 2011641
- [Shir1] T. SHIRANE, A note on splitting numbers for Galois covers and π₁equivalent Zariski k-plets. Proc. Amer. Math. Soc. 145 (2017), 1009–1017. Zbl 1358.14018 MR 3589301
- [Shir2] Connected numbers and the embedded topology of plane curves. *Can. Math. Bull.* **61** (2018), 650–658. Zbl 1429.14019 MR 3831937
- [Zar1] O. ZARISKI, On the Problem of Existence of Algebraic Functions of Two Variables Possessing a Given Branch Curve. Amer. J. Math. 51 (1929), 305–328. JFM 55.0806.01 MR 1506719
- [Zar2] On the irregularity of cyclic multiple planes. Ann. of Math. (2) **32** (1931), 485–511. JFM 57.0440.03 MR 1503012
- [Zar3] On the Poincaré Group of Rational Plane Curves. *Amer. J. Math.* **58** (1936), 607–619. JFM 62.0758.02 MR 1507185

(Reçu le 5 juin 2019)

Benoît GUERVILLE-BALLÉ, Institute of Mathematics, Polish Academy of Sciences, ul. Sniadeckich 8, 00-656 Warsaw, Poland

e-mail: benoit.guerville-balle@math.cnrs.fr

Jean-Baptiste MEILHAN, Université Grenoble Alpes, IF, 38000 Grenoble, France *e-mail:* jean-baptiste.meilhan@univ-grenoble-alpes.fr

© Fondation L'Enseignement Mathématique