Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 66 (2020)

Heft: 1-2

Artikel: A note on the G-Sarkisov program

Autor: Floris, Enrica

DOI: https://doi.org/10.5169/seals-880824

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 30.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

A note on the G-Sarkisov program

Enrica Floris

Abstract. The purpose of this note is to prove the G-equivariant Sarkisov program for a connected algebraic group G following the proof of the Sarkisov program by Hacon and McKernan. As a consequence, we obtain a characterisation of connected subgroups of Bir(Z) acting rationally on Z.

Mathematics Subject Classification (2010). Primary: 14E30, 14E07.

Keywords. Sarkisov program, Cremona group.

1. Introduction

Let Z be a normal projective and \mathbb{Q} -factorial variety. Assume Z is terminal (we refer to Definition 2.1 for the notion of terminality). There is a divisor canonically attached to Z called the canonical divisor and denoted by K_Z . One of the main ideas of birational geometry is that the behaviour of K_Z should determine the geometric properties of Z.

A cornerstone result in this direction is the Cone and Contraction theorem, due to Mori. It describes the curves which have negative intersection with K_Z and says that they can be contracted.

The K_Z -minimal model program, or K_Z -MMP, is a sequence of elementary birational maps

$$Z \longrightarrow Z_1 \longrightarrow Z_2 \longrightarrow \dots$$

which are elementary, in the sense that they are defined using the simplest contractions given by the Cone and Contraction theorem, called *extremal contractions*. The process is expected to end and there are two possible outcomes depending on the geometry of Z. Either the outcome is a variety X with nef canonical bundle, that is, such that $K_X \cdot C \ge 0$ for every curve $C \subseteq X$; or X is a Mori fibre space, that is, there is a fibration $\phi \colon X \to T$ such that $\rho(X) = \rho(T) + 1$ and $-K_X$ restricted to the fibres of ϕ is ample.

Due to the development of the Minimal Model Program, the objects considered are, more than varieties, pairs. A pair (Z,Φ) is the data of a normal variety Z and a \mathbb{Q} -Weil divisor Φ such that $K_Z+\Phi$, called the log-canonical divisor, is \mathbb{Q} -Cartier. The use of pairs originally comes from compactifications. By Hironaka's desingularization theorem, if Z^0 is a non-compact smooth variety, there is a smooth compact variety $Z \supseteq Z^0$ such that $Z \setminus Z^0$ is a simple normal crossing divisor. The datum of Z^0 is then replaced with $(Z, Z \setminus Z^0)$. This is the reason why the Weil divisor Φ is called the boundary. Pairs have many advantages. One of them is that they make the canonical divisor functorial. For instance, it is not true that for a hypersurface $Y \subseteq Z$ we have $K_Z|_Y = K_Y$, but it is true for pairs and log-canonical divisors. Indeed, there are two boundaries Φ_Z on Z and Φ_Y on Y such that $(K_Z + \Phi_Z)|_Y = K_Z + \Phi_Y$, it is sufficient to set $\Phi_Z = Y$ and $\Phi_Y = 0$.

More generally, if $f: Y \to Z$ is a morphism, it is not true that $K_Y = f^*K_Z$, but it is in many cases for log-canonical divisors. For instance, if $f: Y \to Z$ is a \mathbb{P}^n -bundle, it is possible to find boundaries Φ_Z on Z and Φ_Y on Y such that $K_Z + \Phi_Y = f^*(K_Z + \Phi_Z)$.

Moreover, pairs are a crucial tool in the proof of many important results, like Shokurov's base-point-freeness, where it is necessary to be able to modify/perturb the coefficients of Φ , even in the case $\Phi = 0$.


Another nice feature of pairs is that they encode singularities. If Z is normal but not smooth, it is possible to define the canonical divisor as a Weil divisor. Indeed, on the smooth locus Z^{sm} of Z we have $K_{Z^{sm}} = \sum a_i D_i$ and, since $codim_Z(Z \setminus Z^{sm}) \geq 2$, there is a unique Weil divisor on Z whose restriction is $K_{Z^{sm}}$, namely $K_Z = \sum a_i \overline{D}_i^{Zar}$. The divisor K_Z is in general not \mathbb{Q} -Cartier and Φ encodes the lack of "Cartier-ness" of K_Z . Moreover there is a way of measuring the singularities of (Z, Φ) by looking at the coefficients of Φ . We refer to Definition 2.1 for the notion of klt, which is a singularity notion defined in this way. The Cone and Contraction theorem holds for a klt pair and we can run the $K_Z + \Phi$ -minimal model program.

It is still an open problem whether the $K_Z + \Phi$ -MMP stops in general, even when $\Phi = 0$, but there is an important class of varieties for which it is known to terminate, the ones which are covered by curves which have negative intersection with $K_Z + \Phi$. We say in this case that $K_Z + \Phi$ is *not pseudoeffective*. Varieties with this property are uniruled. In this case, after a finite number of steps, we find a Mori fibre space by [BCHM].

The outcome is nevertheless far from being unique. The Sarkisov program describes the relation between two different Mori fibre spaces that are outcomes of two MMP on the same variety:

Theorem 1.1 ([HM], [Cor]). Suppose that $\phi: X \to S$ and $\psi: Y \to T$ are two Mori fibre spaces. Then X and Y are birational if and only if they are related by a sequence of Sarkisov links.

A Sarkisov link between $\phi: X \to S$ and $\psi: Y \to T$ is a diagram of one of the four following types:



The vertical arrows in the diagrams are extremal contractions; the arrows to X or Y are extremal divisorial contractions. The horizontal dotted arrows are isomorphisms in codimension one which are a composition of flops with respect to a suitable boundary. A flop is a composition of birational maps $h_2^{-1} \circ h_1 \colon X_1 \dashrightarrow X_2$ with $h_i \colon X_i \to \overline{X}$ such that the exceptional loci of h_1 and h_2 have codimension at least two, $\rho(X_i) = \rho(\overline{X}) + 1$ and there are divisors D_i on X_i such that (X_i, D_i) is klt and $K_{X_i} + D_i$ has zero intersection with every curve contracted by ϕ_i .

Theorem 1.1 is another case where pairs are used in a fundamental way. Indeed, if X_1/S_1 and X_2/S_2 are two Mori fibre spaces birational to each other, there is a variety Z and two boundaries Θ_1 and Θ_2 on Z such that $K_Z+\Theta_i$ is semiample on Z and induces a morphism $Z\to X_i$. Roughly speaking, the possible birational models are encoded by a combinatorial object (a polytope) in terms of boundaries on Z.

The purpose of this note is to give a proof of Theorem 1.1 in the G-equivariant setting. We prove that two Mori fibre spaces that are outcomes of two G-equivariant MMP on the same G-pair are related by a sequence of G-equivariant Sarkisov links.

Theorem 1.2. Let G be a connected algebraic group. Let (Z, Φ) be a klt pair such that $G < Aut^0(Z)$ and G leaves Φ invariant. Let $\phi: X \to S$ and $\psi: Y \to T$ be two Mori fibre spaces obtained from (Z, Φ) via a G-equivariant MMP. Then X and Y are related by a sequence of G-equivariant Sarkisov links and in every such link the horizontal dotted arrows are compositions of G-equivariant flops with respect to a suitable boundary.

In dimension 3 and for $\Phi = 0$ the interested reader can check that the proof in [Cor] is "G-equivariant".

As a direct consequence we obtain the following characterisation of subgroups of Bir(W) that are maximal among the connected groups acting rationally on W.

Corollary 1.3. Let W be an uniruled variety and let G be a connected algebraic group acting rationally on W. Then G is maximal among the connected groups acting rationally on W if and only if $G = Aut^0(X)$ where $\phi: X \to S$ is a Mori fibre space and for every Mori fibre space $\psi: Y \to T$ which is related to ϕ by a finite sequence of G-Sarkisov links we have $G = Aut^0(Y)$.

We have the following application to subgroups of the Cremona group.

Corollary 1.4. Let G be a connected algebraic group acting rationally on \mathbb{P}^n . Then G is maximal among the connected groups acting rationally on \mathbb{P}^n if and only if $G = Aut^0(X)$ where $\phi: X \to S$ is a rational Mori fibre space and for every Mori fibre space $\psi: Y \to T$ which is related to ϕ by a finite sequence of G-Sarkisov links we have $G = Aut^0(Y)$.

Theorem 1.2 is an essential ingredient in [BFT, Theorem E], the classification of automorphism groups of rational Mori fibre spaces of dimension 3. This result implies that there is a finite number of conjugacy classes of maximal connected algebraic subgroups of $Bir(\mathbb{P}^3)$ acting rationally on \mathbb{P}^3 .

2. MMP and group actions

We refer to [KM] for the definitions of singularities of pairs, of MMP and of Mori fibre space and to [DL] for the definition of rational and regular action.

We give now the definition of pairs and terminal and klt singularities:

Definition 2.1. A pair (Z, Φ) is the data of a normal variety Z and a \mathbb{Q} -divisor Φ such that $K_Z + \Phi$ is \mathbb{Q} -Cartier. A pair is said to be kawamata log terminal, or klt, if for any birational morphism $\mu \colon \widetilde{Z} \to Z$ with \widetilde{Z} smooth we have $K_{\widetilde{Z}} = \mu^*(K_Z + \Phi) + \sum a_E E$ with $a_E > -1$ for all E.

A variety Z is said to be terminal if for any birational morphism $\mu \colon \widetilde{Z} \to Z$ with \widetilde{Z} smooth we have $K_{\widetilde{Z}} = \mu^* K_Z + \sum a_E E$ with $a_E > 0$ for all $E \subseteq Exc(\mu)$.

Definition 2.2. We call a pair (Z, Φ) a G-pair if G acts on Z regularly and for all $g \in G$ we have $g \cdot \Phi = \Phi$.

Remark 2.3. The pair (Z,0) is a G-pair for every subgroup G of Aut(Z).

Remark 2.4. In the next lemma we prove that if (Z, Φ) is a G-pair with G connected, then every MMP on (Z, Φ) is a G-MMP. The situation is thus very different from the one where the group G is finite. Indeed, in the connected case, the action of G on the polyhedral part of NE(X) is trivial, while a finite group can act in a non-trivial way on NE(X) and one has to consider the G-invariant part $NE(X)^G$.

An easy example which illustrates well this difference is $(\mathbb{P}^1 \times \mathbb{P}^1, 0)$ with the action of $\mathbb{Z}/2\mathbb{Z}$ exchanging the two factors. We have $NE(\mathbb{P}^1 \times \mathbb{P}^1)^{\mathbb{Z}/2\mathbb{Z}} = \mathbb{R}_+[l_1+l_2]$ where l_1 and l_2 are fibres of the projections. The only possible $\mathbb{Z}/2\mathbb{Z}$ -invariant contraction is $\mathbb{P}^1 \times \mathbb{P}^1 \to \{p\}$. If we consider instead the diagonal action of PGL(2), we have $NE(\mathbb{P}^1 \times \mathbb{P}^1)^{PGL(2)} = \mathbb{R}_+[l_1] + \mathbb{R}_+[l_2]$ and there are two extremal contractions, the two projections onto the two factors. We refer to [KM] and $[\mathbb{C}^+]$ for a discussion of the case where G is finite, but also to the classical [Man] and to the work of Prokhorov (e.g., [Pro]).

Let G be a connected algebraic group. If we have a klt G-pair (Z, Φ) , we can run a G-equivariant MMP on (Z, Φ) , that is, an MMP where all the birational maps, divisorial contractions and flips, are compatible with the action of the group.

Lemma 2.5. Let G be a connected group and (Z, Φ) a G-pair. Then any MMP on (Z, Φ) is G-equivariant.

Proof. Indeed let $Z \longrightarrow Z_1$ be the first step of the MMP. If it is an extremal contraction, then by the Blanchard's lemma [BSU, Prop 4.2.1] there is an induced action on Z_1 making the map G-equivariant. Equivalently, a connected group acts trivially on the extremal rays contained in the $(K_Z + \Phi)$ -negative part of the Mori cone, that are discrete. Then the extremal ray corresponding to the contraction $Z \to Z_1$ is G-invariant and so is the locus spanned by it.

Assume that the first step is a flip given by the composition of two small contractions $\mu: Z \to Y$ and $\mu^+: Z_1 \to Y$. By the discussion above, there is an action on Y making μ G-equivariant. Moreover, $Z_1 \cong Proj_Y \bigoplus_m \mu_* \mathcal{O}_X(m(K_Z + \Phi))$. Since $K_Z + \Phi$ is G-invariant, the group G acts on $\mathcal{O}_X(m(K_Z + \Phi))$ for m sufficiently divisible, and subsequently on Z_1 .

If $K_Z + \Phi$ is not pseudoeffective, by [BCHM] the outcome of such an MMP is a Mori fibre space $\phi: X \to S$ with a regular action of G on X.


Definition 2.6.

• A *Mori fibre space* is the data of a fibration $\phi: X \to S$ such that X is terminal, $\rho(X) - \rho(S) = 1$ and a boundary Δ on X such that (X, Δ) is klt and $-(K_X + \Delta)$ is ϕ -ample.

- We will call two Mori fibre spaces $\phi: X \to S$ and $\psi: Y \to T$ Sarkisov related if X and Y are outcomes of running the $K_Z + \Phi$ -MMP, for the same \mathbb{Q} -factorial klt pair (Z, Φ) .
- If G is a connected group, a G-Mori fibre space is a Mori fibre space with a regular action of a group G.
- We say that two G-Mori fibre spaces $\phi: X \to S$ and $\psi: Y \to T$ are GSarkisov related if X and Y are results the G-equivariant MMP on (Z, Φ) ,
 for the same \mathbb{Q} -factorial klt G-pair (Z, Φ) .

The following remark will be useful in the next section.

Remark 2.7. If there is a commutative diagram



where f_i is birational and G-equivariant for i = 1, 2 and g_i is a morphism for i = 1, 2, then, by the Blanchard's lemma [BSU, Prop 4.2.1] there are two actions of G on Y. These actions coincide. Indeed, they coincide on an open set contained in the image of the open set where $f_2 \circ f_1^{-1}$ is an isomorphism.

Remark 2.8. If G is connected and X is uniruled, or if G is rationally connected, then all the complete linear systems are G-invariant, although the individual divisors might not be left invariant by the action of G.

3. Proof of Theorem 1.2

In this section we present a proof of Theorem 1.2. Let Z be a projective variety. From now on, we assume the setup of [HM, Section 3].

Definition 3.1. [BCHM] Let V be a finite dimensional affine subspace of the real vector space $WDiv_{\mathbb{R}}(Z)$ of Weil divisors on Z, which is defined over the rationals, and $A \geq 0$ an ample \mathbb{Q} -divisor on Z.

$$\mathcal{L}_A(V) = \{ \Theta = A + B | B \in V, B \ge 0, K_Z + \Theta \text{ is log canonical} \},$$

 $\mathcal{E}_A(V) = \{ \Theta \in \mathcal{L}_A(V) | K_Z + \Theta \text{ is pseudo-effective} \}.$

As in [HM], we assume that there exists $B_0 \in V$ such that $K_Z + \Theta_0 = K_Z + A + B_0$ is big and klt. Given a rational contraction $f: Z \longrightarrow X$, define

$$\mathcal{A}_{A,f}(V) = \{\Theta \in \mathcal{E}_A(V) | f \text{ is the ample model of } (Z,\Theta)\}.$$

In addition, let $C_{A,f}(V)$ denote the closure of $A_{A,f}(V)$. We refer to [HM, Definition 3.1] for the definition of ample model. We recall the following result from [HM] for the benefit of the reader.

Theorem 3.2. [HM, Theorem 3.3] There is a natural number m and there are $f_i: Z \longrightarrow X_i$ rational contractions $1 \le i \le m$ with the following properties:

- (1) $\{A_i = A_{A,f_i} | 1 \le i \le m\}$ is a partition of $\mathcal{E}_A(V)$. A_i is a finite union of relative interiors of rational polytopes. If f_i is birational, then $C_i = C_{A,f_i}$ is a rational polytope.
- (2) If $1 \le i \le m$ and $1 \le j \le m$ are two indices such that $A_j \cap C_i \ne \emptyset$, then there is a contraction morphism $f_{ij}: X_i \to X_j$ and a factorisation $f_j = f_{ij} \circ f_i$.

Now suppose in addition that V spans the Néron-Severi group of Z.

- (3) Pick $1 \le i \le m$ such that a connected component C of C_i intersects the interior of $\mathcal{L}_A(V)$. The following are equivalent:
 - \bullet C spans V.
 - f_i is birational and X_i is \mathbb{Q} -factorial.

The following is a G-equivariant version of [HM, Lemma 4.1].

Lemma 3.3. Let G be a connected group. Let $\phi: X \to S$ and $\psi: Y \to T$ be two G-Sarkisov related G-Mori fibre spaces corresponding to two \mathbb{Q} -factorial klt projective G-pairs (X, Δ) and (Y, Γ) . Then we may find a smooth projective variety Z with a regular action of G, two birational G-equivariant contractions $f: Z \dashrightarrow X$ and $g: Z \dashrightarrow Y$, a klt G-pair (Z, Φ) , an ample \mathbb{Q} -divisor A on Z and a two-dimensional rational affine subspace V of $WDiv_{\mathbb{R}}(Z)$ such that

(1) if $\Theta \in \mathcal{L}_A(V)$ then $\Theta - \Phi$ is ample,

- (2) $A_{A,\phi\circ f}$ and $A_{A,\psi\circ g}$ are not contained in the boundary of $\mathcal{L}_A(V)$,
- (3) V satisfies (1-4) of Theorem 3.2,
- (4) $C_{A,f}$ and $C_{A,g}$ are two dimensional, and
- (5) $C_{A,\phi\circ f}$ and $C_{A,\psi\circ g}$ are one dimensional.

Proof. By assumption we may find a \mathbb{Q} -factorial klt G-pair (Z, Φ) such that $f: Z \longrightarrow X$ and $g: Z \longrightarrow Y$ are both outcomes of the G-equivariant MMP on (Z, Φ) . Let $p: W \to Z$ be any G-equivariant log resolution of (Z, Φ) which resolves the indeterminacy of f and g. Such pair exists for instance by [Kol, Proposition 3.9.1, Theorem 3.35, Theorem 3.36]. The rest of the proof goes as in [HM, Lemma 4.1].

Proof of Theorem 1.2. The proof follows the same lines as [HM, Theorem 1.3] but instead of choosing (Z, Φ) as in [HM, Lemma 4.1] we choose the G-pair given by Lemma 3.3.

We prove now that any X_i as in Theorem 3.2 carries a regular action of G making f_i G-equivariant. Since $\Theta \in \mathcal{L}_A(V)$ implies $\Theta - \Phi$ is ample, by Theorem 3.2(3) and [HM, Lemma 3.6], for every X_i corresponding to a full-dimensional polytope of \mathcal{A}_i , the variety X_i is the result of an MMP on (Z, Φ) . By Lemma 2.5 this MMP is G-equivariant and therefore there is a regular action of G on X_i .

Let X_j be a variety corresponding to a non full-dimensional polytope \mathcal{A}_j . Let \mathcal{A}_i be a full-dimensional polytope such that $\mathcal{C}_i \cap \mathcal{A}_j \neq \emptyset$. By Theorem 3.2 there is a surjective morphism $f_{ij}: X_i \to X_j$. By the Blanchard's lemma [BSU, Prop 4.2.1] there is an action of G on X_j making the morphism f_{ij} G-equivariant. By Remark 2.7 this action does not depend on the choice of i.

The links given by [HM, Theorem 3.7] are G-equivariant. Indeed the maps appearing in the links are either

- morphisms of the form f_{ij} as in Theorem 3.2, and those are G-equivariant by the discussion above; or
- flops (with respect to a suitable boundary) of the form $f_{ij} \circ f_{kj}^{-1}$, and those are again G-equivariant.

4. Proof of Corollary 1.3

Definition 4.1. A connected subgroup G < Bir(Z) is not maximal among the connected groups acting rationally on Z if there is a connected subgroup of Bir(Z) acting rationally on Z such that $G \subsetneq H$. It is maximal among the connected groups acting rationally on Z otherwise. We will say maximal for short.

Proof of Corollary 1.3. By a theorem of Weil [Wei] (see also [Kra]) there is a birational model \widetilde{W} of W, such that G acts regularly on \widetilde{W} . We then run a G-equivariant MMP on \widetilde{W} (see Lemma 2.5) and by [BDPP] and [BCHM] the result is a G-Mori fibre space $\phi: X \to S$.

We prove now that G is maximal if and only if for every Mori fibre space $\psi: Y \to T$ which is related to ϕ by a finite sequence of G-Sarkisov links we have $G = Aut^0(Y)$.

Assume that G is maximal. Let $X/S \dashrightarrow Y/T$ be a composition of G-Sarkisov links and let $\varphi \colon X \dashrightarrow Y$ be the corresponding birational map. Then $\varphi G \varphi^{-1} \subseteq Aut^0(Y)$ and if G is maximal $\varphi G \varphi^{-1} = Aut^0(Y)$.

Assume that G is not maximal and let H be a connected subgroup of Bir(Z) acting rationally on Z and such that $G \subsetneq H$. By a theorem of Weil [Wei] there is a birational model \widehat{W} of \widetilde{W} , such that H acts regularly on \widehat{W} . We run an H-equivariant MMP on \widehat{W} and obtain a Mori fibre space $Y \to T$ such that $H < Aut^0(Y)$. This MMP is also G-equivariant. Therefore $X \to S$ and $Y \to T$ are G-Sarkisov related. By Theorem 1.2, there is a finite sequence of G-Sarkisov links $X \to S \dashrightarrow Y \to T$.

Acknowledgements. The author would like to thank the referee for his or her suggestions and comments which improved the quality of the presentation, Anne-Sophie Kaloghiros, Boris Pasquier and Luca Tasin for discussions, Andrea Fanelli and Ronan Terpereau for their comments on an earlier draft and Jérémy Blanc for his comments and for suggesting the problem and the application to subgroups of the Cremona group. The author is supported by the ANR Project FIBALGA ANR-18-CE40-0003-01.

References

- [BCHM] C. BIRKAR, P. CASCINI, C. D. HACON and J. M^cKernan, Existence of minimal models for varieties of log general type. *J. Amer. Math. Soc.* **23** (2010), 405–468. Zbl 1210.14019 MR 2601039
- [BDPP] S. BOUCKSOM, J.-P. DEMAILLY, M. PĂUN and T. PETERNELL, The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension. *J. Algebraic Geom.* **22** (2013), 201–248. Zbl 1267.32017 MR 3019449
- [BFT] J. Blanc, A. Fanelli and R. Terpereau, Connected algebraic groups acting on 3-dimensional mori fibrations. arXiv:1912.11364

- [BSU] M. BRION, P. SAMUEL and V. UMA, Lectures on the Structure of Algebraic Groups and Geometric Applications. Vol. 1 of CMI Lecture Series in Mathematics. Hindustan Book Agency, New Delhi; Chennai Mathematical Institute (CMI), Chennai (2013). Zbl 1326.14001 MR 3088271
- [C⁺] D. Chan, K. Chan, L. de Thanhoffer de Völcsey, C. Ingalls, K. Jabbusch, S. J. Kovács, R. Kulkarni, B. Lerner, B. Nanayakkara, S. Okawa and M. Van den Bergh, The minimal model program for b-log canonical divisors and applications. arXiv:1707.00834 (2017).
- [Cor] A. Corti, Factoring birational maps of threefolds after Sarkisov. J. Algebraic Geom. 4 (1995), 223–254. Zbl 0866.14007 MR 1311348
- [DL] A. Dubouloz and A. Liendo, Rationally integrable vector fields and rational additive group actions. *Internat. J. Math.* **27** (2016), 1650060, 19. Doi 10.1142/S0129167X16500609. Zbl 1362.14015 MR 3530281
- [HM] C. D. HACON and J. McKernan, The Sarkisov program. J. Algebraic Geom. 22 (2013), 389–405. Doi 10.1090/S1056-3911-2012-00599-2. Zbl 1267.14024 MR 3019454
- [KM] J. Kollár and S. Mori, Birational geometry of algebraic varieties, vol. 134 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge (1998). Doi 10.1017/CBO9780511662560. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original. Zbl 0926.14003 MR 1658959
- [Kol] J. Kollár, Lectures on Resolution of Singularities, vol. 166 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ (2007). Zbl 1113.14013 MR 2289519
- [Kra] H. Kraft, Regularization of rational group actions. arXiv:1808.08729 (2018).
- [Man] J. I. Manin, Rational surfaces over perfect fields. II. *Mat. Sb.* (*N.S.*) **72** (1967), 161–192. Zbl 0182.23701 MR 0225781
- [Pro] Y. Prokhorov, *G*-Fano threefolds, II. *Adv. Geom.* **13** (2013), 419–434. Doi 10.1515/advgeom-2013-0009. Zbl 1291.14025 MR 3100918
- [Wei] A. Weil, On algebraic groups of transformations. *Amer. J. Math.* **77** (1955), 355–391. Doi 10.2307/2372535. Zbl 0065.14201 MR 0074083

(Reçu le 13 juin 2019)

Enrica Floris, Université de Poitiers, Laboratoire de Mathématiques et Applications, UMR CNRS 7348, Téléport 2, Boulevard Marie et Pierre Curie, BP 30179, 86962 Futuroscope Chasseneuil Cedex, France

e-mail: enrica.floris@univ-poitiers.fr