

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 66 (2020)
Heft: 1-2

Artikel: Surface groups in the group of germs of analytic diffeomorphisms in one variable
Autor: Cantat, Serge / Cerveau, Dominique / Guirardel, Vincent
DOI: <https://doi.org/10.5169/seals-880825>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Surface groups in the group of germs of analytic diffeomorphisms in one variable

Serge CANTAT, Dominique CERVEAU, Vincent GUIRADEL
and Juan SOUTO

Abstract. We construct embeddings of surface groups into the group of germs of analytic diffeomorphisms in one variable.

Mathematics Subject Classification (2010). Primary: 37C85, 57M60, 20F34.

Keywords. Germs of diffeomorphisms, fundamental groups of surfaces, codimension one foliations.

Contents

1	Introduction	93
2	Germs of diffeomorphisms and the Koenigs Linearization Theorem . .	97
3	Embedding orientable surface groups	99
4	Non-orientable surface groups	105
5	The final topology on germs of diffeomorphisms	109
6	A large irreducible component of the representation variety	119
7	A p -adic proof	122
8	Complements and open questions	125
9	Appendix: Free groups	128
	References	132

1. Introduction

1.1. The main result. Let \mathbf{C} be the field of complex numbers and $\text{Diff}(\mathbf{C}, 0)$ the group of germs of analytic diffeomorphisms at the origin $0 \in \mathbf{C}$. Choosing a local coordinate z near the origin, every element $f \in \text{Diff}(\mathbf{C}, 0)$ is determined by a unique power series

$$f(z) = a_1z + a_2z^2 + a_3z^3 + \dots + a_nz^n + \dots$$

with $f'(0) = a_1 \neq 0$ and with a positive radius of convergence

$$(1.1) \quad \text{rad}(f) = \left(\limsup_{n \rightarrow +\infty} |a_n|^{1/n} \right)^{-1}.$$

We denote by $\text{Diff}(\mathbf{R}, 0) \subset \text{Diff}(\mathbf{C}, 0)$ the subgroup of real germs in this chart, i.e., with $a_i \in \mathbf{R}$ for all $i \in \mathbf{N}$ (this inclusion depends on the choice of the coordinate z). The main goal of this note is the following result, that answers a question raised by E. Ghys (see [Cer], §3.3, or also [Bru1], Problem 4.15).

Theorem A. *Let Γ be the fundamental group of a closed orientable surface, or of a closed non-orientable surface of genus ≥ 4 . Then Γ embeds in the group $\text{Diff}(\mathbf{R}, 0)$ and in particular in $\text{Diff}(\mathbf{C}, 0)$.*

We shall present three proofs of Theorem A. For simplicity, in this introduction, we restrict to the case where Γ is the fundamental group of an orientable surface of genus 2, and we consider the presentation

$$(1.2) \quad \Gamma_2 = \langle a_1, b_1, a_2, b_2 \mid [a_1, b_1] = [a_2, b_2] \rangle.$$

Our proofs of theorem A are inspired by [BGSS], where it is proved that a compact topological group or a connected Lie group which contains a dense free group of rank 2 contains a dense subgroup isomorphic to Γ_2 .

The surface groups considered in Theorem A are examples of limit groups. Recently, and independently, A. Brudnyi proved a related embedding theorem: limit groups embed into the group of (non converging) formal germs of diffeomorphisms (see [Bru2])

1.2. Compact groups. Let us describe the argument used in [BGSS] to prove the following result.

Theorem 1.1 ([BGSS]). *If a compact group G contains a free group F of rank 2, then there is an embedding $\rho: \Gamma_2 \rightarrow G$ such that $F \subset \rho(\Gamma_2)$.*

Proof. Denote by \mathbf{F}_m the free group on m generators. The first ingredient is a result by Baumslag [Bau] saying that Γ_2 is *fully residually free*; this means that there exists a sequence of morphisms $p_N: \Gamma_2 \rightarrow \mathbf{F}_2$ which is asymptotically injective: for every $g \in \Gamma_2 \setminus \{1\}$, $p_N(g) \neq 1$ if N is large enough.

To be more explicit, we use the presentation (1.2) of Γ_2 , and we note that the subgroup $\langle a_1, b_1 \rangle$ of Γ_2 is a free group $\mathbf{F}_2 = \langle a_1, b_1 \rangle$. Let $p: \Gamma_2 \rightarrow \langle a_1, b_1 \rangle$ be the morphism fixing a_1 and b_1 and sending a_2 and b_2 to a_1 and b_1

respectively. Let $\tau : \Gamma_2 \rightarrow \Gamma_2$ be the Dehn twist around the curve $c = [a_1, b_1]$, i.e. the automorphism that fixes a_1 and b_1 and sends a_2 and b_2 to ca_2c^{-1} and cb_2c^{-1} respectively.

Proposition 1.2 (see [BGSS, Corollary 2.2]). *Given any $g \in \Gamma_2 \setminus \{1\}$, there exists a positive integer n_0 such that $p \circ \tau^N(g) \neq 1$ for all $N \geq n_0$.*

Now, fix an embedding $\iota : \langle a_1, b_1 \rangle \rightarrow G$ such that $\iota(\langle a_1, b_1 \rangle) = F$. Composing $p \circ \tau^N$ with ι , we get a sequence of points $p_N := \iota \circ p \circ \tau^N$ in $\text{Hom}(\Gamma_2, G)$. Now, consider the element $h = \iota(p(c))$ of G , and let T be the closure of the cyclic group $\langle h \rangle$ in the compact group G . For $t \in T$, define a morphism $\rho_t : \Gamma_2 \rightarrow G$ by

$$(1.3) \quad \rho_t(a_1) = \iota(p(a_1)), \quad \rho_t(a_2) = t \circ \left(\iota(p(a_1)) \right) \circ t^{-1}$$

$$(1.4) \quad \rho_t(b_1) = \iota(p(b_1)), \quad \rho_t(b_2) = t \circ \left(\iota(p(b_1)) \right) \circ t^{-1}$$

these representations are well defined and satisfy $\rho_t = \iota \circ p \circ \tau^N$ when $t = h^N$. Moreover, on the subgroup $\langle a_1, b_1 \rangle$, ρ_t coincides with $\iota \circ p$, so $F \subset \rho_t(\Gamma_2)$. Thus, $(\rho_t)_{t \in T}$ is a compact subset $\mathcal{R}(T) \subset \text{Hom}(\Gamma_2, G)$ that contains the sequence of points p_N . For every g in $\Gamma_2 \setminus \{1\}$, the subset $\mathcal{R}(T)_g = \{\rho_t \mid \rho_t(g) \neq 1\}$ is open, and Proposition 1.2 shows that it is dense because $\{h^n \mid n \geq n_0\}$ is dense in T for every integer n_0 . By the Baire theorem, the subset of injective representations ρ_t is a dense G_δ in $\mathcal{R}(T)$, and this proves Theorem 1.1. \square

The group $\text{Diff}(\mathbf{R}, 0)$ contains non-abelian free groups (this is well known, see Section 3.3), and one may want to copy the above argument for $G = \text{Diff}(\mathbf{R}, 0)$ instead of a compact group. The Koenigs linearization theorem says that if $f \in \text{Diff}(\mathbf{R}, 0)$ satisfies $f'(0) > 1$, then f is conjugate to the homothety $z \mapsto f'(0)z$; in particular, there is a flow of diffeomorphisms $(\varphi^t)_{t \in \mathbf{R}}$ for which $\varphi^1 = f$. In our argument, the compact group T introduced to prove Theorem 1.1 will be replaced by such a flow, hence by a group isomorphic to $(\mathbf{R}, +)$. Also, in that proof, $h = \iota(p(c))$ was a commutator, and the derivative of any commutator in $\text{Diff}(\mathbf{R}, 0)$ is equal to 1 at the origin, so that Koenigs theorem can not be applied to a commutator. Thus, we need to change p_N into a different sequence of morphisms: the Dehn twist τ will be replaced by another automorphism of Γ_2 , twisting along three non-separating curves.

This argument will be described in details in Sections 2 and 3; the reader who wants the simplest proof of Theorem A in the case of orientable surfaces only needs to read these sections. Non orientable surfaces are dealt with in Section 4.

1.3. Lie groups. Now, let us look at representations in a linear algebraic subgroup G of $\mathrm{GL}_m(\mathbf{R})$. Assuming that there is a faithful representation $\iota: \mathbf{F}_2 \rightarrow G$ with dense image, we shall construct a faithful representation $\Gamma_2 \rightarrow G$.

The representation variety $\mathrm{Hom}(\Gamma_2, G)$ is an algebraic subset of G^4 . Let \mathcal{R} be the irreducible component containing the trivial representation. Let $p_N: \Gamma_2 \rightarrow \mathbf{F}_2$ be an asymptotically injective sequence of morphisms, as given by Baumslag's proposition. When the image of ρ is dense, one can prove that $\iota \circ p_N$ is in \mathcal{R} for arbitrarily large values of N . For $g \in \Gamma_2 \setminus \{1\}$, the subset $\mathcal{R}_g \subset \mathcal{R}$ of homomorphisms killing g is algebraic, and it is a proper subset because it does not contain $\iota \circ p_N$ for some large N . Then, a Baire category argument in \mathcal{R} implies that a generic choice of $\rho \in \mathcal{R}$ is faithful.

To apply this argument to $G = \mathrm{Diff}(\mathbf{R}, 0)$, one needs a good topology on $\mathrm{Diff}(\mathbf{R}, 0)$, and a good "irreducible variety" $\mathcal{R} \subset \mathrm{Hom}(\Gamma_2, G)$ containing $\iota \circ p_N$, in which a Baire category argument can be used. This approach may seem difficult because $\mathrm{Hom}(\Gamma_2, G)$ is a priori far from being an irreducible analytic variety but, again, the Koenigs linearization theorem will provide the key ingredient.

First, we shall adapt an idea introduced by Leslie in [Les] to define a useful group topology on $\mathrm{Diff}(\mathbf{R}, 0)$ (see Section 5). With this topology, $\mathrm{Diff}(\mathbf{R}, 0)$ is an increasing union of Baire spaces, which will be enough for our purpose. Denote by $\mathrm{Cont}(\mathbf{R}, 0) \subset \mathrm{Diff}(\mathbf{R}, 0)$ the set of elements $f \in \mathrm{Diff}(\mathbf{R}, 0)$ with $|f'(0)| < 1$; Cont stands for "contractions". Consider the set \mathcal{R} of representations $\rho: \Gamma_2 \rightarrow \mathrm{Diff}(\mathbf{R}, 0)$ with $\rho(a_1)$ tangent to the identity, and $\rho(b_1) \in \mathrm{Cont}(\mathbf{R}, 0)$. Then, the key fact is that the map

$$\begin{aligned} \Psi: \mathcal{R} &\rightarrow \mathrm{Cont}(\mathbf{R}, 0) \times \mathrm{Diff}(\mathbf{R}, 0) \times \mathrm{Diff}(\mathbf{R}, 0) \\ \rho &\mapsto (\rho(b_1), \rho(a_2), \rho(b_2)) \end{aligned}$$

is a continuous bijection. Indeed, the defining relation of Γ is equivalent to $a_1 b_1 a_1^{-1} = [a_2, b_2] b_1$. Given $(g_1, f_2, g_2) \in \mathrm{Cont}(\mathbf{R}, 0) \times \mathrm{Diff}(\mathbf{R}, 0) \times \mathrm{Diff}(\mathbf{R}, 0)$, the germs g_1 and $[f_2, g_2]g_1$ have the same derivative at the origin and, from the Koenigs linearization theorem, there is a unique $f_1 \in \mathrm{Diff}(\mathbf{R}, 0)$ tangent to the identity solving the equation $f_1 g_1 f_1^{-1} = [f_2, g_2]g_1$: by construction there is a unique morphism $\rho: \Gamma_2 \rightarrow \mathrm{Diff}(\mathbf{R}, 0)$ that maps the a_i to the f_i , and the b_i to the g_i , and this representation satisfies $\Psi(\rho) = (g_1, f_2, g_2)$. With this bijection Ψ and the topology of Leslie, we can identify \mathcal{R} with a union of Baire spaces, in which the Baire category argument applies.

1.4. Other fields. Let \mathbf{k} be a finite field with p elements. The group $\mathrm{Diff}^1(\mathbf{k}, 0)$, also known as the Nottingham group, is the group of power series tangent to the identity and with coefficients in the finite field \mathbf{k} . It is a compact group containing a free group (see [Sze]). Thus, by [BGSS], it contains a surface group.

Now, let p be a prime number, and let \mathbf{Q}_p be the field of p -adic numbers.

Consider the subgroup $\text{Diff}^1(\mathbf{Z}_p, 0) \subset \text{Diff}(\mathbf{Q}_p, 0)$ of formal power series tangent to the identity and with coefficients in \mathbf{Z}_p . First, note that all elements f of $\text{Diff}^1(\mathbf{Z}_p, 0)$ satisfy $\text{rad}(f) \geq 1$, so that $\text{Diff}^1(\mathbf{Z}_p, 0)$ acts faithfully as a group of (p -adic analytic) homeomorphisms on $\{z \in \mathbf{Z}_p ; |z| < 1\}$. So, in that respect, $\text{Diff}^1(\mathbf{Z}_p, 0)$ is much better than the group of germs of diffeomorphisms $\text{Diff}(\mathbf{C}, 0)$. Moreover, with the topology given by the product topology on the coefficients $a_n \in \mathbf{Z}_p$ of the power series, the group $\text{Diff}^1(\mathbf{Z}_p, 0)$ becomes a compact group. And this compact group contains a free group. By the result of [BGSS] described in Section 1.2, it contains a copy of the surface group Γ_2 . So, we get a surface group acting faithfully as a group of p -adic analytic homeomorphisms on $\{z \in \mathbf{Z}_p ; |z| < 1\}$. In Section 7 we give a third proof of Theorem A that starts with the case of p -adic coefficients.

1.5. Organisation. The article is split in four parts.

- I. Sections 2 to 4 give a first proof of Theorem A; Section 4 is the only place where we deal with non-orientable surfaces. We refer to Theorem B in Section 4.3 for a stronger result, in which the field \mathbf{R} is replaced by any non-discrete, complete valued field \mathbf{k} .
- II. Section 5 and 6 present our second proof, based on the construction of a group topology on $\text{Diff}(\mathbf{C}, 0)$.
- III. Then, our p -adic proof is presented in Section 7.
- IV. Section 8 draws some consequences and list a few open problems, while the appendix shows how to construct free groups in $\text{Diff}(\mathbf{C}, 0)$, or $\text{Diff}(\mathbf{k}, 0)$ for any non-discrete and complete valued field.

Part I

2. Germs of diffeomorphisms and the Koenigs Linearization Theorem

2.1. Formal diffeomorphisms. Let \mathbf{k} be a field (of arbitrary characteristic). Denote by $\mathbf{k}[[z]]$ the ring of formal power series in one variable with coefficients in \mathbf{k} . For every integer $n \geq 0$, let $A_n: \mathbf{k}[[z]] \rightarrow \mathbf{k}$ denote the n -th coefficient function:

$$(2.1) \quad A_n: f = \sum a_n z^n \mapsto A_n(f) = a_n.$$

A *formal diffeomorphism* is a formal power series $f \in \mathbf{k}[[z]]$ such that $A_0(f) = 0$ and $A_1(f) \neq 0$. The composition $f \circ g$ determines a group law on the set

$$(2.2) \quad \widehat{\text{Diff}}(\mathbf{k}, 0) = \{f \in \mathbf{k}[[z]] \mid A_0(f) = 0 \text{ and } A_1(f) \neq 0\}$$

of all formal diffeomorphisms.

For each $n \geq 1$, there is a polynomial $P_n \in \mathbf{Z}[A_1, B_1, \dots, A_n, B_n]$ such that if $f = \sum a_n z^n$ and $g = \sum b_n z^n$ then $f \circ g = \sum_{n \geq 1} P_n(a_1, b_1, \dots, a_n, b_n) z^n$. Similarly, there are polynomials $Q_n \in \mathbf{Z}[A_1, \dots, A_n][A_1^{-1}]$ such that $f^{-1} = \sum_{n \geq 1} Q_n(a_1, \dots, a_n) z^n$ if $f = \sum a_n z^n$; the polynomial function Q_n is given by the following *inversion formula*:

$$\frac{1}{a_1^n} \sum_{k_1, k_2, \dots} (-1)^{k_1 + k_2 + \dots} \cdot \frac{(n+1) \cdots (n-1 + k_1 + k_2 + \dots)}{k_1! k_2! \cdots} \cdot \left(\frac{a_2}{a_1}\right)^{k_1} \left(\frac{a_3}{a_1}\right)^{k_2} \cdots$$

where $a_i = A_i(f)$ and the sum is over all sequences of integers k_i such that

$$k_1 + 2k_2 + 3k_3 + \cdots = n - 1.$$

We refer to [Jen] where this is proved for f and g tangent to the identity; the general case easily follows.

To encapsulate this kind of properties, we introduce the following definition. Let m be a positive integer. By definition, a function $Q: \widehat{\text{Diff}}(\mathbf{k}, 0)^m \rightarrow \mathbf{k}$ is a *polynomial function* with integer coefficients, if there is an integer n , and a polynomial $q \in \mathbf{Z}[A_{1,1}, A_{2,1}, \dots, A_{m-1,n}, A_{m,n}][A_{1,1}^{-1}, \dots, A_{m,1}^{-1}]$ such that

$$(2.3) \quad Q(f_1, \dots, f_m) = q(A_1(f_1), \dots, A_n(f_m))$$

for all m -tuples $(f_1, \dots, f_m) \in \widehat{\text{Diff}}(\mathbf{k}, 0)^m$; we denote by $\mathbf{Z}[\widehat{\text{Diff}}(\mathbf{k}, 0)^m]$ this ring of polynomial functions.

Let $\mathbf{F}_m = \langle e_1, \dots, e_m \rangle$ be the free group of rank m . To every word $w = e_{i_1}^{n_1} \cdots e_{i_k}^{n_k}$ in \mathbf{F}_m , with exponents $n_i \in \mathbf{Z}$, we associate the *word map* $w: \widehat{\text{Diff}}(\mathbf{k}, 0)^m \rightarrow \widehat{\text{Diff}}(\mathbf{k}, 0)$,

$$(2.4) \quad (g_1, \dots, g_m) \mapsto w(g_1, \dots, g_m) \stackrel{\text{def}}{=} g_{i_1}^{n_1} \circ \cdots \circ g_{i_k}^{n_k}.$$

Since composition and inversion are polynomial functions on $\widehat{\text{Diff}}(\mathbf{k}, 0)$, we get:

Lemma 2.1. *Let $w: \widehat{\text{Diff}}(\mathbf{k}, 0)^m \rightarrow \widehat{\text{Diff}}(\mathbf{k}, 0)$ be the word map given by some element of the free group \mathbf{F}_m . For each $n \geq 1$, there is a polynomial function $Q_{w,n} \in \mathbf{Z}[\widehat{\text{Diff}}(\mathbf{k}, 0)^m]$ such that*

$$A_n(w(g_1, \dots, g_m)) = Q_{w,n}(g_1, \dots, g_m)$$

for all $g_1, \dots, g_m \in \widehat{\text{Diff}}(\mathbf{k}, 0)$.

2.2. Diffeomorphisms and Koenigs linearization Theorem. Suppose now that \mathbf{k} is endowed with an absolute value $|\cdot|: \mathbf{k} \rightarrow \mathbf{R}_+$. Then \mathbf{k} becomes a metric space with the distance induced by $|\cdot|$. We shall almost always assume that

- \mathbf{k} is not discrete, equivalently there is an element $x \in \mathbf{k}$ with $|x| \neq 0, 1$;
- \mathbf{k} is complete.

Let $\mathbf{k}\{z\}$ be the subring of $\mathbf{k}[[z]]$ consisting of power series $f(z) = \sum a_n z^n$ whose radius of convergence $\text{rad}(f)$ is positive (see Equation (1.1)). When \mathbf{k} is complete, the series $\sum a_n z^n$ converges uniformly in the closed disk $\mathbb{D}_r = \{z \in \mathbf{k} \mid |z| \leq r\}$ for every $r < \text{rad}(f)$. The group of germs of analytic diffeomorphisms is the intersection $\text{Diff}(\mathbf{k}, 0) := \widehat{\text{Diff}}(\mathbf{k}, 0) \cap \mathbf{k}\{z\}$; it is a subgroup of $\widehat{\text{Diff}}(\mathbf{k}, 0)$.

A germ $f \in \text{Diff}(\mathbf{k}, 0)$ is *hyperbolic* if $|A_1(f)| \neq 1$. The following result is proved in [Mil, Chapter 8] and [HY, Theorem 1, p. 423] (see also [Sie, Theorem 1] or [Koe]).

Theorem 2.2 (Koenigs linearization theorem). *Let $(\mathbf{k}, |\cdot|)$ be a complete, non-discrete valued field. Let $f \in \text{Diff}(\mathbf{k}, 0)$ be a hyperbolic germ of diffeomorphism. There is a unique germ of diffeomorphism $h \in \text{Diff}(\mathbf{k}, 0)$ such that $h(f(z)) = A_1(f) \cdot h(z)$ and $A_1(h) = 1$.*

3. Embedding orientable surface groups

3.1. Abstract setting. Our strategy to construct embeddings of surface groups relies on the following simple remark. Let Γ be a countable group, and G be any group. Consider a non-empty topological space \mathcal{R} , with a map $\Phi: s \in \mathcal{R} \mapsto \Phi_s \in \text{Hom}(\Gamma, G)$. Given $g \in \Gamma$, set $\mathcal{R}_g = \{s \in \mathcal{R} \mid \Phi_s(g) = 1\}$.

Lemma 3.1. *Assume that \mathcal{R} has the following three properties:*

- (1) **Baire:** \mathcal{R} is a Baire space;
- (2) **Separation:** for every $g \neq 1$ in Γ , $\Phi_s(g) \neq 1$ for some $s \in \mathcal{R}$;
- (3) **Irreducibility:** for every $g \in \Gamma$, either $\mathcal{R}_g = \mathcal{R}$ or \mathcal{R}_g is closed with empty interior.

Then the set of $s \in \mathcal{R}$ such that Φ_s is an injective homomorphism is a dense G_δ in \mathcal{R} ; in particular, it is non-empty.

Proof. For any $g \in \Gamma \setminus \{1\}$, one has $\mathcal{R}_g \neq \mathcal{R}$ by (2), so \mathcal{R}_g is closed with empty interior by (3). By the Baire property, $\mathcal{R} \setminus (\cup_{g \in \Gamma \setminus \{1\}} \mathcal{R}_g)$ is a dense G_δ . But $\mathcal{R} \setminus (\cup_{g \in \Gamma \setminus \{1\}} \mathcal{R}_g)$ is precisely the set of $s \in \mathcal{R}$ such that Φ_s is injective. \square

3.2. Baumslag Lemma. As explained in the introduction, it is proved in [Bau] that the fundamental group of an orientable surface is fully-residually free. We need a precise version of this result; to obtain it, the main technical input is the Baumslag's Lemma (see [Ols, Lemma 2.4]):

Lemma 3.2 (Baumslag's Lemma). *Let $n \geq 1$ be a positive integer. Let g_0, \dots, g_n be elements of \mathbf{F}_k , and let c_1, \dots, c_n be elements of $\mathbf{F}_k \setminus \{1\}$. Assume that for all $1 \leq i \leq n-1$, $g_i^{-1}c_i g_i$ does not commute with c_{i+1} . Then for N large enough,*

$$g_0 c_1^N g_1 c_2^N \dots c_{n-1}^N g_{n-1} c_n^N g_n \neq 1.$$

Sketch of proof (I). The group $\mathrm{PSL}_2(\mathbf{R})$ acts on the hyperbolic plane \mathbb{H} by isometries, and contains a subgroup Γ such that (0) Γ is isomorphic to \mathbf{F}_k , (1) every element $g \neq \mathrm{Id}$ in Γ is a loxodromic isometry of \mathbb{H} , and (2) two elements g and h in $\Gamma \setminus \{\mathrm{Id}\}$ commute if and only if they have the same axis, which happens if and only if they share a common fixed point on $\partial\mathbb{H}$. One can find such a group in any lattice of $\mathrm{PSL}_2(\mathbf{R})$. To prove the lemma, we prove it in Γ .


Fix a base point $x \in \mathbb{H}$, denote by α_i and ω_i the repulsive and attracting fixed points of c_i in $\partial\mathbb{H}$, and consider the word

$$g_0 c_1^N g_1 c_2^N g_2.$$

For m large enough, $c_2^m g_2$ maps x to a point which is near ω_2 . If $g_1(\omega_2)$ were equal to α_1 , then c_1 and $g_1 c_2 g_1^{-1}$ would share the common fixed point α_1 , and they would commute. Thus, $g_1(\omega_2) \neq \alpha_1$ and then $g_0 c_1^{m'} g_1 c_2^m g_2$ maps x to a point which is near $g_0(\omega_1)$ if m' is large enough. Thus, $g_0 c_1^N g_1 c_2^N g_2(x) \neq x$ for large N . The proof is similar if n is larger than 2. \square

Sketch of proof (II). We rephrase this proof, using the action of \mathbf{F}_k on its boundary, because this boundary will also be used in the proof of Proposition 3.3.

Fix a basis a_1, \dots, a_k of \mathbf{F}_k , and denote by $\partial\mathbf{F}_k$ the boundary of \mathbf{F}_k . The elements of $\partial\mathbf{F}_k$ are represented by infinite reduced words in the generators a_i and their inverses. If g is an element of \mathbf{F}_k and α is an element of $\partial\mathbf{F}_k$ the concatenation $g \cdot \alpha$ is an element of $\partial\mathbf{F}_k$: this defines an action of \mathbf{F}_k by homeomorphisms on the Cantor set $\partial\mathbf{F}_k$. If g is a non-trivial, its action on $\partial\mathbf{F}_k$ has exactly two fixed points, given by the infinite words $\omega(g) = g \dots g \dots$ and $\alpha(g) = g^{-1} \dots g^{-1} \dots$ (there are no simplifications if g is given by a reduced and cyclically reduced word). Then we get: (1) every element $g \neq \mathrm{Id}$ in \mathbf{F}_k has a north-south dynamics on $\partial\mathbf{F}_k$, every orbit $g^n \cdot \beta$ converging to $\omega(g)$, except when $\beta = \alpha(g)$, and (2) two elements g and h in $\mathbf{F}_k \setminus \{\mathrm{Id}\}$ commute if and only if they have the same fixed points, which happens if and only if they share



$$\begin{aligned}
 a_0 &= \bar{a}_0 = \alpha_0 \\
 a_1 &= \eta_1 \alpha_1 \eta_1^{-1} \\
 \bar{a}_1 &= \bar{\eta}_1 \alpha_1 \bar{\eta}_1^{-1} \\
 a_2 &= \eta_2 \alpha_2 \eta_2^{-1} \\
 \bar{a}_2 &= \bar{\eta}_2 \alpha_2 \bar{\eta}_2^{-1} \\
 t_1 &= \eta_1 \bar{\eta}_1^{-1} \\
 t_2 &= \eta_2 \bar{\eta}_2^{-1}
 \end{aligned}$$

FIGURE 1

The fundamental group Γ_2 . – The α_i are three loops, while the η_j and $\bar{\eta}_j$ are four paths. The figure is symmetric with respect to the plane cutting the surface along the loops α_i .

a common fixed point on $\partial\mathbf{F}_k$. One can then repeat the previous proof with the action of \mathbf{F}_k on its boundary. \square

Write the surface of genus 2 as the union of two pairs of pants as in Figure 1, with respective fundamental groups

$$(3.1) \quad \langle a_0, a_1, a_2 \mid a_0 a_1 a_2 = 1 \rangle \text{ and } \langle \bar{a}_0, \bar{a}_1, \bar{a}_2 \mid \bar{a}_0 \bar{a}_1 \bar{a}_2 = 1 \rangle.$$

This gives the presentation

$$(3.2) \quad \Gamma_2 = \left\langle \begin{array}{l} a_0, a_1, a_2, \\ \bar{a}_0, \bar{a}_1, \bar{a}_2, \\ t_1, t_2 \end{array} \left| \begin{array}{l} a_0 a_1 a_2 = 1, \\ \bar{a}_0 \bar{a}_1 \bar{a}_2 = 1, \\ \bar{a}_0 = a_0, \bar{a}_1 = t_1^{-1} a_1 t_1, \bar{a}_2 = t_2^{-1} a_2 t_2 \end{array} \right. \right\rangle$$

which can be rewritten as

$$(3.3) \quad \Gamma_2 = \langle a_0, a_1, a_2, t_1, t_2 \mid a_0 a_1 a_2 = 1, a_0 t_1^{-1} a_1 t_1 t_2^{-1} a_2 t_2 = 1 \rangle.$$

Denote by $p : \Gamma_2 \rightarrow \langle a_0, a_1, a_2 \rangle \simeq \mathbf{F}_2$ the morphism defined by $p(a_i) = a_i$, $p(\bar{a}_i) = a_i$, and $p(t_1) = p(t_2) = 1$. Let $\tau : \Gamma_2 \rightarrow \Gamma_2$ be the (left) Dehn twist along the three curves a_0 , a_1 , and a_2 , i.e. the automorphism fixing a_i and sending t_i to $a_i t_i a_i^{-1}$ for $i = 1, 2$. Note the following facts:

- τ sends \bar{a}_i to $a_0 \bar{a}_i a_0^{-1}$; in particular, if g is a word in the \bar{a}_i , then $\tau^N(g) = a_0^N g a_0^{-N}$;
- $p \circ \tau^N$ fixes a_i for every $i = 0, 1, 2$, and

$$(3.4) \quad p \circ \tau^N(t_j) = a_j^N a_0^{-N}$$

for $j = 1, 2$.

Proposition 3.3. *For every $g \in \Gamma_2 \setminus \{1\}$, there exists a positive integer n_0 such that $p \circ \tau^N(g) \neq 1$ for all $N \geq n_0$.*

Proof. To uniformize notations, we define $t_0 = 1$ so that for all $i \in \{0, 1, 2\}$ the relation $t_i \bar{a}_i t_i^{-1} = a_i$ holds, and τ maps t_i to $a_i t_i a_0^{-1}$. Let $A = \langle a_0, a_1, a_2 \rangle$ and $\bar{A} = \langle \bar{a}_0, \bar{a}_1, \bar{a}_2 \rangle$. Write g as a shortest possible word of the following form:

$$(3.5) \quad g = g_0 t_{i_1} g_1 t_{i_2}^{-1} g_2 t_{i_3} \cdots g_{n-1} t_{i_n}^{-1} g_n$$

where n is even, $i_k \in \{0, 1, 2\}$ for all $k \leq n$, $g_k \in A$ for k even, $g_k \in \bar{A}$ for k odd, and the exponent of t_{i_k} is $(-1)^{k+1}$ (we allow $g_k = 1$). One easily checks that g can be written in this form because all generators can (for instance $t_1 = 1 \cdot t_1 \cdot 1 \cdot t_0^{-1} \cdot 1$).

If k is such that $i_k = i_{k+1}$, then $g_k \notin \langle a_{i_k} \rangle$ if k is even (resp $g_k \notin \langle \bar{a}_{i_k} \rangle$ if k is odd) as otherwise, one could shorten the word using the relation $t_{i_k} a_{i_k} t_{i_k}^{-1} = \bar{a}_{i_k}$.

First claim. *If $k \in \{2, \dots, n-2\}$ is even, $g_k^{-1} a_{i_k} g_k$ does not commute to $a_{i_{k+1}}$.*

If $i_k \neq i_{k+1}$, this is because $g_k \in A \simeq \mathbf{F}_2$ and no pair of A -conjugates of a_{i_k} and $a_{i_{k+1}}$ commute. If $i_k = i_{k+1}$, then $g_k \notin \langle a_{i_k} \rangle$ as we have just seen; since a_{i_k} is not a proper power in A , this shows that $g_k \cdot (a_{i_k}^{+\infty}) \neq a_{i_k}^{+\infty}$ in the boundary at infinity of the free group A , so $g_k^{-1} a_{i_k} g_k$ does not commute with a_{i_k} , and the claim follows.

Similarly, using the fact that $g_k \in \bar{A}$ for odd indices, we obtain:

Second claim. *If $k \leq n-1$ is odd, $g_k^{-1} \bar{a}_{i_k} g_k$ does not commute to $\bar{a}_{i_{k+1}}$.*

We have $\tau^N(g_k) = g_k$ if k is even, and $\tau^N(g_k) = a_0^N g_k a_0^{-N}$ if k is odd. After simplifications, one has

$$(3.6) \quad \tau^N(g) = g_0 a_{i_1}^N t_{i_1} g_1 t_{i_2}^{-1} a_{i_2}^{-N} g_2 a_{i_3}^N t_{i_3} \cdots g_{n-1} t_{i_n}^{-1} a_{i_n}^{-N} g_n.$$

For k odd, denote by $g'_k \in \mathbf{F}_r$ the image of g_k under p . Applying p , we thus get

$$(3.7) \quad p \circ \tau^N(g) = g_0 a_{i_1}^N g'_1 a_{i_2}^{-N} g_2 a_{i_3}^N g'_3 \cdots g'_{n-1} a_{i_n}^{-N} g_n,$$

with $g'_i := p(g_i)$. Let us check that the hypotheses of the Baumslag Lemma 3.2 apply. For k even, the first claim shows that $g_k^{-1} a_{i_k} g_k$ does not commute to $a_{i_{k+1}}$, as required. For k odd, we use that p is injective on \bar{A} and that \bar{A} contains $g_k^{-1} \bar{a}_{i_k} g_k$ and $\bar{a}_{i_{k+1}}$, and we apply the second claim to deduce that $g_k'^{-1} a_{i_k} g'_k$ does not commute to $a_{i_{k+1}}$. Applying Baumslag's Lemma, we conclude that $p \circ \tau^N(g) \neq 1$ for N large enough. \square

3.3. Embeddings of free groups. The group $\text{Diff}(\mathbf{R}, 0)$ contains non-abelian free groups. This has been proved by arithmetic means [Whi, Gla], by looking at the monodromy of generic polynomial planar vector fields [IP], and by a dynamical argument [MRR]. We shall need the following precise version of that result.

Theorem 3.4. *Let $(\mathbf{k}, |\cdot|)$ be a complete non-discrete valued field. For every pair (λ_1, λ_2) in \mathbf{k}^* , there exists a pair $f_1, f_2 \in \text{Diff}(\mathbf{k}, 0)$ that generates a free group and satisfies $f_1'(0) = \lambda_1$ and $f_2'(0) = \lambda_2$.*

This result is proved in [BCLN, Proposition 4.3] for generic pairs of derivatives (λ_1, λ_2) . We provide a proof of Theorem 3.4 in the Appendix, extending the argument of [MRR]. We refer to Section 7.1 below for other approaches.

3.4. Embedding orientable surface groups. We can now prove Theorem A for orientable surfaces:

Theorem 3.5. *Let Γ_g be the fundamental group of a closed, orientable surface of genus g . Then, there exists an injective morphism $\Gamma_g \rightarrow \text{Diff}(\mathbf{R}, 0)$.*

The group Γ_0 is trivial. The group Γ_1 is isomorphic to \mathbf{Z}^2 , hence it embeds in the group of homotheties $z \mapsto \lambda z$, $\lambda \in \mathbf{R}_+^*$. If $g \geq 2$, then Γ_g embeds in Γ_2 . To see this, fix a surjective morphism $\Gamma_2 \rightarrow \mathbf{Z}$, and take the preimage $\Lambda \subset \Gamma_2$ of the subgroup $(g-1)\mathbf{Z} \subset \mathbf{Z}$. Then, Λ is a normal subgroup of index $g-1$ in Γ_2 , and it is the fundamental group of a closed surface Σ , given by a Galois cover of degree $g-1$ of the surface of genus 2. Since the Euler characteristic is multiplicative, the genus of Σ satisfies $-2(g-1) = 2 - 2g(\Sigma)$. Thus, $g(\Sigma) = g$ and Λ is isomorphic to Γ_g . Thus, we now restrict to the case $g = 2$.

By Theorem 3.4, we can fix an injective morphism

$$(3.8) \quad \rho_0 : \mathbf{F}_2 = \langle a_0, a_1, a_2 \mid a_0 a_1 a_2 = 1 \rangle \rightarrow \text{Diff}(\mathbf{R}, 0)$$

such that the images $f_1 = \rho_0(a_1)$, $f_2 = \rho_0(a_2)$, and $f_0 = \rho_0(a_0) = f_2^{-1} f_1^{-1}$ satisfy

$$(3.9) \quad f_1'(0) = \lambda_1 > 1, \quad f_2'(0) = \lambda_2 > 1, \quad f_0'(0) = \lambda_0 < 1$$

for some real numbers λ_1 and $\lambda_2 > 1$ and $\lambda_0 = (\lambda_1 \lambda_2)^{-1}$. In particular, f_0 , f_1 , and f_2 are hyperbolic. For $\lambda \in \mathbf{R}^*$, denote by $m_\lambda(z) = \lambda z$ the corresponding homothety. For $i \in \{0, 1, 2\}$, the Koenigs linearization theorem shows that f_i is conjugate to the homothety m_{λ_i} : there is a germ of diffeomorphism $h_i \in \text{Diff}(\mathbf{R}, 0)$ such that $f_i = h_i \circ m_{\lambda_i} \circ h_i^{-1}$. Thus f_i extends to the multiplicative flow $\varphi_i : \mathbf{R}_+^* \rightarrow \text{Diff}(\mathbf{R}, 0)$ defined by $\varphi_i^s = h_i \circ m_s \circ h_i^{-1}$ for $s \in \mathbf{R}_+^*$; by construction,

$\varphi_i^{\lambda_i} = f_i$ and φ_i^s commutes with f_i for all $s > 0$. We note that $s \mapsto \varphi_i^s$ is polynomial in the sense that for all $k \in \mathbf{N}$, $s \mapsto A_k(\varphi_i^s)$ is a polynomial function with real coefficients in the variables s and s^{-1} .

Set $\mathcal{R} = (\mathbf{R}_+^*)^3$. As in Section 3.2, consider the presentation

$$(3.10) \quad \Gamma_2 = \langle a_0, a_1, a_2, t_1, t_2 \mid a_0 a_1 a_2 = 1, a_0 t_1^{-1} a_1 t_1 t_2^{-1} a_2 t_2 = 1 \rangle.$$

Given $s = (s_0, s_1, s_2) \in (\mathbf{R}_+^*)^3$, we define a morphism $\Phi_s : \Gamma_2 \rightarrow \text{Diff}(\mathbf{R}, 0)$ by

$$(3.11) \quad \Phi_s(a_i) = f_i \quad \text{for } i \in \{0, 1, 2\}$$

$$(3.12) \quad \Phi_s(t_i) = \varphi_i^{s_i} \varphi_0^{s_0} \quad \text{for } i \in \{1, 2\}$$

This provides a well defined homomorphism because φ_i commutes with f_i . As we shall see below, this morphism Φ_s is constructed to coincide with $\rho_0 \circ p \circ \tau^N$ for $s = (\lambda_0^N, \lambda_1^N, \lambda_2^N)$ (see Equation (3.4)).

Remark 3.6. For every $s \in \mathcal{R}$, the image of Φ_s contains f_1 and f_2 , hence the free group $\rho_0(\mathbf{F}_2)$. This will be used in Section 4.3.

To conclude, we check that the three assumptions of Lemma 3.1 hold for this family of morphisms $(\Phi_s)_{s \in \mathcal{R}}$.

Clearly, \mathcal{R} is a Baire space.

To check the irreducibility property, consider $g \in \Gamma_2$ and assume that $\mathcal{R}_g \neq \mathcal{R}$: this means that there exists a parameter $s \in \mathcal{R}$ and an index $k \geq 1$ such that $A_k(\Phi_s(g)) \neq A_k(\text{Id})$. The map $s = (s_0, s_1, s_2) \mapsto A_k(\Phi_s(g)) - A_k(\text{Id})$ is a polynomial function in the variables $s_0^{\pm 1}$, $s_1^{\pm 1}$, and $s_2^{\pm 1}$ that does not vanish identically on \mathcal{R} , so its zero set is a closed subset with empty interior.

We now check that \mathcal{R} has the separation property. As in Section 3.2, denote by $p : \Gamma_2 \rightarrow \mathbf{F}_2 = \langle a_0, a_1, a_2 \mid a_0 a_1 a_2 = 1 \rangle$ the morphism obtained by killing t_1 and t_2 . For the parameter $s = (1, 1, 1)$, Φ_s is equal to $\rho_0 \circ p$. More generally, setting $s_N = (\lambda_0^N, \lambda_1^N, \lambda_2^N)$ for $N \in \mathbf{N}$, the morphism $\Phi_{s_N} : \Gamma_2 \rightarrow \text{Diff}(\mathbf{R}, 0)$ satisfies

$$(3.13) \quad \Phi_{s_N}(a_i) = f_i \quad \text{for } i \in \{0, 1, 2\}$$

$$(3.14) \quad \Phi_{s_N}(t_i) = \varphi_i^{u_i^N} \varphi_0^{u_0^N} = f_i^N f_0^N \quad \text{for } i \in \{1, 2\}.$$

This means that $\Phi_{s_N} = \rho_0 \circ p \circ \tau^N$ where, as in Section 3.2, $\tau : \Gamma_2 \rightarrow \Gamma_2$ is the Dehn twist along the three curves a_i . By Proposition 3.3, for all $g \in \Gamma_2 \setminus \{1\}$ there exists $N \in \mathbf{N}$ such that $p \circ \tau^N(g) \neq 1$. Since ρ_0 is injective, this implies that $\Phi_{s_N}(g) \neq 1$ which shows that \mathcal{R} has the separation property.

4. Non-orientable surface groups

Theorem 4.1. *Let N_g be the fundamental group of a closed non-orientable surface of genus $g \geq 4$. There exists an injective morphism $N_g \rightarrow \text{Diff}(\mathbf{R}, 0)$.*

Remark 4.2. The fundamental group N_3 of the non-orientable surface of genus 3 is not fully residually free, and our methods do not apply to this group. (See [Lyn, Proposition 9].)

4.1. Even genus. We first treat the case of an even genus $g \geq 4$. In this case, the group N_g embeds in N_4 . Indeed, the non-orientable surface of genus 4 is the connected sum of a torus $\mathbf{R}^2/\mathbf{Z}^2$ with two projective planes $\mathbb{P}^2(\mathbf{R})$. Taking a cyclic cover of the torus of degree k , we get a surface homeomorphic to the connected sum of $\mathbf{R}^2/\mathbf{Z}^2$ with $2k$ copies of $\mathbb{P}^2(\mathbf{R})$, hence a non-orientable surface of genus $2(k+1)$. Thus, it suffices to prove that N_4 embeds in $\text{Diff}(\mathbf{R}, 0)$.

The non-orientable surface of genus 4 is homeomorphic to the connected sum of 4 copies of $\mathbb{P}^2(\mathbf{R})$, and this gives the presentation (see Figure 2)

$$(4.1) \quad N_4 = \langle a_1, a_2, b_1, b_2 \mid a_1^2 a_2^2 b_1^2 b_2^2 = 1 \rangle.$$

Let $p : N_4 \rightarrow \langle a_1, a_2 \rangle$ be the morphism fixing a_1, a_2 and sending b_1 and b_2 to a_1^{-1} and a_2^{-1} respectively. Let $\tau : N_4 \rightarrow N_4$ be the Dehn twist around the curve $\gamma = (a_1^2 a_2^2)^{-1}$, i.e., the automorphism that fixes a_1 and a_2 and sends b_1 and b_2 to $\gamma b_1 \gamma^{-1}$ and $\gamma b_2 \gamma^{-1}$ respectively.

Lemma 4.3. *Given any $g \in N_4 \setminus \{1\}$, there exists $n_0 \in \mathbf{N}$ such that for all $N \geq n_0$, $p \circ \tau^N(g) \neq 1$.*




FIGURE 2

The fundamental group N_4 . – The base point is represented by \bullet , the 4 generators are a_1, a_2, b_1, b_2 , and the curve γ is used to construct the Dehn twist τ .

For the proof. The proof of this statement is completely analogous to the proof of [BGSS, Corollary 2.2], using Baumslag Lemma, we leave it as an exercise to the reader. See also [CG, Proposition 4.13]. \square

Using Theorem 3.4, we fix two germs of diffeomorphisms f_1 and $f_2 \in \text{Diff}(\mathbf{R}, 0)$ generating a free group and satisfying $f_1'(0) > 1$ and $f_2'(0) > 1$. We denote by

$$(4.2) \quad \rho_0 : \mathbf{F}_2 = \langle a_1, a_2 \rangle \rightarrow \text{Diff}(\mathbf{R}, 0)$$

the injective morphism sending a_i to f_i for $i \in \{1, 2\}$. In particular,

$$(4.3) \quad \rho_0(\gamma) = (f_1^2 \circ f_2^2)^{-1}$$

is a hyperbolic germ: its derivative $\lambda = ((f_1^2 \circ f_2^2)'(0))^{-1}$ is < 1 . The Koenigs linearization theorem gives an element $h \in \text{Diff}(\mathbf{R}, 0)$ such that $\rho_0(\gamma) = h \circ m_\lambda \circ h^{-1}$. Consider the multiplicative flow $\varphi : \mathbf{R}_+^* \rightarrow \text{Diff}(\mathbf{R}, 0)$ defined by $\varphi^s = g \circ m_s \circ g^{-1}$. As above, $\varphi^\lambda = \rho_0(\gamma)$, φ^s commutes with $\rho_0(\gamma)$ for all $s > 0$, and $s \mapsto \varphi^s$ is a polynomial map: for all $k \in \mathbf{N}$, $s \mapsto A_k(\varphi^s)$ is a polynomial in the variables s and s^{-1} .

Set $\mathcal{R} = \mathbf{R}_+^*$. Given $s \in \mathbf{R}_+^*$, consider the morphism $\rho_s : N_4 \rightarrow \text{Diff}(\mathbf{R}, 0)$ defined by

$$\begin{array}{ll} a_1 \mapsto f_1 & a_2 \mapsto f_2 \\ b_1 \mapsto \varphi^s f_1^{-1} \varphi^{-s} & b_2 \mapsto \varphi^s f_2^{-1} \varphi^{-s}. \end{array}$$

This gives a well defined homomorphism because φ^s commutes with $f_1^2 f_2^2$.

We now check the three assumptions of Lemma 3.1. Clearly, \mathcal{R} is a Baire space. The irreducibility is a consequence of the fact that for any $g \in N_4$, and any $k \in \mathbf{N}$ the map $s \mapsto A_k(\varphi_s(g))$ is a polynomial function in the variables $s^{\pm 1}$. The separation property follows from Lemma 4.3 together with the fact that $\rho_{\lambda^N} = \rho_0 \circ p \circ \tau^N$ and that ρ_0 is injective.

4.2. Odd genus. We now treat the case of a non-orientable surface of odd genus $g = 2k + 1$, $k \geq 2$. One can write N_{2k+1} as (see Figure 3 below)

$$(4.4) \quad N_{2k+1} = \langle a_1, \dots, a_k, c, b_1, \dots, b_k \mid a_1^2 \dots a_k^2 c^2 b_k^2 \dots b_1^2 = 1 \rangle.$$

This group splits as a double amalgam of free groups

$$(4.5) \quad N_{2k+1} = \langle a_1, \dots, a_k \rangle_{a_1^2 \dots a_k^2 = \gamma^{-1}} \ast_{c^{-2}\gamma = b_k^2 \dots b_1^2} \langle \gamma, c \rangle \ast_{c^{-2}\gamma = b_k^2 \dots b_1^2} \langle b_1, \dots, b_k \rangle.$$

We shall use the following notation to refer to this amalgam structure:




FIGURE 3

The fundamental group N_{2k+1}

- $A_1 = \langle a_1, \dots, a_k \rangle$ and $e_{1,2} = (a_1^2 \dots a_k^2)^{-1}$;
- $A_2 = \langle \gamma, c \rangle$ and $e_{2,1} = \gamma$ and $e_{2,3} = c^{-2}\gamma = \delta$;
- $A_3 = \langle b_1, \dots, b_k \rangle$ and $e_{3,2} = b_k^2 \dots b_1^2$.

So, each of the A_i is a free group and the amalgamation is given by $e_{1,2} = e_{2,1}$ and $e_{2,3} = e_{3,2}$.

Define a morphism $p: N_{2k+1} \rightarrow \langle a_1, \dots, a_k \rangle \simeq \mathbf{F}_k$ by

$$\begin{aligned} a_i &\mapsto a_i & \text{for } i \leq k & & c &\mapsto a_k^{-2} \\ b_i &\mapsto a_i^{-1} & \text{for } i \leq k-1 & & b_k &\mapsto a_k \end{aligned}$$

(the structure of amalgam shows that p is well defined).

Lemma 4.4. *The morphism $p: N_{2k+1} \rightarrow \mathbf{F}_k$ is injective in restriction to each of the three subgroups of the amalgam (4.5).*

Proof. By construction, it is injective in restriction to $\langle a_1, \dots, a_k \rangle$ and in restriction to $\langle b_1, \dots, b_k \rangle$. Then, note that $p(\langle \gamma, c \rangle) = \langle a_1^2 \dots a_k^2, a_k^{-2} \rangle$ is isomorphic to \mathbf{F}_2 because it is a non-abelian subgroup of a free group. Since \mathbf{F}_2 is Hopfian, p is necessarily injective in restriction to $\langle \gamma, c \rangle$. \square

Consider $\delta = b_k^2 \dots b_1^2 = c^{-2}\gamma$ and note that $p(\delta) = a_k^2 a_{k-1}^{-2} \dots a_1^{-2}$. Let τ be the Dehn twist corresponding to the decomposition above, i.e. the automorphism fixing a_i , sending c to $\gamma c \gamma^{-1}$ and sending b_i to $(\gamma \delta) b_i (\gamma \delta)^{-1}$. Since τ is the composition of the twists given by γ and δ and these two twists commute we get

$$\tau^N(b) = (\gamma^N \delta^N) b (\gamma^N \delta^N)^{-1}, \quad \forall b \in A_3.$$

In this situation, one can prove the following lemma in a similar way to Proposition 3.3.

Lemma 4.5. *Given any $g \in N_{2g+1} \setminus \{1\}$, there exists $n_0 \in \mathbf{N}$ such that for all $N \geq n_0$, $p \circ \tau^N(g) \neq 1$.*

Proof. Write g as a word in the graph of groups, i.e., $g = s_0 \dots s_n$ with $s_k \in A_{r_k}$ (we allow $s_k = 1$) for some $r_k \in \{1, 2, 3\}$, with $r_{k+1} = r_k \pm 1$, and $r_0 = r_n = 1$. We take this word of minimal possible length among words satisfying these constraints. If k is such that $r_{k-1} = r_{k+1}$, then $s_k \notin \langle e_{r_k, r_{k+1}} \rangle$ since otherwise, one could shorten the word using the structure of amalgam (in particular $s_k \neq 1$ in this case). Now one easily checks that

$$(4.6) \quad \tau^N(g) = s_0 d_1^{\varepsilon_1 N} s_1 d_2^{\varepsilon_2 N} s_2 \dots d_n^{\varepsilon_n N} s_n$$

where $d_k = e_{r_{k-1}, r_k} \in \{\gamma, \delta\}$, and $\varepsilon_k = r_k - r_{k-1} \in \{\pm 1\}$.

We claim that $s_k^{-1} d_k s_k$ does not commute with d_{k+1} . If $d_k \neq d_{k+1}$, this follows from the fact that γ commutes with no conjugate of δ in $A_2 = \langle c, \delta \rangle$. If $d_k = d_{k+1}$, then $r_{k-1} = r_{k+1}$, so $s_k \notin \langle e_{r_k, r_{k+1}} \rangle = \langle d_k \rangle$. If $[s_k^{-1} d_k s_k, d_k] = 1$, then s_k preserves the axis of d_k in the Cayley graph of the free group A_{r_k} , so s_k is a power of d_k , because $d_k \in \{\gamma, \delta\}$ is not a proper power; this contradicts that $s_k \notin \langle d_k \rangle$.

Denote by $\bar{s}_k, \bar{d}_k \in \mathbf{F}_r$ the images of s_k, d_k under p . Since p is injective on each A_{r_k} , $\bar{s}_k^{-1} \bar{d}_k \bar{s}_k$ does not commute with \bar{s}_{k+1} , so the hypotheses of Baumslag Lemma apply to the word

$$(4.7) \quad p \circ \tau^N(g) = \bar{d}_0^{\varepsilon_0 N} \bar{s}_1 \bar{d}_1^{\varepsilon_1 N} \bar{s}_2 \dots \bar{d}_{n-1}^{\varepsilon_{n-1} N} \bar{s}_n \bar{d}_n^{\varepsilon_n N}$$

so $p \circ \tau^N(g) \neq 1$ for N large enough. □

Now consider k elements f_1, \dots, f_k of $\text{Diff}(\mathbf{R}, 0)$ generating a free group of rank k with $f_i'(0) > 1$ for all $i \in \{1, \dots, k\}$, and $f_k'(0) < f_1'(0)$. Such a set can be obtained from two generators g_1 and g_2 of a free group of rank 2 with $g_i'(0) > 1$, as in Theorem 3.4, by taking $f_i = g_1^i \circ g_2^2 \circ g_1^{-i}$ for $i < k$ and $f_k = g_1^k \circ g_2 \circ g_1^{-k}$. Let $\rho_0 : \mathbf{F}_k = \langle a_1, \dots, a_k \rangle \rightarrow \text{Diff}(\mathbf{R}, 0)$ be the injective morphism sending a_i to f_i for $i \leq k$. In particular, $\rho_0(\gamma) = (f_1^2 \circ \dots \circ f_k^2)^{-1}$ and $\rho_0(p(\delta)) = f_k^2 \circ f_{k-1}^{-2} \circ \dots \circ f_1^{-2}$ are hyperbolic. Using Koenigs linearization theorem as above, there exists two multiplicative flows φ and $\psi : \mathbf{R}_+^* \rightarrow \text{Diff}(\mathbf{R}, 0)$ and a pair of positive real numbers λ and μ such that (1) $\varphi^\lambda = \rho_0(\gamma)$ and $\psi^\mu = \rho_0(p(\delta))$, and (2) $s \mapsto \varphi^s$ and $s \mapsto \psi^s$ are polynomial mappings.

Set $\mathcal{R} = (\mathbf{R}_+^*)^2$ and, for every $(s, s') \in \mathcal{R}$, define a morphism $\rho_{s, s'} : \mathbf{N}_{2k+1} \rightarrow \text{Diff}(\mathbf{R}, 0)$ by

$$\begin{aligned} a_i &\mapsto f_i & \text{for } i \leq k & & c &\mapsto \varphi^s f_k^{-2} \varphi^{-s} \\ b_i &\mapsto \varphi^s \psi^{s'} f_i^{-1} (\varphi^s \psi^{s'})^{-1} & \text{for } i \leq k-1 & & b_k &\mapsto \varphi^s \psi^{s'} f_k (\varphi^s \psi^{s'})^{-1} \end{aligned}$$

(this is well defined because φ^s and $\psi^{s'}$ commute with $\rho_0(\gamma) = (f_1^2 \circ \dots \circ f_k^2)^{-1}$ and $\rho_0(p(\delta)) = f_k^2 \circ f_{k-1}^{-2} \circ \dots \circ f_1^{-2}$ respectively).

The assumptions of Lemma 3.1 hold: \mathcal{R} is a Baire space, and the irreducibility follows from the fact that the maps $s \mapsto \varphi_s$ and $s' \mapsto \varphi_{s'}$ are polynomials in the variables $s^{\pm 1}, s'^{\pm 1}$. The separation property follows from Lemma 4.5 together with the fact that $\rho_{\lambda^N, \mu^N} = \rho_0 \circ p \circ \tau^N$, and that ρ_0 is injective.

4.3. Embeddings in $\text{Diff}(\mathbf{k}, 0)$. The proofs just given provide the following statement.

Theorem B. *Let $(\mathbf{k}, |\cdot|)$ be a non-discrete and complete valued field.*

- (1) *Let Γ be the fundamental group of a closed orientable surface, or a closed non-orientable surface of genus ≥ 4 . Then, there is an embedding of Γ into $\text{Diff}(\mathbf{k}, 0)$.*
- (2) *Let $F \subset \text{Diff}(\mathbf{k}, 0)$ be a free group of rank 2, generated by two germs f and g with $|f'(0)| > 1$ and $|g'(0)| > 1$. Then, there is an embedding of Γ_2 , the fundamental group of a closed, orientable surface of genus 2, into $\text{Diff}(\mathbf{k}, 0)$ whose image contains F .*

Proof. For the first assertion, we just have to replace \mathbf{R} by \mathbf{k} in the proofs of Theorem 3.5 and 4.1. The parameter space is $\mathcal{R} = (\mathbf{k}^*)^3$ or \mathbf{k}^* or $(\mathbf{k}^*)^2$, and it is a Baire space because $(\mathbf{k}, |\cdot|)$ is complete.

For the second assertion, we start with a representation ρ_0 in Equation (3.8) whose image is equal to F . Remark 3.6 shows that all the injective morphisms Φ_s that we get satisfy also $\Phi_s(\Gamma_2) \supset F$. \square

Part II

5. The final topology on germs of diffeomorphisms

Let $(\mathbf{k}, |\cdot|)$ be a complete field. This section introduces a new topology on $\mathbf{k}\{z\}$ and $\text{Diff}(\mathbf{k}, 0)$, which will be used in our second proof of Theorem A. The reader may very well skip this section on a first reading.

5.1. The final topology over the complex numbers. Until Section 5.4, we focus on the case $\mathbf{k} = \mathbf{C}$. Let r be a positive real number. Consider the subalgebra \mathcal{A}_r of $\mathbf{C}\{z\}$ consisting of those power series $f(z) = \sum_n a_n z^n$ which converge on the open unit disk \mathbb{D}_r (i.e., $\text{rad}(f) \geq r$) and extend continuously to the closed unit disk $\overline{\mathbb{D}}_r$. When endowed with the norm

$$(5.1) \quad \|f\|_{\mathcal{A}_r} = \max_{z \in \overline{\mathbb{D}}_r} |f(z)|,$$

\mathcal{A}_r is a Banach algebra. If $s < r$, the restriction of functions $f \in \mathcal{A}_r$ to the smaller disk $\overline{\mathbb{D}}_s$ determines a 1-Lipschitz embedding $\mathcal{A}_r \rightarrow \mathcal{A}_s$.

The space $\mathbf{C}\{z\}$ is the union of the algebras \mathcal{A}_r and can be thus endowed with the final topology associated to the colimit

$$(5.2) \quad \mathbf{C}\{z\} = \varinjlim \mathcal{A}_r.$$

This means that a subset $\mathcal{U} \subset \mathbf{C}\{z\}$ is open if its intersection with \mathcal{A}_r is open for every $r > 0$. Equivalently, a map $\varphi: \mathbf{C}\{z\} \rightarrow X$ to a topological space is continuous if and only if its composition with the embedding $\mathcal{A}_r \rightarrow \mathbf{C}\{z\}$ is continuous for all r . Unless we say it explicitly, open sets, neighborhoods, and continuous maps refer, from now on, to this topology. A word of warning: for $r > s$, the inclusion $\mathcal{A}_r \rightarrow \mathcal{A}_s$ is not a homeomorphism to its image, and neither is the inclusion $\mathcal{A}_r \rightarrow \mathbf{C}\{z\}$.

The goal of this section is to obtain several basic properties of this topology. For instance, we are going to prove that there is a filtration of $\mathbf{C}\{z\}$ by compact subsets $\mathbf{C}_c\{z\}$ so that the continuity can be checked in restriction to each $\mathbf{C}_c\{z\}$.

Remark 5.1. If $s < r$, the homomorphism $\mathcal{A}_r \rightarrow \mathcal{A}_s$ is compact: by Montel theorem, the ball of radius 1 in \mathcal{A}_r is mapped into a compact subset K_1 of \mathcal{A}_s .

Let $K \subset \mathcal{A}_r$ be a bounded subset. Then, the closure $cl_s(K)$ of (the image of) K in \mathcal{A}_s is compact. If $t \leq s$, the image of $cl_s(K)$ in \mathcal{A}_t is compact, hence closed; this implies that $cl_s(K) = cl_t(K)$ in $\mathbf{C}\{z\}$. Thus, the closure \overline{K} of K in $\mathbf{C}\{z\}$ coincides with the closure $cl_s(K)$ of K in \mathcal{A}_s for any $s < r$. As a consequence, \overline{K} is compact.

We denote by $B_{\mathcal{A}_r}(\varepsilon)$ the open ball centred at 0 and of radius ε in \mathcal{A}_r , which we also view as a subset of $\mathbf{C}\{z\}$. If $s \leq r$ and $\varepsilon \leq \varepsilon'$, then $B_{\mathcal{A}_r}(\varepsilon) \subset B_{\mathcal{A}_s}(\varepsilon') \subset \mathbf{C}\{z\}$. Given any finite set of such balls $B_{\mathcal{A}_{r_j}}(\varepsilon_j)$, the sum $\sum_{j \geq 1}^n B_{\mathcal{A}_{r_j}}(\varepsilon_j)$ is the subset of $\mathbf{C}\{z\}$ whose elements are sums $f_1 + \cdots + f_n$ with $f_j \in B_{\mathcal{A}_{r_j}}(\varepsilon_j)$ for all j .

Lemma 5.2. *A subset \mathcal{U} of $\mathbf{C}\{z\}$ is a neighborhood of 0 if and only if there are decreasing sequences (r_n) and (ϵ_n) tending to 0 such that \mathcal{U} contains the set*

$$\mathcal{B} = \bigcup_n \sum_{j \geq 1}^n B_{\mathcal{A}_{r_j}}(\epsilon_j).$$

Lemma 5.2 shows that the topology defined in this section is the same as the topology introduced by Leslie in [Les], except that we consider germs of analytic functions at the origin in \mathbf{C} instead of real analytic functions on a compact analytic manifold.

Proof. First we argue that any set \mathcal{B} as in the statement of Lemma 5.2 is a neighborhood of 0 in $\mathbf{C}\{z\}$. To do so we need to check that $\mathcal{B} \cap \mathcal{A}_r$ contains a neighborhood of 0 for all r . The sum $\sum_{j=1}^n B_{\mathcal{A}_{r_j}}(\epsilon_j)$ is a subset of $\mathbf{C}\{z\}$ which is contained in \mathcal{A}_{r_n} . It is open in \mathcal{A}_{r_n} because one of the summands, namely $B_{\mathcal{A}_{r_n}}(\epsilon_n)$, is itself open. Now, the continuity of the inclusion $\mathcal{A}_r \rightarrow \mathcal{A}_{r_n}$ for $r_n < r$ implies that $\sum_{j=1}^n B_{\mathcal{A}_{r_j}}(\epsilon_j) \cap \mathcal{A}_r$ is also open in \mathcal{A}_r . Since $\sum_{j=1}^n B_{\mathcal{A}_{r_j}}(\epsilon_j) \cap \mathcal{A}_r$ is contained in $\mathcal{B} \cap \mathcal{A}_r$, the latter is a neighborhood of 0, as we needed to prove.

Suppose now that \mathcal{U} is a neighborhood of the origin in $\mathbf{C}\{z\}$, and fix a decreasing sequence (r_n) tending to 0. For each $n \geq 1$, set $\mathcal{U}_n = \mathcal{U} \cap \mathcal{A}_{r_n}$.

We first claim that there is a ball B_1 in \mathcal{A}_{r_1} such that $\overline{B_1} \subset \mathcal{U}$. Since \mathcal{U}_2 is open in \mathcal{A}_{r_2} , consider $\epsilon > 0$ such that $B_{\mathcal{A}_{r_2}}(\epsilon) \subset \mathcal{U}_2$. Now let $B_1 = B_{\mathcal{A}_{r_1}}(\epsilon/2)$. Then for all $\eta > 0$, $\overline{B_1} \subset B_1 + B_{\mathcal{A}_{r_2}}(\eta)$ so taking $\eta = \epsilon/2$, we get $\overline{B_1} \subset B_{\mathcal{A}_{r_2}}(\epsilon/2) + B_{\mathcal{A}_{r_2}}(\epsilon/2) \subset B_{\mathcal{A}_{r_2}}(\epsilon) \subset \mathcal{U}$, which proves our claim.

We now construct by induction open balls $B_n \subset \mathcal{A}_{r_n}$ such that for all n , $\overline{B_1} + \dots + \overline{B_n} \subset \mathcal{U}$. Given such a set of balls B_1, \dots, B_n , the set $K = \overline{B_1} + \dots + \overline{B_n}$ provides a compact subset of $\mathcal{A}_{r_{n+2}}$ contained in \mathcal{U}_{n+2} . Let ϵ be the distance from K to the complement of \mathcal{U}_{n+2} in $\mathcal{A}_{r_{n+2}}$; by compactness, $\epsilon > 0$, and $K + B_{\mathcal{A}_{r_{n+2}}}(\epsilon/2) \subset \mathcal{U}$. We then define $B_{n+1} = B_{\mathcal{A}_{r_{n+1}}}(\epsilon/4)$. Then $K + \overline{B_{n+1}} \subset K + B_{\mathcal{A}_{r_{n+2}}}(\epsilon/2) \subset \mathcal{U}$. This concludes the induction step and the proof. \square

5.2. Coefficient functions. Recall that the coefficients of $f \in \mathcal{A}_r$ can be computed via the Cauchy integral formula:

$$(5.3) \quad A_n(f) = \frac{1}{2\pi i} \int_{\{|z|=r\}} \frac{f(z)}{z^{n+1}} dz.$$

This implies that the linear form A_n is continuous on each algebra \mathcal{A}_r with operator norm $\|A_n\|_{\mathcal{A}_r^*} \leq \frac{1}{2\pi} r^{-(n+1)}$, i.e. $|A_n(f)| \leq \frac{1}{2\pi} r^{-(n+1)} \|f\|_{\mathcal{A}_r}$ for all $f \in \mathcal{A}_r$. Since the maps A_n separate points in $\mathbf{C}\{z\}$, we obtain:

Lemma 5.3. *For each $n \geq 0$, the map $A_n : \mathbf{C}\{z\} \rightarrow \mathbf{C}$ is continuous. The topological space $\mathbf{C}\{z\}$ is Hausdorff.*

More generally, we have:

Lemma 5.4. *If $\sum_n \theta_n z^n$ is a power series with infinite convergence radius, then the quantity*

$$(5.4) \quad \Theta(f) = \sum_n \theta_n |A_n(f)|$$

is well defined for every $f \in \mathbf{C}\{z\}$ and the function $\Theta : \mathbf{C}\{z\} \rightarrow \mathbf{R}_+$ is continuous.

Proof. The estimate $\|A_n\|_{\mathcal{A}_r^*} \leq \frac{1}{2\pi} r^{-(n+1)}$ implies that the map

$$(5.5) \quad \mathcal{A}_r \rightarrow \mathbf{C}, \quad f \mapsto \sum_n \theta_n |A_n(f)|$$

is continuous for any power series $\sum_n \theta_n z^n$ with convergence radius greater than $\frac{1}{r}$. By definition of the topology on $\mathbf{C}\{z\}$ we get that this map is continuous on the whole space if the power series in question has infinite convergence radius. \square

5.3. Another filtration. We now introduce another filtration of $\mathbf{C}\{z\}$. If c is any positive real number, we define

$$(5.6) \quad \mathbf{C}_c\{z\} = \{f \in \mathbf{C}\{z\} \text{ with } |A_n(f)| \leq c^{n+1} \text{ for all } n\}.$$

Then $\mathbf{C}_c\{z\} \subset \mathbf{C}_{c'}\{z\}$ for $c \leq c'$, and $\mathbf{C}\{z\}$ is the increasing union of all $\mathbf{C}_c\{z\}$.

Lemma 5.5. *$\mathbf{C}_c\{z\}$ is compact, and contained in \mathcal{A}_r for all $r < c^{-1}$. Every compact subset $\Lambda \subset \mathbf{C}\{z\}$ is contained in some $\mathbf{C}_c\{z\}$.*

By compactness, the topology on $\mathbf{C}_c\{z\}$ induced by \mathcal{A}_r and by $\mathbf{C}\{z\}$ agree.

Proof. From Lemma 5.3, we deduce that $\mathbf{C}_c\{z\}$ is closed in $\mathbf{C}\{z\}$. If $f \in \mathbf{C}_c\{z\}$ and $r < c^{-1}$, then

$$(5.7) \quad \|f\|_{\mathcal{A}_r} \leq \sum_n c^{n+1} r^n \leq \frac{c}{1-cr}$$

This means that $\mathbf{C}_c\{z\}$ is a bounded subset in \mathcal{A}_r . Since the inclusion $\mathcal{A}_r \rightarrow \mathcal{A}_s$ is compact for $r > s$, $\mathbf{C}_c\{z\}$ has compact closure in \mathcal{A}_s , hence in $\mathbf{C}\{z\}$. Since $\mathbf{C}_c\{z\}$ is closed, it is compact.

To prove the second assertion, assume by contradiction that there is a compact subset $\Lambda \subset \mathbf{C}\{z\}$ such that for every integer $m > 0$ there exists $f_m \in \Lambda \setminus \mathbf{C}_m\{z\}$.

By definition, there is an index $n_m \geq 0$ with $|A_{n_m}(f_m)| > m^{n_m+1}$. By Lemma 5.4, each individual coefficient is continuous and thus bounded on the compact Λ . It follows that n_m goes to $+\infty$ as m does. We can thus assume, passing to a subsequence if necessary, that the n_m 's are pairwise distinct.

Set $\theta_{n_m} = (\frac{1}{m})^{n_m}$, and $\theta_n = 0$ if n is not one of the indices n_m . Then $\theta_n^{1/n}$ converges towards 0 as n goes to $+\infty$, meaning that the power series $\sum_n \theta_n z^n$ has infinite convergence radius. By Lemma 5.4, the map $f \mapsto \Theta(f) = \sum_n \theta_n |A_n(f)|$ is continuous on $\mathbf{C}\{z\}$ and thus bounded on our compact set Λ . On the other hand we have

$$\Theta(f_m) \geq \theta_{n_m} |A_{n_m}(f_m)| \geq m.$$

This yields the desired contradiction. \square

Remark 5.6. Given $r > 0$, introduce

$$(5.8) \quad \mathcal{B}_r = \left\{ f \in \mathbf{C}\{z\} \mid \text{rad}(f) \geq r, \sup_{\mathbb{D}_r} |f| \leq \frac{1}{r} \right\}.$$

This is the closure in $\mathbf{C}\{z\}$ of a ball in \mathcal{A}_r and is therefore compact (Remark 5.1). There are functions c_1, c_2, r_1 , and $r_2 : \mathbf{R}_+^* \rightarrow \mathbf{R}_+^*$ such that

$$\mathbf{C}_{c_1(r)}\{z\} \subset \mathcal{B}_r \subset \mathbf{C}_{c_2(r)}\{z\} \text{ and } \mathcal{B}_{r_1(c)} \subset \mathbf{C}_c\{z\} \subset \mathcal{B}_{r_2(c)}.$$

It follows that one could equivalently state the results of this section in terms of the filtration $(\mathcal{B}_r)_{r>0}$ instead of $(\mathbf{C}_c\{z\})_{c>0}$.

The following corollary allows us to view the final topology on $\mathbf{C}\{z\}$ as the weak topology associated to the filtration by the compact sets $\mathbf{C}_c\{z\}$.

Corollary 5.7. *A subset $F \subset \mathbf{C}\{z\}$ is closed if and only if for all $c > 0$, $F \cap \mathbf{C}_c\{z\}$ is closed. A map $F : \mathbf{C}\{z\} \rightarrow X$ to a topological space is continuous if and only if its restriction to $\mathbf{C}_c\{z\}$ is continuous for all $c > 0$.*

Proof. Clearly, it suffices to prove the first assertion. If F is closed, so is $F \cap \mathbf{C}_c\{z\}$. Assume conversely that $F \cap \mathbf{C}_c\{z\}$ is closed for all $c > 0$, and let us prove that F is closed. By definition of the final topology, we need to prove that given $r > 0$, its preimage $j_r^{-1}(F)$ under the inclusion $j_r : \mathcal{A}_r \rightarrow \mathbf{C}\{z\}$ is closed. It suffices to prove that for any $R > 0$, its intersection with the ball $B_{\mathcal{A}_r}(R)$ is closed in \mathcal{A}_r . Since $B_{\mathcal{A}_r}(R)$ has compact closure, there exists $c > 0$ such that $B_{\mathcal{A}_r}(R) \subset \mathbf{C}_c\{z\}$. Since $F \cap \mathbf{C}_c\{z\}$ is closed, $B_{\mathcal{A}_r}(R) \cap j_r^{-1}(F) = B_{\mathcal{A}_r}(R) \cap j_r^{-1}(F \cap \mathbf{C}_c\{z\})$ is a closed subset of \mathcal{A}_r which concludes the proof. \square

Although one can show that the topology on $\mathbf{C}\{z\}$ is not metrizable, each space $\mathbf{C}_c\{z\}$ is a metric space. Being compact, the topology on $\mathbf{C}_c\{z\}$ can be described in many equivalent ways:

Proposition 5.8. *Let c be a positive real number. Let (f_m) be a sequence in $\mathbf{C}_c\{z\}$ and let f_∞ be an element of $\mathbf{C}_c\{z\}$. The following are equivalent:*

- (1) (f_m) converges to f_∞ in $\mathbf{C}_c\{z\}$;
- (2) for some (any) $r < c^{-1}$, (f_m) converges uniformly toward f_∞ on $\overline{\mathbb{D}}_r$;
- (3) (f_m) converges toward f_∞ uniformly on every compact subset of $\mathbb{D}_{c^{-1}}$;
- (4) for every index n , $A_n(f_m)$ converges toward $A_n(f_\infty)$.

Proof. As seen before, $\mathbf{C}_c\{z\}$ is contained in \mathcal{A}_r for all $r < c^{-1}$ and the topology induced by $\|\cdot\|_{\mathcal{A}_r}$ agrees with the topology induced by $\mathbf{C}_c\{z\}$. This proves the equivalence of the first three assertions.

To prove the equivalence with the last assertion, consider the map $\Phi : \mathbf{C}_c\{z\} \rightarrow [0, 1]^{\mathbb{N}}$ defined by $\Phi(f) = (\frac{A_n(f)}{c^{n+1}})_{n \in \mathbb{N}}$, where $[0, 1]^{\mathbb{N}}$ is endowed with the product topology. This map being continuous and injective, it is a homeomorphism to its image, and the result follows. \square

5.4. The final topology on a field with an absolute value. In this section, we explain that the final topology induced by the filtration $\mathbf{C}_c\{z\}$ makes sense for every field \mathbf{k} with an absolute value $|\cdot|$; but the results based on Montel theorem (Remark 5.1) may fail for fields $\mathbf{k} \neq \mathbf{C}$.

Let \mathbf{k} be a complete field \mathbf{k} for some absolute value $|\cdot| : \mathbf{k} \rightarrow \mathbf{R}_+$. By Ostrowski's Theorem, \mathbf{k} is either \mathbf{R} or \mathbf{C} , or the absolute value is non-archimedean: $|x + y| \leq \max(|x|, |y|)$ for all $x, y \in \mathbf{k}$. The algebra $\mathbf{k}\{z\}$ of convergent power series is filtrated by the family of subsets

$$(5.9) \quad \mathbf{k}_c\{z\} = \{f \in \mathbf{k}\{z\} \text{ with } |A_n(f)| \leq c^{n+1} \text{ for all } n\}$$

for $c > 0$. We endow $\mathbf{k}_c\{z\}$ with the product topology, via the embedding $f \in \mathbf{k}_c\{z\} \rightarrow (A_n(f))_n \in \mathbf{k}^{\mathbb{N}}$: a sequence $(f_k)_{k \in \mathbb{N}}$ of elements of $\mathbf{k}_c\{z\}$ converges to $f_\infty \in \mathbf{k}_c\{z\}$ if and only if $A_n(f_k) \rightarrow A_n(f_\infty)$ for all n . For $c \leq c'$, $\mathbf{k}_c\{z\}$ is closed in $\mathbf{k}_{c'}\{z\}$ and the inclusion is a homeomorphism to its image. We then endow $\mathbf{k}\{z\}$ with the topology associated to this filtration: a subset $F \subset \mathbf{k}\{z\}$ is closed if and only if $F \cap \mathbf{k}_c\{z\}$ is closed in $\mathbf{k}_c\{z\}$. Equivalently, a map $\varphi : \mathbf{k}\{z\} \rightarrow X$ to a topological space is continuous if and only if its restriction to $\mathbf{k}_c\{z\}$ is continuous for every $c > 0$. By construction, the maps $f \mapsto A_n(f)$ are continuous on $\mathbf{k}\{z\}$. Proposition 5.8 shows that, when $\mathbf{k} = \mathbf{C}$, this topology agrees with the final topology defined in Section 5.1.

If \mathbf{k} is locally compact, each $\mathbf{k}_c\{z\}$ is compact. In general, since $\mathbf{k}_c\{z\}$ is a countable product of complete metric spaces, we get:

Proposition 5.9. *If \mathbf{k} is a complete field, then $\mathbf{k}_c\{z\}$ is a metrizable complete space. In particular, it is a Baire space.*

On the other hand, $\mathbf{k}\{z\}$ is not a Baire space since it is a countable union of $\mathbf{k}_c\{z\}$, each of which is closed and has an empty interior.

5.5. The topological group of germs of diffeomorphisms. Any $f \in \text{Diff}(\mathbf{k}, 0)$ can be written as $f = \lambda(z + z^2\tilde{f})$ for some $\tilde{f} \in \mathbf{k}\{z\}$ or equivalently as

$$(5.10) \quad f = \lambda(z + \tilde{a}_2z^2 + \cdots + \tilde{a}_kz^k + \dots)$$

for some $\lambda \in \mathbf{k}^*$ and $\tilde{a}_n \in \mathbf{k}$. Thus, we define the maps $\tilde{A}_n : \text{Diff}(\mathbf{k}, 0) \rightarrow \mathbf{k}$ by

$$(5.11) \quad \tilde{A}_n(f) = A_n(f)/A_1(f) = \tilde{a}_n.$$

Given two real numbers $c > 0$ and $\lambda_0 > 1$, we define the two subsets

$$(5.12) \quad \text{Diff}_c(\mathbf{k}, 0) = \{f \in \text{Diff}(\mathbf{k}, 0) ; |\tilde{A}_n(f)| \leq c^{n-1} \text{ for all } n\}$$

and

$$(5.13) \quad \text{Diff}_{\lambda_0, c}(\mathbf{k}, 0) = \left\{ f \in \text{Diff}_c(\mathbf{k}, 0) ; \frac{1}{\lambda_0} \leq |A_1(f)| \leq \lambda_0 \right\}$$

Observe that if we denote by $m_\alpha : z \mapsto \alpha z$ the multiplication by some scalar $\alpha \in \mathbf{k}^*$ then we have

$$(5.14) \quad m_\alpha \text{Diff}_c(\mathbf{k}, 0) m_\alpha^{-1} = \text{Diff}_{c\alpha}(\mathbf{k}, 0)$$

and

$$(5.15) \quad m_\alpha \text{Diff}_{\lambda_0, c}(\mathbf{k}, 0) m_\alpha^{-1} = \text{Diff}_{\lambda_0, c\alpha}(\mathbf{k}, 0)$$

Lemma 5.10. *A map $\varphi : \text{Diff}(\mathbf{k}, 0) \rightarrow X$ to a topological space is continuous if and only if it is continuous in restriction to $\text{Diff}_c(\mathbf{k}, 0)$ (or equivalently to $\text{Diff}_{\lambda_0, c}(\mathbf{k}, 0)$) for every $c > 0$ and $\lambda_0 > 1$.*

Proof. It suffices to check the continuity of φ on the open set $U_{\lambda_0} = \{f ; \frac{1}{\lambda_0} < |A_1(f)| < \lambda_0\}$ for all $\lambda_0 > 1$. By definition of the final topology, it suffices to check its continuity on $U_{\lambda_0} \cap \mathbf{k}_c\{z\}$ for every $c > 1$. But $U_{\lambda_0} \cap \mathbf{k}_c\{z\}$ is a subset of $\text{Diff}_{\lambda_0, c'}(\mathbf{k}, 0)$ as soon as $c' \geq \max(\lambda_0, c^3)$; since we know that φ is continuous on $\text{Diff}_{\lambda_0, c'}(\mathbf{k}, 0)$, this proves the lemma. \square

Proposition 5.11. *If \mathbf{k} is a complete field, then $\text{Diff}_{\lambda_0, c}(\mathbf{k}, 0)$ and $\text{Diff}_c(\mathbf{k}, 0)$ are complete metric spaces. In particular, they are Baire spaces.*

Proof. By definition, $\text{Diff}_{c, \lambda_0}(\mathbf{k}, 0)$ is homeomorphic to a countable product of closed subsets of \mathbf{k} ; so, \mathbf{k} being complete, its topology is induced by a complete metric. Since \mathbf{k}^* is homeomorphic to the closed subset $\{(x, y) | xy - 1\} \subset \mathbf{k}^2$, the same argument applies to $\text{Diff}_c(\mathbf{k}, 0)$. \square

Theorem 5.12. *Let $(\mathbf{k}, |\cdot|)$ be a field with a complete absolute value. With the final topology, $\text{Diff}(\mathbf{k}, 0)$ is a topological group.*

Lemma 5.13. *For every real number $c > 1$, there exists a real number $c' > 1$ such that the following holds: if f and g are in $\text{Diff}_{c, c}(\mathbf{k}, 0)$, then $f \circ g$ and f^{-1} lie in $\text{Diff}_{c', c'}(\mathbf{k}, 0)$.*

Proof. Let $f = \lambda(z + \sum_{n \geq 2} \tilde{a}_n z^n)$, $g = \mu(z + \sum_{n \geq 2} \tilde{b}_n z^n)$ with $|\tilde{a}_n|, |\tilde{b}_n| \leq c^{n-1}$ and $|\lambda|, |\mu|, |\lambda|^{-1}, |\mu|^{-1} \leq c$. Let $F = c(z + \sum_{n \geq 2} c^{n-1} z^n) = \frac{cz}{1-cz} \in \mathbf{R}\{z\}$ so that the absolute value of the coefficients of f and g are bounded by the coefficients of F . Then the absolute value of the coefficients of $f \circ g = \sum_{n \geq 1} a_n (\sum_{m \geq 1} b_m z^m)^n$ are bounded by the coefficients of $F \circ F = \frac{cz(1-cz)}{1-cz-c^2z}$. Since $F \circ F$ has positive convergence radius, there exists $c' \geq c^2$ such that $\tilde{A}_n(F \circ F) \leq c'^{n-1}$ for all $n \geq 2$. The first assertion follows.

We now prove the second assertion. Let $f = \lambda(z + \sum_{n \geq 2} \tilde{a}_n z^n)$, and let $f^{-1} = \lambda^{-1}(z + \sum_{n \geq 2} \tilde{b}_n z^n)$. The inversion formula from Section 2.1 gives

$$\begin{aligned} |\tilde{b}_n| &\leq \frac{|\lambda|}{|\lambda|^n} \sum_{k_1, k_2, \dots} \frac{(n+1) \cdots (n-1+k_1+k_2+\dots)}{k_1! k_2! \cdots} \cdot |\tilde{a}_2|^{k_1} |\tilde{a}_3|^{k_2} \cdots \\ &\leq c^{n-1} \sum_{k_1, k_2, \dots} \frac{(n+1) \cdots (n-1+k_1+k_2+\dots)}{k_1! k_2! \cdots} (c)^{k_1} (c)^{2k_2} \cdots \\ &= c^{2n-2} \sum_{k_1, k_2, \dots} \frac{(n+1) \cdots (n-1+k_1+k_2+\dots)}{k_1! k_2! \cdots}. \end{aligned}$$

Thus, we have to bound the quantity

$$K_n := \sum_{k_1, k_2, \dots} \frac{(n+1) \cdots (n-1+k_1+k_2+\dots)}{k_1! k_2! \cdots}.$$

But the numbers K_n are the coefficients of the power series expansion of the reciprocal diffeomorphism g^{-1} of

$$g(z) = z - z^2 - z^3 - z^4 \cdots = z \left(2 - \frac{1}{1-z} \right) = \frac{z - 2z^2}{1-z}.$$

In close form, we obtain

$$g^{-1}(y) = \frac{(1+y)}{4} - \frac{1}{4}\sqrt{1-6y+y^2}.$$

Since g^{-1} has positive convergence radius, there exists c_0 such that for all $n \geq 2$, $K_n \leq c_0^{n-1}$ hence $|\tilde{b}_n| \leq (c_0 c^2)^{n-1}$ and the result follows. \square

Proof of Theorem 5.12. By definition of the topology, given $c > 0$, one only needs to check the continuity of the group laws in restriction to $\text{Diff}_{c,c}(\mathbf{k}, 0)$.

Since $A_n(f \circ g)$ and $A_n(f^{-1})$ are given by polynomials in the coefficients $\tilde{A}_i(f)$, $\tilde{A}_i(g)$, $A_1(f)^{\pm 1}$, $A_1(g)^{\pm 1}$, the maps $(f, g) \mapsto A_n(f \circ g)$ and $f \mapsto A_n(f^{-1})$ are continuous on $\text{Diff}_{c,c}(\mathbf{k}, 0)$. By Lemma 5.13 there exists c' such that for all $f, g \in \text{Diff}_{c,c}(\mathbf{k}, 0)$, $f \circ g$ and f^{-1} lie in $\text{Diff}_{c',c'}(\mathbf{k}, 0)$. Since the topology on $\text{Diff}_{c',c'}(\mathbf{k}, 0)$ is the product topology, the continuity of the coefficients implies the continuity of the group laws. \square

5.6. Other topologies. First, we would like to point out that there are other reasonable and useful topologies on $\mathbf{C}\{z\}$, but for which the group laws are not continuous. This the case for the so-called *Takens topology* [MRR, BT]; this is the topology induced by the distance

$$(5.16) \quad \text{dist}(f, g) = \sup_n |A_n(f) - A_n(g)|^{1/n}.$$

Note that in particular the convergence radius of $f - g$ is large if f and g are close to each other in the Takens topology, and this implies that the right translation $R_f: g \mapsto g \circ f$ is not continuous if the radius of convergence of f is finite. Indeed, a small perturbation $g(z) + \epsilon z$ is mapped to $R_f(g + \epsilon z) = g \circ f + \epsilon f$, and the difference ϵf is not small in the Takens topology because its radius of convergence does not depend on ϵ .

We comment now on another important topology on $\text{Diff}(\mathbf{C}, 0)$, but for which the Baire property fails. Let $\text{Jets}_\ell(\mathbf{C}, 0)$ be the group of ℓ -jets of diffeomorphisms $a_1 z + \dots + a_\ell z^\ell \pmod{(z^{\ell+1})}$, with $a_1 \neq 0$; it can be considered as a solvable algebraic group and thus as a solvable complex Lie group. Let

$$(5.17) \quad j_\ell: \text{Diff}(\mathbf{C}, 0) \rightarrow \text{Jets}_\ell(\mathbf{C}, 0)$$

denote the homomorphism that maps a power series $f = \sum_n a_n z^n$ to $\sum_{n=1}^\ell a_n z^n$. We can then define a topology on $\text{Diff}(\mathbf{C}, 0)$ (resp. on $\overline{\text{Diff}}(\mathbf{C}, 0)$): the weakest topology for which all projections j_ℓ are continuous. With this topology, $\text{Diff}(\mathbf{C}, 0)$ is a topological group, because the projections j_ℓ are homomorphisms. Moreover, a sequence (f_m) converges toward a germ of diffeomorphism g if and only if the coefficients $A_n(f_m)$ converge to $A_n(g)$ for all n . In other words, this is the

topology of simple convergence on the coefficients. In particular, $\text{Diff}(\mathbf{C}, 0)$ is not a closed subset of $\widehat{\text{Diff}}(\mathbf{C}, 0)$ for this topology. With this topology, $\widehat{\text{Diff}}(\mathbf{C}, 0)$ is a Baire space, but $\text{Diff}(\mathbf{C}, 0)$ is not (Proposition 5.11 fails if $\text{Diff}(\mathbf{C}, 0)$ is endowed with this topology).

5.7. Continuity in the Koenigs linearization Theorem. A contraction $f \in \text{Diff}(\mathbf{k}, 0)$ is an element with $|A_1(f)| < 1$. In this case, Koenigs theorem says that the unique formal diffeomorphism h_f tangent to the identity that conjugates f to the homothety $z \mapsto A_1(f)z$ has positive convergence radius. The following result shows that $f \mapsto h_f$ is continuous for the final topology on the set of contractions

$$(5.18) \quad \text{Cont}(\mathbf{k}, 0) = \{f \in \text{Diff}(\mathbf{k}, 0) \mid |A_1(f)| < 1\}.$$

Theorem 5.14. *Let \mathbf{k} be a field with a complete non-trivial absolute value. For every germ $f \in \text{Cont}(\mathbf{k}, 0)$, the unique formal diffeomorphism h_f such that*

$$h_f(f(z)) = A_1(f) \cdot h_f(z) \quad \text{and} \quad A_1(h_f) = 1$$

has positive convergence radius, and the map

$$h : f \in \text{Cont}(\mathbf{k}, 0) \mapsto h_f \in \text{Diff}(\mathbf{k}, 0)$$

is continuous for the final topology. The coefficients of h_f are polynomial functions with integer coefficients in the variables $A_i(f)$ and $(A_1(f)^j - 1)^{-1}$, for $i, j \geq 1$.

When $\mathbf{k} = \mathbf{C}$, it is shown in [Mil, Chapter 8] that h_f is convergent and its coefficients depend holomorphically on f . Theorem 5.14 is just a variation on this classical result.

Proof. We refer to [Sie] for the real and complex cases, and to [HY] for the non-archimedean ones.

The coefficients of h_f can be computed inductively and turn out to be polynomials with integer coefficients in the variables $A_i(f)$ and $(A_1(f)^j - 1)^{-1}$, for $i, j \geq 1$ (see for instance [Sie, Eq. 4]). If $|A_1(f)| \leq \alpha$ for some α in the interval $[0, 1[$, then

$$(5.19) \quad |(A_1(f))^n - 1| \geq 1 - \alpha$$

for all $n > 0$. By [Sie, Theorem 1] and [HY, Theorem 1] in the archimedean and non-archimedean cases respectively, h_f is convergent and for all $c, \lambda > 1$, there exists c' such that $h_f \in \text{Diff}_{c'}(\mathbf{k}, 0)$ if $f \in \text{Diff}_{\lambda, c}(\mathbf{k}, 0)$.

The topology on $\text{Diff}_{c'}(\mathbf{C}, 0)$ is the product topology on the coefficients. Since the coefficients of h_f are continuous functions of f , it follows that the restriction of $f \mapsto h_f$ to $\text{Diff}_{\lambda, c}(f)$ is continuous. By definition of the final topology, this proves the continuity of h . \square

6. A large irreducible component of the representation variety

This section describes our second proof strategy for Theorem A. For simplicity, we consider only the fundamental group of a closed orientable surface of genus 2, but we work over any field \mathbf{k} with a complete absolute value $|\cdot|$.

6.1. An irreducible set of representations. Using the presentation

$$(6.1) \quad \Gamma_2 = \langle a, b, \bar{a}, \bar{b} \mid [a, b] = [\bar{a}, \bar{b}] \rangle,$$

we get an identification

$$(6.2) \quad \text{Hom}(\Gamma_2; \text{Diff}(\mathbf{k}, 0)) = \{(f, g, \bar{f}, \bar{g}) \in \widehat{\text{Diff}}(\mathbf{k}, 0)^4 \mid [f, g] = [\bar{f}, \bar{g}]\}.$$

Let $\mathbb{X} \subset \text{Hom}(\Gamma_2, \text{Diff}(\mathbf{k}, 0))$ be the set of representations $\rho : \Gamma_2 \rightarrow \text{Diff}(\mathbf{k}, 0)$ such that $\rho(a)$ is tangent to Id and $\rho(b)$ is a contraction. As in Equation (5.18), we denote by $\text{Cont}(\mathbf{k}, 0)$ the set of contractions. For $c > 0$, we let $\mathbb{X}_c = \mathbb{X} \cap \text{Diff}_c(\mathbf{C}, 0)$. Set

$$(6.3) \quad \mathcal{R} = \text{Cont}(\mathbf{k}, 0) \times \text{Diff}(\mathbf{k}, 0) \times \text{Diff}(\mathbf{k}, 0),$$

$$(6.4) \quad \mathcal{R}(c) = \text{Cont}_c(\mathbf{k}, 0) \times \text{Diff}_c(\mathbf{k}, 0) \times \text{Diff}_c(\mathbf{k}, 0),$$

and denote by $\pi : \mathbb{X} \rightarrow \mathcal{R}$ the projection

$$(6.5) \quad \pi(\rho) = (\rho(b), \rho(\bar{a}), \rho(\bar{b}))$$

Proposition 6.1. *The map π is a homeomorphism for the final topology, and its inverse*

$$\pi^{-1} : (g, \bar{f}, \bar{g}) \mapsto (f, g, \bar{f}, \bar{g})$$

is a polynomial map, in the following sense: for each $n \in \mathbf{N}^*$, the map $(g, \bar{f}, \bar{g}) \mapsto A_n(f)$ is polynomial in (finitely many of) the variables $A_k(g)$, $A_k(\bar{f})$, $A_k(\bar{g})$, $A_1(g)^{-1}$, $A_1(\bar{f})^{-1}$, $A_1(\bar{g})^{-1}$ and $(A_1(g)^k - 1)^{-1}$ ($k \geq 1$).

Proof. The projection π is continuous because both \mathbb{X} and \mathcal{R} come with the topology induced by the same topology on $\text{Diff}(\mathbf{k}, 0)$.

Consider a triple $(g, \bar{f}, \bar{g}) \in \text{Cont}(\mathbf{k}, 0) \times \text{Diff}(\mathbf{k}, 0) \times \text{Diff}(\mathbf{k}, 0)$. Since $[\bar{f}, \bar{g}]$ is tangent to the identity, the germs g and $[\bar{f}, \bar{g}] \circ g$ have the same derivative $\lambda = A_1(g)$ at 0. Since $|\lambda| < 1$, we can apply Koenigs Theorem 5.14: we get two germs h_1 and $h_2 \in \text{Diff}(\mathbf{k}, 0)$ tangent to the identity such that

$$(6.6) \quad h_1 \circ g \circ h_1^{-1} = m_\lambda \quad \text{and} \quad h_2 \circ ([\bar{f}, \bar{g}] \circ g) \circ h_2^{-1} = m_\lambda$$

where $m_\lambda(z) = \lambda z$ is the multiplication by λ . Then, the map $f := h_2^{-1} \circ h_1$ conjugates g to $[\bar{f}, \bar{g}] \circ g$ so

$$f \circ g \circ f^{-1} = [\overline{f}, \overline{g}] \circ g \quad \text{and} \quad [f, g] = [\overline{f}, \overline{g}].$$

This means that one can define the preimage $\pi^{-1}(g, \overline{f}, \overline{g}) \in \text{Hom}(\Gamma_2, \text{Diff}(\mathbf{k}, 0))$ by the 4-tuple $(f, g, \overline{f}, \overline{g})$: the fact that $\pi^{-1} \circ \pi = \text{Id}_{\mathcal{R}}$ follows from uniqueness in Koenigs Theorem.

The continuity of π^{-1} is a consequence of the continuity of the conjugacy in Koenigs Theorem 5.14 and of the continuity of the map $(g, \overline{f}, \overline{g}) \mapsto [\overline{f}, \overline{g}] \circ g$. The fact that $A_n(f)$ is polynomial in the given variables is a direct consequence of the corresponding fact in Koenigs Theorem, and the fact that group operations are polynomial mappings. \square

We denote the inverse map π^{-1} by Φ :

$$(6.7) \quad \forall s \in \mathcal{R}, \quad \Phi_s = \pi^{-1}(s).$$

Thus, if $s = (g, \overline{f}, \overline{g})$, then Φ_s is the morphism $\Gamma_2 \rightarrow \text{Diff}(\mathbf{k}, 0)$ such that $\Phi_s(b) = g$, $\Phi_s(\overline{a}) = \overline{f}$, $\Phi_s(\overline{b}) = \overline{g}$, and $\Phi_s(a)$ is the unique germ of diffeomorphism f which is tangent to the identity and satisfies the relation $[f, g] = [\overline{f}, \overline{g}]$. To conclude the proof, our goal now is to prove that for every $c > 0$, the family of morphisms Φ_s , for $s \in \mathcal{R}(c)$, satisfies the assumptions of Lemma 3.1. Proposition 5.11 shows that $\mathcal{R}(c)$ is a Baire space. The following corollary proves the irreducibility of $\mathcal{R}(c)$.

Corollary 6.2. *For any $w \in \Gamma_2$, denote by $\mathcal{R}(c)_w \subset \mathcal{R}(c)$ the set of homomorphisms in $\mathcal{R}(c)$ that kill w . Then either $\mathcal{R}(c)_w = \mathcal{R}(c)$ or $\mathcal{R}(c)_w$ is a closed subset of $\mathcal{R}(c)$ with empty interior.*

Proof. Since the functions $s \in \mathcal{R} \mapsto A_k(\Phi_s(w)) - A_k(\text{Id})$ are continuous, $\mathcal{R}(c)_w$ is closed. Now, assume that $\mathcal{R}(c)_w \neq \mathcal{R}(c)$: there exists $k \geq 1$ and a point $s = (g_0, \overline{f}_0, \overline{g}_0)$ in $\mathcal{R}(c)$ such that $A_k(\Phi_s(w)) \neq A_k(\text{Id})$. According to Proposition 6.1 the map $s \mapsto A_k(\Phi_s(w)) - A_k(\text{Id})$ is a polynomial function in finitely many of

- the coefficients $A_n(g_0)$, $A_n(\overline{f}_0)$ and $A_n(\overline{g}_0)$ ($n \geq 1$),
- the inverses $A_1(g_0)^{-1}$, $A_1(\overline{f}_0)^{-1}$, $A_1(\overline{g}_0)^{-1}$ and $(A_1(g_0)^k - 1)^{-1}$ ($k \geq 1$)

(note that $A_1(g_0)$, $A_1(\overline{f}_0)$, $A_1(\overline{g}_0)$ and $(A_1(g_0)^k - 1)$ do not vanish on \mathcal{R}). Our assumption says that this function does not vanish identically on $\mathcal{R}(c)$. Assume that $\mathcal{R}(c)_w$ contains a non-empty open subset \mathcal{U} , and choose a point $s = (g_1, \overline{f}_1, \overline{g}_1)$ in \mathcal{U} . If $\mathbf{k} = \mathbf{R}$ or \mathbf{C} , we denote by $B_{\mathbf{k}}$ the interval $[0, 1] \subset \mathbf{R}$; in the non-archimedean case we set $B_{\mathbf{k}} = \{t \in \mathbf{k}, |t| \leq 1\}$. Then, we consider the convex combination

$$(6.8) \quad s_t = \left(g_t = tf_1 + (1-t)f_0, \bar{f}_t = t\bar{f}_1 + (1-t)\bar{f}_0, \bar{g}_t = t\bar{g}_1 + (1-t)\bar{g}_0 \right)$$

with t in $B_{\mathbf{k}}$. According to Lemma 6.3 below, g_t , \bar{f}_t and \bar{g}_t are in $\mathcal{R}(c')$ for some $c' \geq c$, and $t \mapsto s_t$ is continuous; thus $\{t ; s_t \in \mathcal{U}\}$ is an open neighborhood of 1.

The function $t \mapsto \mathcal{A}_n(\varphi_{s_t}(w)) - A_n(\text{Id})$ does not vanish for $t = 0$, it is the restriction of a rational function of the variable t to the interval $[0, 1]$, and it vanishes identically on the open set $\{t ; s_t \in \mathcal{U}\}$. This is a contradiction, which shows that the interior of $\mathcal{R}(c)_w$ is empty. \square

Let $B_{\mathbf{k}}$ be the interval $[0, 1] \subset \mathbf{R}$ if $\mathbf{k} = \mathbf{R}$ or \mathbf{C} , or the ball $\{t \in \mathbf{k}, |t| \leq 1\}$ in the non-archimedean case.

Lemma 6.3. *Let $f_0 \in \text{Diff}_{c_0}(\mathbf{k}, 0)$ and $f_1 \in \text{Diff}_{c_1}(\mathbf{k}, 0)$, and for $t \in \mathbf{k}$, let $f_t = (1-t)f_0 + tf_1$. Let $p \in \mathbf{k}$ be the value of t (if any) such that $f'_p(0) = 0$.*

If $c_0 \leq c_1$ and $c_0|f'_0(0)| \leq c_1|f'_1(0)|$, then for all $t \in B_{\mathbf{k}} \setminus \{p\}$, $f_t \in \text{Diff}_{c_1}(\mathbf{k}, 0)$.

Proof. Denote $\lambda_0 = |f'_0(0)|$ and $\lambda_1 = |f'_1(0)|$. By assumption, for all $n \geq 2$, $|A_n(f_0)| \leq \lambda_0 c_0^{n-1}$ and $|A_n(f_1)| \leq \lambda_1 c_1^{n-1}$.

Consider first the case $\mathbf{k} = \mathbf{R}$ or \mathbf{C} . Since $\lambda_0 c_0 \leq \lambda_1 c_1$, we get for all $t \in [0, 1]$,

$$(6.9) \quad |A_n(f_t)| \leq (1-t)\lambda_0 c_0^{n-1} + t\lambda_1 c_1^{n-1}$$

$$(6.10) \quad \leq (1-t)\lambda_1 c_1 c_0^{n-2} + t\lambda_1 c_1^{n-1} \leq \lambda_1 c_1^{n-1}.$$

This shows that $f_t \in \text{Diff}_{c_1}(\mathbf{k})$ as soon as $f'_t(0) \neq 0$.

In the non-archimedean case, one has $|t-1| \leq 1$ for $t \in B_{\mathbf{k}}$. Similarly, we get

$$(6.11) \quad |A_n(f_t)| \leq \max\{|1-t|\lambda_0 c_0^{n-1}, |t|\lambda_1 c_1^{n-1}\}$$

$$(6.12) \quad \leq \max\{\lambda_1 c_1 c_0^{n-2}, \lambda_1 c_1^{n-1}\} \leq \lambda_1 c_1^{n-1}.$$

This shows that $f_t \in \text{Diff}_{c_1}(\mathbf{k})$ as soon as $f'_t(0) \neq 0$. Then, the continuity follows from the continuity of the coefficients $t \mapsto A_n(f_t)$. \square

6.2. Separation. To conclude the proof, we fix $c > 0$ and prove that $\mathcal{R}(c)$ satisfies the separation condition of Lemma 3.1. We thus fix $g \in \Gamma_2 \setminus \{1\}$ and show that $\mathcal{R}(c)$ contains a representation that does not kill g . Write the orientable surface group of genus 2 as $\Gamma_2 = \langle a, b, \bar{a}, \bar{b} \mid [a, b] = [\bar{a}, \bar{b}] \rangle$, and let $p : \Gamma_2 \rightarrow \langle a, b \rangle$ be the morphism fixing a and b and sending \bar{a} and \bar{b} to a and b respectively. Let $\tau : \Gamma_2 \rightarrow \Gamma_2$ be the Dehn twist around the curve $c = [a, b]$, i.e., the automorphism that fixes a, b and sends \bar{a} and \bar{b} to $c\bar{a}c^{-1}$ and $c\bar{b}c^{-1}$.

respectively. According to Proposition 1.2, there exists a positive integer n_0 such that $p \circ \tau^N(g) \neq 1$ for all $N \geq n_0$.

Apply Theorem 3.4 to get a pair f_1, f_2 of germs of diffeomorphisms generating a free group $\langle f_1, f_2 \rangle$ of rank 2 and satisfying $f_1'(0) > 1$ and $f_2'(0) > 1$. Define a morphism $\rho : \langle a, b \rangle \rightarrow \text{Diff}(\mathbf{k}, 0)$ by $\rho(a) = [f_1, f_2]$ and $\rho(b) = f_2^{-1}$. Then ρ is injective, $\rho(a)$ is tangent to the identity, and $\rho(b)$ is a contraction. Set $\rho_N := \rho \circ p \circ \tau^N$. For $N \geq n_0$ $\rho_N(g) \neq 1$. Thus, $\pi(\rho_N)$ lies in $\mathcal{R} \setminus \mathcal{R}_g$; but it might not lie in $\mathcal{R}(c)$.

Let $c_N > 0$ be such that $\pi(\rho_N) \in \mathcal{R}(c_N)$. Given $\alpha \in \mathbf{k}^*$, let ad_α be the inner automorphism of $\text{Diff}(\mathbf{k}, 0)$ given by $f \mapsto m_\alpha \circ f \circ m_\alpha^{-1}$. As noticed in Equation (5.12), we have $\text{ad}_\alpha(\text{Diff}_{c_N}(\mathbf{k}, 0)) = \text{Diff}_{\alpha c_N}(\mathbf{k}, 0)$. Thus, the representation $\rho'_N = \text{ad}_\alpha \circ \rho_N$ satisfies $\pi(\rho'_N) \in \mathcal{R}(c)$ if α is sufficiently small. Since $\rho'_N(g) \neq 1$ this concludes that $\mathcal{R}(c)$ satisfies the separation condition of Lemma 3.1.

6.3. Conclusion. The family of representations Φ_s , with $s \in \mathcal{R}(c)$ satisfies the Baire property, the irreducibility property, and the separation property of Lemma 3.1. This lemma implies that a generic element of $s \in \mathcal{R}(c)$ gives an embedding $\Phi_s : \Gamma_2 \rightarrow \text{Diff}(\mathbf{k}, 0)$, proving Theorem A for the group Γ_2 .

Part III

7. A p -adic proof

7.1. Free groups with integer coefficients. A theorem of White [Whi] shows that the homeomorphisms of \mathbf{R} defined by $f : z \mapsto z + 1$ and $g : z \mapsto z^3$ generate a free group. Conjugating the maps f and gfg^{-1} by $z \mapsto \frac{1}{3z}$, as in [Gla]¹, one gets two formal diffeomorphisms

$$(7.1) \quad \begin{aligned} f_0(z) &= \frac{z}{1 + 3z} = \sum_{n=1}^{\infty} (-3)^{n-1} z^n \\ g_0(z) &= \frac{z}{(1 + (3z)^3)^{1/3}} = \sum_{n=0}^{\infty} \binom{-1/3}{n} 3^{3n} z^{3n+1} \end{aligned}$$

that generate a non-abelian free group $\langle f_0, g_0 \rangle \subset \widehat{\text{Diff}}(\mathbf{Q}, 0) \subset \widehat{\text{Diff}}(\mathbf{k}, 0)$. It is remarkable that f_0 and g_0 are tangent to the identity at the origin and have integer coefficients:

¹This conjugacy is called the ‘‘Wilson trick’’ in [Gla].

Theorem 7.1. *The group $\widehat{\text{Diff}}(\mathbf{Q}, 0)$ contains a non-abelian free group, all of whose elements are tangent to the identity and have integer coefficients.*

Thus, one can produce an explicit free group in $\widehat{\text{Diff}}(\mathbf{k}, 0)$ for every field \mathbf{k} of characteristic 0. In characteristic $p > 0$, Szegedy proved that almost every pair of elements in the Nottingham group $\widehat{\text{Diff}}(\mathbf{Z}/p\mathbf{Z}, 0)$ generates a free group [Sze].

7.2. Subgroups of $\text{Diff}(\mathbf{Q}_p, 0)$. In this section, p is a prime number, and \mathbf{Q}_p is the field of p -adic numbers, with its absolute value $|\cdot|$ normalized by $|p| = 1/p$.

Let G_p denote the set of elements $f = \sum_{n \geq 1} a_n z^n$ in $\text{Diff}(\mathbf{Q}_p, 0)$ such that

$$(7.2) \quad a_n \in \mathbf{Z}_p \quad \forall n \quad \text{and} \quad |a_1| = 1.$$

Every element $f \in G_p$ satisfies $\text{rad}(f) \geq 1$. The ultrametric inequality and the Inversion formula show that G_p is a subgroup of $\text{Diff}(\mathbf{Q}_p, 0)$. With the product topology on coefficients (as in Section 5.6), it is a compact topological group, and the morphism $j_\ell: G_p \rightarrow \text{Jets}_\ell(\mathbf{Q}_p, 0)$ is continuous for every integer $\ell \geq 1$. The kernel of j_ℓ will be denoted $G_{p,\ell}$.

From Theorem 7.1, we know that G_p contains a free group of rank two generated by two germs f_0 and g_0 whose coefficients are in \mathbf{Z} .

Corollary 7.2. *Let p be a prime number and ℓ be a positive integer. The group $G_{p,\ell}$ contains a non-abelian free group. The group G_p contains a free group $\langle f, g \rangle$ of rank 2 such that $A_1(f)$ is a transcendental number while g is tangent to the identity up to order ℓ .*

Proof. Start with a non-abelian free group F in G_p . Since the group of jets $\text{Jets}_\ell(\mathbf{Q}_p, 0)$ is solvable, the restriction of j_ℓ to F is not injective. Its kernel is a free group (as any subgroup of F), and if ℓ is large it is not cyclic. Thus, the kernel is a non-abelian free group. This proves the first statement.

Set $\mathcal{R} = \{t \in \mathbf{Z}_p; |t| = 1\}$. Now, take a pair of generators f_0 and g_0 of a free group of rank 2 in $G_{p,\ell}$, and for $t \in \mathcal{R}$ consider the family of representations $\rho_t: \mathbf{F}_2 = \langle a, b \rangle \rightarrow G_p$ defined by $\rho_t(a) = m_t \circ f_0$ and $\rho_t(b) = g_0$ (here, as usual, $m_t(z) = tz$). If w is an element of \mathbf{F}_2 , and n is a positive integer, then $A_n(\rho_t(w))$ is a polynomial function in t and $1/t$ (see Section 2.1). If $w \neq 1$, there is an integer $n \geq 1$ such that $A_n(\rho_1(w)) \neq A_n(\text{Id})$. Thus, the set $\mathcal{R}_w \subset \mathcal{R}$ of parameters s such that $\rho_s(w) = \text{Id}$ is finite, the union $\cup_{w \neq 1} \mathcal{R}_w$ is at most countable, and there are transcendental numbers in its complement. For such a parameter t , ρ_t is injective and $A_1(\rho_t(a)) = t$ is transcendental. \square

Now, we apply the result of [BGSS] described in Section 1.2 to get:

Theorem 7.3. *Let p be a prime number. Let $\Gamma_2 = \langle a, b, \bar{a}, \bar{b} \mid [a, b] = [\bar{a}, \bar{b}] \rangle$ be the fundamental group of a closed orientable surface of genus 2. Then*

- (1) *For every integer $\ell \geq 1$, the group Γ_2 embeds in the compact group $G_{p,\ell}$.*
- (2) *There is an embedding $\rho: \Gamma_2 \rightarrow G_p$ such that $\rho(a)'(0) = \rho(\bar{a})'(0)$ is a transcendental number while $\rho(b)$ and $\rho(\bar{b})$ are tangent to the identity up to order ℓ .*

7.3. Back to complex coefficients. The field \mathbf{Q}_p , and thus the ring \mathbf{Z}_p , embeds (although not continuously) into \mathbf{C} ; such an embedding induces an embedding, coefficient by coefficient, of $\mathbf{Z}_p[[z]]$ into $\mathbf{C}[[z]]$. Thus, the surface groups constructed in Theorem 7.3 provide surface groups in $\widehat{\text{Diff}}(\mathbf{C}, 0)$. This construction does not preserve the convergence of power series, but it preserves the order of tangency to Id. Since there are transcendental complex numbers with modulus < 1 , we obtain:

Corollary 7.4. *Let ℓ be a positive integer. There is an embedding $\rho: \Gamma_2 \rightarrow \widehat{\text{Diff}}(\mathbf{C}, 0)$ such that $|\rho(a)'(0)| = |\rho(\bar{a})'(0)| < 1$ while $\rho(b)$ and $\rho(\bar{b})$ are tangent to the identity up to order ℓ .*

We can now prove the following version of Theorem A. This will be our third and last proof of it.

Theorem 7.5. *There is an embedding of Γ_2 in $\text{Diff}(\mathbf{C}, 0)$ such that $|\rho(a)'(0)| = |\rho(\bar{a})'(0)| < 1$ while $\rho(b)$ and $\rho(\bar{b})$ are tangent to the identity up to order ℓ .*

Proof. The first step is to choose a sequence $\mathcal{C} = (a_1, a_2, a_3, \dots)$ of complex numbers such that

- (a) the set $\{a_1, a_2, \dots\}$ is algebraically free: if $m \geq 1$ and $P \in \mathbf{Z}[x_1, \dots, x_m]$, and if $P(a_1, \dots, a_m) = 0$, then $P = 0$;
- (b) $|a_n| \leq 2^{-n}$ for all $n \geq 1$.

Such a sequence exists because \mathbf{C} is uncountable. Concrete examples can be obtained from the Lindemann-Weierstrass theorem (see also [Wal] for the constructions of von Neumann, Perron, Kneser, and Durand of uncountably many, algebraically free complex numbers). We shall consider the a_i as indeterminates for the field of rational functions $\mathbf{Q}(a_1, a_2, \dots)$. Armed with such a set we consider the following three formal diffeomorphisms

$$(7.3) \quad g = a_1 z + \sum_{i=1}^{\infty} a_{3i+1} z^{i+1}, \quad \bar{f} = z + \sum_{i=\ell}^{\infty} a_{3i+2} z^{i+1}, \quad \bar{g} = z + \sum_{i=\ell}^{\infty} a_{3i+3} z^{i+1}.$$

From the decay relation (b), these three power series have a positive radius of convergence. Since $|a_1| \leq 1/2$, the Koenigs linearization theorem gives a unique element $\bar{f} \in \text{Diff}(\mathbf{C}, 0)$ with $\bar{f}'(0) = 1$ such that

$$(7.4) \quad fgf^{-1} = [\bar{f}, \bar{g}]g.$$

The four elements (f, g, \bar{f}, \bar{g}) determine a representation φ of Γ_2 into $\text{Diff}(\mathbf{C}, 0)$.

Let us prove that this representation is faithful. Fix a non-trivial element w of Γ_2 , and write it as a word in a, b, \bar{a}, \bar{b} and their inverses. For every integer n , the coefficient $A_n(\varphi(w))$ is a polynomial function $Q_{w,n}$ in the variables a_n (for $n \geq 1$), a_1^{-1} , and the $(a_1^k - 1)^{-1}$ (for $k \geq 1$) with integer coefficients.

Now, take a faithful representation $\rho: \Gamma_2 \rightarrow \widehat{\text{Diff}}(\mathbf{C}, 0)$ that satisfies the conclusion of Corollary 7.4. There is an integer $n \geq 1$ such that $A_n(\rho(w)) \neq A_n(\text{Id})$. This implies that $Q_{w,n} \neq A_n(\text{Id})$ when we specialize the indeterminates a_i to the coefficients of the generators $\rho(\bar{a}), \rho(b)$, and $\rho(\bar{b})$. Since $Q_{w,n} \neq A_n(\text{Id})$, $\rho(w) \neq \text{Id}$ and ρ is the identity. \square

Part IV

8. Complements and open questions

8.1. Takens' theorem and smooth diffeomorphisms. To conclude this chapter, we mention the following result which allows to realize *any* faithful representation of a surface group in the group of formal germs as a group of C^∞ germs. Note that the p -adic method provides many embeddings of surface groups in $\widehat{\text{Diff}}(\mathbf{R}, 0)$ (see Corollary 7.4).

Recall that Γ_g denotes the fundamental group of the closed orientable surface of genus g .

Theorem C. *Let $\hat{\rho}: \Gamma_g \rightarrow \widehat{\text{Diff}}(\mathbf{R}, 0)$ be a faithful representation of the surface group Γ_g in the group of formal diffeomorphisms in one real variable. Then, there exists a faithful representation $\rho: \Gamma_g \rightarrow \text{Diff}^\infty(\mathbf{R}, 0)$ into the group of germs of C^∞ -diffeomorphisms such that the Taylor expansion of $\rho(w)$ coincides with $\hat{\rho}(w)$ for every $w \in \Gamma_g$.*

The proof will be a consequence of the following result (this theorem is easily derived from the Sternberg linearization theorem and Theorem 2 of [Tak]):

Theorem 8.1 (Sternberg [Ste], Takens, [Tak]). *Let $f, g: (\mathbf{R}, 0) \rightarrow (\mathbf{R}, 0)$ be two germs of C^∞ -diffeomorphisms, and let \hat{f} and \hat{g} denote their Taylor expansions. Suppose that f is not flat to the identity, that is $\hat{f} \neq \text{Id}$. Then, if \hat{f} and \hat{g} are conjugate by a formal diffeomorphism \hat{h} , there exists a germ of C^∞ -diffeomorphism $h: (\mathbf{R}, 0) \rightarrow (\mathbf{R}, 0)$ such that*

- *the Taylor expansion of h coincides with \hat{h} ;*
- *h conjugates f to g .*

Proof of Theorem C. Denote by $\hat{a}_i, \hat{b}_i, 1 \leq i \leq g$ the images of the standard generators of Γ_g by the representation ρ ; they satisfy the relation

$$(8.1) \quad \hat{a}_1 \circ \hat{b}_1 \circ \hat{a}_1^{-1} = \left(\prod_{j=2}^g [\hat{a}_j, \hat{b}_j] \right) \circ \hat{b}_1.$$

By the theorem of Borel and Peano, one can find germs of diffeomorphisms b_1 and $a_j, b_j, j \geq 2$, whose respective Taylor expansions coincide with \hat{b}_1, \hat{a}_j , and \hat{b}_j respectively. Then, Theorem 8.1 provides a germ of diffeomorphism a_1 such that $a_1 \circ b_1 \circ a_1^{-1} = \left(\prod_{j=2}^g [a_j, a_j] \right) \circ b_1$. Thus, one gets a representation ρ of Γ_2 into $\text{Diff}^\infty(\mathbf{R}, 0)$ with Taylor expansion equal to $\hat{\rho}$. Since the initial representation $\hat{\rho}$ is injective, so is ρ . \square

8.2. Conjugacy classes. Two subgroups Γ and Γ' of $\text{Diff}(\mathbf{C}, 0)$ are *topologically conjugate* if there is a germ of homeomorphism $\varphi: (\mathbf{C}, 0) \rightarrow (\mathbf{C}, 0)$ such that $\varphi \circ \Gamma \circ \varphi^{-1} = \Gamma'$, and are *formally conjugate* if there is a formal diffeomorphism $\hat{\varphi}$ such that $\hat{\varphi} \circ \Gamma \circ \hat{\varphi}^{-1} = \Gamma'$. A germ of homeomorphism φ is *anti-holomorphic* if its complex conjugate $z \mapsto \overline{\varphi(z)}$ is holomorphic.

Theorem 8.2 (Nakai, Cerveau-Moussu). *Let Γ and Γ' be two subgroups of $\text{Diff}(\mathbf{C}, 0)$ which are not solvable.*

- (1) *If φ is a local homeomorphism that conjugates Γ to Γ' , then φ is holomorphic, or anti-holomorphic.*
- (2) *If $\hat{\varphi}$ is a formal conjugacy between Γ and Γ' , then $\hat{\varphi}$ converges and is therefore a holomorphic conjugacy.*

Thus, (the images of) two embeddings of Γ_g in $\text{Diff}(\mathbf{C}, 0)$ are topologically or formally conjugate if and only if they are analytically conjugate.

8.3. Two questions.

8.3.1. It would be interesting to exhibit an embedding α of the group Γ_g , $g \geq 2$, into the group of analytic diffeomorphisms of the circle \mathbf{R}/\mathbf{Z} fixing the origin $o \in \mathbf{R}/\mathbf{Z}$. If such an embedding exists, the suspension of this representation α gives a compact manifold M_α of dimension 3 that fibers over Σ_g , together with a foliation \mathcal{F}_α of co-dimension 1 which is transverse to the fibration $\pi: M_\alpha \rightarrow \Sigma_g$ and whose monodromy is given by τ . The fixed point gives a compact leaf of \mathcal{F}_α with holonomy given by the same representation τ .

Question. Does there exist an embedding of Γ_2 into the group of analytic diffeomorphisms of the circle fixing the origin ?

This question was the original motivation of Cerveau and Ghys when they asked for a proof of Theorem A (see [Cer]).

Remark 8.3. According to Theorem 7.3, there is an embedding ρ of Γ_2 in $\text{Diff}(\mathbf{Q}_p, 0)$ such that $\rho(a)'(0)$ and $\rho(\bar{a})'(0)$ have modulus 1 while $\rho(b)$ and $\rho(\bar{b})$ are tangent to the identity. Conjugate ρ by the homothety $a \mapsto p^N z$ for some positive integer N . If N is large enough, the coefficients a_n , $n \geq 2$, of all elements of $\rho(\Gamma_2)$ have norm < 1 , and the ultrametric inequality shows that $\rho(\Gamma_2)$ preserves the open disks $\{z \in \mathbf{C}_p; |z| < 1 - \epsilon\}$ for every $\epsilon > 0$. Thus, it preserves arbitrary thin annuli $\{z \in \mathbf{C}_p; 1 - \epsilon|z| \leq 1\}$. (Here \mathbf{C}_p is the completion of the algebraic closure of \mathbf{Q}_p .)

A related, but a priori simpler question is: does there exist an embedding of Γ_2 into the group of increasing, real analytic diffeomorphisms of $[0, 1]$ fixing 0 and 1 ? Here, we demand that the diffeomorphisms extend to germs of real analytic diffeomorphisms on neighbourhoods of 0 and 1. If we replace real analytic diffeomorphisms by C^∞ diffeomorphisms, interesting examples have been constructed in [MS]. We refer to the introduction of [MS] for a description of the difficulties in trying to apply the strategy of [BGSS]: this is related to the question of deciding when a diffeomorphism f of $[0, 1]$ is contained in the flow of a smooth vector field, hence to Mather's invariant (see [EBN, Yoc]).

8.3.2. The derived subgroup of $\text{Diff}(\mathbf{C}, 0)$ is the kernel of the morphism $j_1: f \mapsto f'(0)$:

Theorem 8.4. *Let \mathbf{k} be a complete, non-discrete valued field. An element f of $\text{Diff}(\mathbf{k}, 0)$ is a commutator if and only if $f'(0) = 1$. All higher terms of the lower central series coincide with the kernel of $j_1: \text{Diff}(\mathbf{k}, 0) \rightarrow \text{Jets}_1(\mathbf{k}, 0)$.*

Proof. If f is a commutator $[g, h]$ then $f'(0) = 1$. If $f'(0) = 1$, compose f with the homothety $m_\lambda(z) = \lambda z$ for some $\lambda \in \mathbf{k}^*$ of norm $|\lambda| \neq 1$, and

apply Koenigs linearization theorem to find an element $h \in \text{Diff}(\mathbf{k}, 0)$ such that $m_\lambda \circ f = h \circ m_\lambda h^{-1}$ and $h'(0) = 1$. Then $f = [h, m_\lambda]$. This proves that the derived subgroup of $\text{Diff}(\mathbf{k}, 0)$ is the kernel of j_1 ; since h is in the kernel of j_1 , all subsequent terms of the lower central series coincide with the derived subgroup. \square

Now, consider the upper central series. The first terms are $\text{Diff}(\mathbf{C}, 0)$ and its derived subgroup $\text{Diff}(\mathbf{k}, 0)^{(1)}$. Then comes

$$(8.2) \quad \text{Diff}(\mathbf{k}, 0)^{(2)} := [\text{Diff}(\mathbf{k}, 0)^{(1)}, \text{Diff}(\mathbf{k}, 0)^{(1)}].$$

The group of jets of order 3 which are tangent to identity, i.e., jets of the form $j(z) = z + a_2 z^2 + a_3 z^3$ modulo z^4 , is an abelian group; at the level of formal germs, it is known that the kernel of j_3 in $\widehat{\text{Diff}}(\mathbf{k}, 0)^{(1)}$ coincides with the derived subgroup $\widehat{\text{Diff}}(\mathbf{k}, 0)^{(2)}$ (see [Cam], §3, for the description of the upper central series of $\widehat{\text{Diff}}(\mathbf{k}, 0)$). We don't know if a similar statement holds for germs of diffeomorphisms:

Question. Does the kernel of j_3 coincide with the second derived subgroup of $\text{Diff}(\mathbf{C}, 0)^{(2)}$? More generally, what is the upper central series of $\text{Diff}(\mathbf{C}, 0)$?

9. Appendix: Free groups

The following theorem, and its proof, are strongly inspired by [MRR]. The proof given in [MRR] is somewhat difficult because it makes use of a topology on $\text{Diff}(\mathbf{C}, 0)$ which is not compatible with the group law. We adapt the same proof, without reference to such a topology.

Theorem 9.1. *Let $(\mathbf{k}, |\cdot|)$ be a complete, non-discrete valued field. Let f and g be elements of $\text{Diff}(\mathbf{k}, 0)$ of infinite order. Let w be a non-trivial element of the free group \mathbf{F}_2 . Then, there is a polynomial germ of diffeomorphism h such that $w(hf h^{-1}, g) \neq \text{Id}$.*

If $w = a^{n_\ell} b^{n_{\ell-1}} \dots a^{n_2} b^{n_1}$, one can choose h of the form $z + \epsilon z^2 P(z)$ with an arbitrarily small ϵ and a polynomial function $P \in \mathbf{k}[z]$ such that $\deg(P) \leq (2\ell)!$ and $|P(x)| \leq 1$ for all $x \in \mathbb{D}_1$.

Before proving this result, let us introduce some vocabulary and notation. Write w as a reduced word in the generators a and b of the free group:

$$(9.1) \quad w = a^{n_\ell} b^{n_{\ell-1}} \dots a^{n_2} b^{n_1}$$

where the n_i are in $\mathbf{Z} \setminus \{0\}$, except maybe if n_1 or n_ℓ is zero, but conjugating w by a power of a , we only need to consider the case $n_1 n_\ell \neq 0$. Set $N = \max |n_i|$.

Let h be an element of $\text{Diff}(\mathbf{k}, 0)$, and set $f_h = h^{-1} \circ f \circ h$. Let $r > 0$ be smaller than the convergence radius of h , f , g and their inverses. Choose $R > 0$ such that all these germs, and all their compositions of length $\leq 3N\ell$ map \mathbb{D}_R inside \mathbb{D}_r . If z is a point in \mathbb{D}_R , then its orbit under the action of f_h and g stays in \mathbb{D}_r for all compositions of these germs given by words of length $\leq N\ell$ in \mathbf{F}_2 ; in this situation, we say that the orbit of z is *well defined* up to length $N\ell$. In particular, if we look at the composition $w(f_h, g)$, and pick a point z in \mathbb{D}_R , we get a sequence of points

$$(9.2) \quad z_0 = z, z_1 = g^{n_1}(z_0), z_2 = f_h^{n_2}(z_1), \dots, z_\ell = w(f_h, g)(z_0).$$

To prove the theorem, we construct a triple (h, R, z) such that the orbit of z is well defined and the z_i are pairwise distinct; in particular, $z_\ell \neq z_0$ and $w(f_h, g) \neq \text{Id}$.

Proof. We do a recursion on the length ℓ , proving the existence of a triple (h, R, z) such that the z_i are pairwise distinct for $0 \leq i \leq \ell$. Since f and g have infinite order, the union of all fixed points of f^m and g^m in \mathbb{D}_r for $-N \leq m \leq N$ is a finite set F . For $j = 1$, we just pick a point z_0 sufficiently near the origin with $z_1 := g^{n_1}(z_0) \neq z_0$; the only constraint is to take z_0 in the complement of F . The points z_0 and z_1 will be kept fixed in the recursion.

Assume that a polynomial germ of diffeomorphism h_k has been constructed, in such a way that (a) the points $z_0, z_1, z_2, \dots, z_{2k}$, and z_{2k+1} are pairwise distinct (we just initialized the recursion for $k = 0$), and (b) $h_k(z) = z + \epsilon_k R_k(z)$ for some small $\epsilon_k \in \mathbf{k}$ and some element $R_k \in \mathbf{k}[z]$ of degree $\leq (2k)!$ which is divisible by z^2 . Consider a polynomial germ

$$(9.3) \quad P_k(z) = z + \eta_k z^2 \prod_{j=0}^{2k} (z - z_j)$$

with a small $\eta_k \in \mathbf{k}$; then

- P_k fixes z_j for all $j \leq 2k$,
- $P_k(z_{2k+1}) = a_k \eta_k + b_k$ for some pair $(a_k, b_k) \in \mathbf{k}^2$ with $a_k \neq 0$,
- as η_k goes to 0, the radius of convergence of P_k and its inverse P_k^{-1} go to infinity.

If we compose h_k with P_k then $H = h_k \circ P_k$ is a new polynomial germ such that the orbit of z_0 under f_H and g gives the same sequence z_0, z_1, \dots , up to z_{2k+1} . The next point is

$$(9.4) \quad z_{2k+2} = f_H^{n_{2k+2}}(z_{2k+1})$$

and we want to exclude the possibility $z_{2k+2} \in \{z_0, \dots, z_{2k+1}\}$; since

$$(9.5) \quad z_{2k+2} = (P_k^{-1} \circ f_{h_k}^{n_{2k+2}} \circ P_k)(z_{2k+1})$$

we want to avoid the inclusion

$$(9.6) \quad f_{h_k}^{n_{2k+2}}(P_k(z_{2k+1})) \subset P_k\{z_0, \dots, z_{2k+1}\},$$

and for that we just need to choose the parameter η_k in the definition of P_k in such a way that $f_{h_k}^{n_{2k+2}}(P_k(z_{2k+1}))$ is not in $\{z_0, \dots, z_{2k}\}$ and $P_k(z_{2k+1})$ is not a fixed point of $f_{h_k}^{n_{2k+2}}$. These constraints are satisfied for all small non-zero values of η_k because $f_{h_k}^{n_{2k+2}}$ is not the identity and the coefficient a_k in $P_k(z_{2k+1}) = a_k \eta_k + b_k$ is not zero.

The next point is $z_{2k+3} = g^{n_{2k+3}}(z_{2k+2})$ and we want it to be disjoint from $\{z_0, \dots, z_{2k+1}, z_{2k+2}\}$. For this, we do a second perturbation of the conjugacy. Let

$$(9.7) \quad Q_k(z) = z + \beta_k z^2 \prod_{j=0}^{2k+1} (z - z_j)$$

with a small $\beta_k \in \mathbf{k}$; then

- Q_k fixes z_j for all $j \leq 2k + 1$,
- $Q_k(z_{2k+2}) = c_k \beta_k + d_k$ for some pair $(c_k, d_k) \in \mathbf{k}^2$ with $c_k \neq 0$,
- as β_k goes to 0, the radius of convergence of Q_k and its inverse Q_k^{-1} go to infinity.

Now, we set $h_{k+1} = Q_k \circ H$. This does not change the sequence z_i for $0 \leq i \leq 2k + 1$, but the last point z_{2k+2} is replaced by $c_k \beta_k + d_k$. Since $g^{n_{2k+3}} \neq \text{Id}$ and $c_k \neq 0$ any non-zero, small enough value of β_k assures that $z_{2k+3} \notin \{z_0, \dots, z_{2k+1}, z_{2k+2}\}$.

To sum up, if we set $h_{k+1} = Q_k \circ P_k \circ h_k$ then the sequence z_0, \dots, z_{2k+3} is now made of pairwise distinct points. Moreover, when the parameters η_k and β_k go to zero, the germ $Q_k \circ P_k$ and its inverse converge uniformly to the identity on the disk \mathbb{D}_{2R} , so we can assume that the orbit of z_0 is well defined for all composition of h_{k+1} , f , g , and their inverses of length $\leq 3N\ell$. The germ $Q_k \circ P_k$ is equal to $z + S_k(z)$ where S_k is divisible by z^2 and $\deg(S_k) \leq (2k + 1) \times (2k + 2)$. Thus,

$$(9.8) \quad h_{k+1}(z) = z + P_{k+1}(z)$$

where z^2 divides P_{k+1} and

$$(9.9) \quad \deg(P_{k+1}) \leq \deg(P_k) \times (2k + 1) \times (2k + 2) \leq (2k + 2)!$$

This proves the recursion and finishes the proof of the theorem. \square

Theorem 9.2. *Let $(\mathbf{k}, |\cdot|)$ be a complete, non-discrete valued field. If f and g are elements of $\text{Diff}(\mathbf{k}; 0)$ of infinite order, there exists an element h of $\text{Diff}(\mathbf{k}; 0)$ such that $f_h := h \circ f \circ h^{-1}$ and g generate a free group of rank 2. One can choose h such that $h'(0) = 1$.*

Note that Theorem 3.4 is a direct corollary of that result; one just need to start with $f = \lambda_1 z$ or $\lambda_1 z + z^2$ if λ_1 is a root of unity, and similarly for g .

Proof. Denote by a_n and b_n the coefficients of f and g respectively. Let $L \subset \mathbf{k}$ be the field generated by the a_n and b_n . Since \mathbf{k} is not discrete, it has no isolated point; being complete with no isolated point, it is uncountable (a simple consequence of Baire's theorem [Oxt]), and it follows that its transcendental degree over L is infinite: it contains an infinite sequence (c_i) of algebraically independent numbers (over the prime field of \mathbf{k} , see [Lan, Chapter VIII]). We can moreover assume that all c_i are in the unit disk. Set $h_0(z) = \sum_{n \geq 1} c_n z^n$.

Consider a non-trivial element w of \mathbb{F}_2 . The N -th coefficient function

$$(9.10) \quad h \mapsto A_N(w(h \circ f \circ h^{-1}, g))$$

is a polynomial function on $\widehat{\text{Diff}}(\mathbf{k}, 0)$ in the sense of Section 2.1; this means that it is a polynomial function in the coefficients of h and $A_1(h)^{-1}$ (here, f and g are fixed). If $A_N(w(h_0 \circ f \circ h_0^{-1}, g))$ vanishes (resp. is equal to 1), then $A_N(w(h \circ f \circ h^{-1}, g)) = 0$ (resp. 1) for all formal diffeomorphisms h , because the c_i are algebraically independent over \mathbf{k} . Thus, Theorem 9.1 implies that $w(h_0 \circ f \circ h_0^{-1}, g) \neq \text{Id}$, and this shows that $f_{h_0} := h_0 \circ f \circ h_0^{-1}$ and g generate a free group of rank 2.

In this argument, we could start with $h_0 = z + \sum_{n \geq 2} c_n z^n$, because we can choose the germ h in Theorem 9.1 with the additional constraint $h'(0) = 1$. \square

Acknowledgements. Thanks to Yulij Ilyashenko, Frank Loray and Ludovic Marquis for several discussions on this subject, and to the participants of the seminars of Dijon, Moscou, Paris, and Toulouse during which the results of this paper were presented. We thank the referee for interesting comments and references, and for his suggestion leading to the last paragraph of Section 8.3.1. This work was conducted within the framework of the Henri Lebesgue Center ANR-11-LABX-0020-01. The first and second authors thank the del Duca Foundation, from the French Academy of Sciences, for its support.

References

- [Bau] G. BAUMSLAG, On generalised free products. *Math. Z.* **78** (1962), 423–438. Zbl 0104.24402 MR 0140562
- [BCLN] M. BERTHIER, D. CERVEAU and A. LINS NETO, Sur les feuilletages analytiques réels et le problème du centre. *J. Differential Equations* **131** (1996), 244–266. Zbl 0886.58090 MR 1419014
- [BGSS] E. BREUILLARD, T. GELANDER, J. SOUTO and P. STORM, Dense embeddings of surface groups. *Geom. Topol.* **10** (2006), 1373–1389. Zbl 1132.22011 MR 2255501
- [Bru1] A. BRUDNYI, Some algebraic aspects of the center problem for ordinary differential equations. *Qual. Theory Dyn. Syst.* **9** (2010), 9–28. Zbl 1202.37111 MR 2737360
- [Bru2] ——— Subgroups of the group of formal power series with the big powers condition. Preprint arXiv:1908.04918 (2019).
- [BT] H. W. BROER and F. M. TANGERMAN, From a differentiable to a real analytic perturbation theory, applications to the Kupka Smale theorems. *Ergodic Theory Dynam. Systems* **6** (1986), 345–362. Zbl 0582.58019 MR 0863199
- [Cam] R. CAMINA, The Nottingham group. In *New horizons in pro- p groups*, vol. 184 of *Progr. Math.* Birkhäuser Boston, Boston, MA (2000), pp. 205–221. Zbl 0977.20020 MR 1765121
- [Cer] D. CERVEAU, Quelques problèmes en géométrie feuilletée pour les 60 années de l'IMPA. *Bull. Braz. Math. Soc. (N.S.)* **44** (2013), 653–679. Zbl 1333.37061 MR 3167127
- [CG] C. CHAMPETIER and V. GUIARDEL, Limit groups as limits of free groups. *Israel J. Math.* **146** (2005), 1–75. Zbl 1103.20026 MR 2151593
- [EBN] H. EYNARD-BONTEMPS and A. NAVAS, Mather invariant, distortion, and conjugates for diffeomorphisms of the interval. Preprint, arXiv:1912.09305 (2019), 1–59.
- [Gla] A. M. W. GLASS, The ubiquity of free groups. *Math. Intelligencer* **14** (1992), 54–57. Zbl 0791.20001 MR 1184319
- [HY] M. HERMAN and J.-C. YOCOZ, Generalizations of some theorems of small divisors to non-Archimedean fields. In *Geometric dynamics (Rio de Janeiro, 1981)*, vol. 1007 of *Lecture Notes in Math.* Springer, Berlin (1983), pp. 408–447. Zbl 0528.58031 MR 0730280
- [IP] Y. S. ILYASHENKO and A. S. PYARTLI, The monodromy group at infinity of a generic polynomial vector field on the complex projective plane. *Russian J. Math. Phys.* **2** (1994), 275–315. Zbl 0915.34005 MR 1330871
- [Jen] S. A. JENNINGS, Substitution groups of formal power series. *Canadian J. Math.* **6** (1954), 325–340. Zbl 0058.02201 MR 0061610
- [Koe] G. KOENIGS, Nouvelles recherches sur les équations fonctionnelles. *Ann. Sci. École Norm. Sup. (3)* **2** (1885), 385–404. Zbl 17.0370.02 MR 1508770

- [Lan] S. LANG, *Algebra*, vol. 211 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, third ed. (2002). Zbl 0984.00001 MR 1878556
- [Les] J. LESLIE, On the group of real analytic diffeomorphisms of a compact real analytic manifold. *Trans. Amer. Math. Soc.* **274** (1982), 651–669. Zbl 0513.58017 MR 0675073
- [Lyn] R. C. LYNDON, The equation $a^2b^2 = c^2$ in free groups. *Michigan Math. J.* **6** (1959), 89–95. <http://projecteuclid.org/euclid.mmj/1028998143> Zbl 0084.02803 MR 0103218
- [Mil] J. MILNOR, *Dynamics in One Complex Variable*. Vol. 160 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, third ed. (2006). Zbl 1085.30002 MR 2193309
- [MRR] J.-F. MATTEI, J. C. REBELO and H. REIS, Generic pseudogroups on $(\mathbb{C}, 0)$ and the topology of leaves. *Compos. Math.* **149** (2013), 1401–1430. Zbl 1314.32033 MR 3103071
- [MS] L. MARQUIS and J. SOUTO, Surface groups of diffeomorphisms of the interval. *Israel J. Math.* **227** (2018), 379–396. Zbl 06976582 MR 3846328
- [Ols] A. Y. OL'SHANSKIĬ, On residualizing homomorphisms and G -subgroups of hyperbolic groups. *Internat. J. Algebra Comput.* **3** (1993), 365–409. MR 1250244
- [Oxt] J. C. OXTOBY, *Measure and Category*. Vol. 2 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, Berlin, second ed. (1980). A survey of the analogies between topological and measure spaces. Zbl 0435.28011 MR 0584443
- [Sie] C. L. SIEGEL, Iteration of analytic functions. *Ann. of Math. (2)* **43** (1942), 607–612. Zbl 0061.14904 MR 0007044
- [Ste] S. STERNBERG, On the structure of local homeomorphisms of euclidean n -space. *Amer. J. Math.* **80** (1958), 623–631. Zbl 0083.31406 MR 0096854
- [Sze] B. SZEGEDY, Almost all finitely generated subgroups of the Nottingham group are free. *Bull. London Math. Soc.* **37** (2005), 75–79. Zbl 1073.20022 MR 2105821
- [Tak] F. TAKENS, Normal forms for certain singularities of vector fields. *Ann. Inst. Fourier* **23** (1973), 163–195. Zbl 0266.34046 MR 0365620
- [Wal] M. WALDSCHMIDT, Indépendance algébrique des nombres de liouville. *Lecture Notes in Math.* **1415** (1990), 225–235. Zbl 0729.11035 MR 1044117
- [Whi] S. WHITE, The group generated by $x \mapsto x + 1$ and $x \mapsto x^p$ is free. *J. Algebra* **118** (1988), 408–422. Zbl 0662.20024 MR 0969681
- [Yoc] J.-C. YOCOZ, Centralisateurs et conjugaison différentiable des difféomorphismes du cercle. 231 (1995). pp. 89–242. Petits diviseurs en dimension 1. MR 1367354

(Reçu le 19 septembre 2019)

Serge CANTAT, Univ Rennes, CNRS,
IRMAR – UMR 6625, F-35000 Rennes, France
e-mail: serge.cantat@univ-rennes1.fr

Dominique CERVEAU, Univ Rennes, CNRS,
IRMAR – UMR 6625, F-35000 Rennes, France
e-mail: dominique.cerveau@univ-rennes1.fr

Vincent GUIARDEL, Univ Rennes, CNRS,
IRMAR – UMR 6625, F-35000 Rennes, France
e-mail: vincent.guirardel@univ-rennes1.fr

Juan SOUTO, Univ Rennes, CNRS,
IRMAR – UMR 6625, F-35000 Rennes, France
e-mail: juan.souto@univ-rennes1.fr