

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 8 (1962)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SURVEY OF COBORDISM THEORY
Autor: Milnor, J.
DOI: <https://doi.org/10.5169/seals-37949>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A SURVEY OF COBORDISM THEORY¹

by J. MILNOR

This paper will start out with a discussion of known results and then will taper off into a discussion of unsolved problems.

The theory of cobordism was initiated by L. Pontrjagin and V. A. Rohlin [10, 12]. It came of age with the work of R. Thom [17]. The basic question in this theory is the following. Let \mathcal{M} be some class of compact manifolds. Given $V \in \mathcal{M}$ how can one decide whether or not V is the boundary of some other manifold in \mathcal{M} ? Of course a necessary condition is that V itself must be a *closed* manifold: that is the boundary ∂V must be *vacuous*.

1. THE CLASSICAL COBORDISM GROUPS N_k AND Ω_k .

As a first illustration of this problem let \mathcal{D} denote the class of all compact differentiable manifolds. The manifolds $V \in \mathcal{D}$ need not be connected or orientable, and are allowed to have boundaries.

THEOREM 1 (Pontrjagin, Thom). — *A closed k -dimensional manifold $V \in \mathcal{D}$ is the boundary of some $(k+1)$ -dimensional manifold in \mathcal{D} if and only if the Stiefel-Whitney numbers $\omega_{i_1} \dots \omega_{i_n} [V]$ are all zero.*

(Explanation: The Stiefel-Whitney cohomology classes²⁾ $\omega_i \in H^i(V; J_2)$ are defined for example in Steenrød [15]. If $i_1 + \dots + i_n = k$ is any partition of k then the cup product $\omega_{i_1} \dots \omega_{i_n}$ is a top dimensional cohomology class. Applying the canonical "integration" homomorphism

$$[V]: H^k(V; J_2) \rightarrow J_2$$

we obtain a "Stiefel-Whitney number" $\omega_{i_1} \dots \omega_{i_n} [V] \in J_2$.)

¹⁾ Talk delivered at the Zurich Colloquium on Differential Geometry and Topology, June 1960.

²⁾ The notation J will be used for the integers and J_2 for the integers modulo 2.

The *non-oriented cobordism group* $N_k = H_k(\mathcal{D})$ is constructed as follows. Given two k -manifolds $V, V' \in \mathcal{D}$ the *sum* $V + V'$ will mean the (disjoint) topological sum, provided with a differentiable structure in the obvious way.

Definition. Two closed manifolds $V, V' \in \mathcal{D}$ are *congruent modulo* \mathcal{D} if $V + V'$ is the boundary of some manifold in \mathcal{D} . The set of all congruence classes of closed k -manifolds, under the composition operation $+$, forms the required group N_k . We will also use the notation $H_k(\mathcal{D})$ for this group since it is something like a homology group. (The Russian term for “cobordism” is “intrinsic homology”.)

It follows from Theorem 1 that each N_k is a finite abelian group of the form $J_2 \oplus \dots \oplus J_2$.

The cartesian product operation between differentiable manifolds gives rise to a bilinear pairing

$$N_k \oplus N_l \rightarrow N_{k+l}.$$

Thus the graded group $N_* = (N_0, N_1, \dots)$ has the structure of a graded ring.

THEOREM 2 (Thom). — *The non-oriented cobordism ring N_* has the structure of a polynomial algebra*

$$J_2 [X_2, X_4, X_5, X_6, X_8, X_9, \dots]$$

with one generator $X_k \in N_k$ for each dimension which is not of the form $2^m - 1$.

If k is even then the real projective k -space can be taken as generator. For k odd generators have been constructed by Dold [4].

Thom's proof of Theorems 1 and 2 involves a brilliant mixture of algebra and geometry. A key step in the argument is his proof that N_k is isomorphic to a certain homotopy group. I will not try to give details.

Next consider the class \mathcal{D}_o consisting of all *oriented* compact differentiable manifolds.

THEOREM 1'. — *A closed manifold in \mathcal{D}_o is the boundary of a manifold in \mathcal{D}_o if and only if both its Stiefel-Whitney numbers and its Pontrjagin numbers are zero.*

This result is due to Pontrjagin, Thom, Milnor, Averbuh, and Wall. (See [2, 9, 19].) For the definition of the Pontrjagin numbers $p_{i_1} \dots p_{i_n} [V] \in J$ the reader is referred to Hirzebruch [6]. These numbers are defined only if the dimension k is a multiple of 4.

The *oriented cobordism ring* $\Omega_* = H_*(\mathcal{D}_o)$ is defined as follows. For $V \in \mathcal{D}_o$ let $-V$ denote the same manifold V with the opposite orientation. We will say that

$$V \equiv V' \pmod{\partial \mathcal{D}_o}$$

if $(-V) + V'$ is the boundary of some manifold in \mathcal{D}_o . As an example, for any closed manifold V we have $V \equiv V \pmod{\partial \mathcal{D}_o}$ since

$$(-V) + V \approx \partial(V \times I)$$

where I denotes the unit interval. The set of all such congruence classes form the required group Ω_k . Again the cartesian product operation makes $\Omega_* = (\Omega_0, \Omega_1, \dots)$ into a graded ring.

It follows from Theorem 1' that Ω_k is a finitely generated group of the form

$$J \oplus \dots \oplus J \oplus J_2 \oplus \dots \oplus J_2$$

where infinite cyclic summands can occur only if $k \equiv 0 \pmod{4}$.

THEOREM 2'. — *The ring Ω_* , modulo the ideal consisting of 2-torsion elements, is a polynomial ring $J[Y_4, Y_8, Y_{12}, \dots]$ with one generator in each dimension divisible by 4.*

The complex projective space of real dimension $4m$ can be taken as generator for $m = 1, 2, 3$. However a different generator is needed in dimension 16.

For a description of the 2-torsion in Ω_* the reader is referred to Wall's paper.

2. MANIFOLDS WITH X -STRUCTURE.

In this section we will define the concept of an " X -structure" on the tangent bundle of a differentiable manifold; and study the corresponding cobordism theory.

First recall Steenrod's definition of a tensor field [15, § 6.4 and § 9.1 with mild alterations]. Every differentiable k -manifold V can be made Riemannian and hence has a tangent bundle with structural group O_k . Let X be any topological space on which the group O_k acts. Then we can form the weakly associated bundle with base space V and fibre X . This may be called the "tensor bundle of type X " and its cross-sections are "tensor fields". As an example, if $k = 2m$, then O_{2m} acts on the coset space O_{2m}/U_m .

A cross-section of the corresponding bundle is called a *quasi*-(or almost) *complex structure* on V . (See [15, § 41.10].)

We will modify this definition as follows, so that it makes sense for all dimensions simultaneously. Let O denote the union of the orthogonal groups $O_1 \subset O_2 \subset O_3 \subset \dots$ in the fine topology. Then we require that this infinite orthogonal group O act on the space X . It follows that each O_k acts on X . Hence there is a tensor bundle of type X over any manifold $V \in \mathcal{D}$.

Definition: A homotopy class of cross-sections of the tensor bundle with fibre X over V is called an X -structure on V . A manifold $V \in \mathcal{D}$ together with an X -structure on V is called an X -manifold. We will still use the single symbol V to denote this pair.

Now if V is an X -manifold then ∂V is also. Given any closed X -manifold V one can define a second X -manifold $-V$ so that

$$\partial(V \times I) \approx V + (-V).$$

Thus one can define a cobordism group for the class of X -manifolds. The resulting group will be denoted by $N_k(X)$ and called the X -cobordism group. (Following Atiyah [1] this could also be called the k -th "bordism group" of the O -space X .)

Example 1. Let O/U denote the union of the spaces

$$O_2/U_1 \subset O_4/U_2 \subset O_6/U_3 \subset \dots$$

in the fine topology with O acting on O/U in the usual way. Then a manifold with an O/U -structure will be called a *weakly complex manifold*. (Compare Hirzebruch [7].) For example

any complex manifold is quasi-complex and hence weakly complex. Any sphere can be given an O/U -structure although only S^2 and S^6 possess quasi-complex structures.

The following results are due to Milnor and Novikov.

THEOREM 1''. — *A closed weakly complex manifold V is the boundary of a weakly complex manifold if and only if its Chern numbers $c_{i_1} \dots c_{i_n}[V]$ are all zero.*

(Explanation: an O/U -structure on V determines a preferred U -bundle over V . Hence Chern classes are defined.) It follows that $N_k(O/U)$ is zero for k odd and is free abelian for k even.

THEOREM 2''. — *The graded group $N_*(O/U)$ has a natural ring structure, making it into a polynomial ring $J[Y_2, Y_4, Y_6, \dots]$ with one generator in each even dimension.*

As generators one can take certain algebraic varieties with their natural complex structures. (Compare [7]. It is not known whether connected varieties will suffice.)

Example 2. More generally one could use any subgroup G of the infinite orthogonal group in place of U . For example using the infinite symplectic group Sp we would obtain a cobordism ring $N_*(O/Sp)$ which is appropriate for the study of "weakly quaternionic manifolds". The following six groups seem particularly interesting:

$$1 \subset Sp \subset SU \subset U \subset SO \subset 0.$$

Starting from the right, the ring $N_*(O/O)$ is just the non-oriented cobordism ring N_* and $N_*(O/SO)$ is the oriented cobordism ring Ω_* . The rings $N_*(O/SU)$ and $N_*(O/Sp)$ are more or less unknown. (Compare the concluding remarks in [9].)

The ring $N_*(O/1) = N_*(O)$ has essentially been studied by Pontrjagin [11]. An O -structure on V is a trivialization of the tangent O -bundle of V (the "stable" tangent bundle). Manifolds which admit such a structure are called " π -manifolds". It turns out that $N_k(O)$ is isomorphic to the stable homotopy groups $\pi_{k+n}(S^n)$ of the n -sphere, with n large. This fact is the basis for Pontrjagin's method of studying homotopy groups.

Example 3. Let X be a space on which O operates trivially. Then an X -structure on V is just a preferred homotopy class of maps $V \rightarrow X$. As cases of particular interest X might be an Eilenberg-MacLane space or the classifying space of a group. How does one compute the groups $N_k(X)$?

The above definitions can be modified slightly by admitting only oriented manifolds. Thus one obtains groups $\Omega_k(X)$ where X is any space on which the rotation group SO acts. Again I do not know how to compute these groups. (Added in proof: See Conner and Floyd [21].)

Example 4. Let P denote the infinite real projective space, with the infinite rotation group SO acting in the natural way. The cobordism groups $\Omega_k(P)$ for oriented manifolds with P -structure can be called the *spinor cobordism groups*. This name is appropriate since a P -structure is roughly a "lifting" of the structural group of the tangent bundle to the infinite spinor group. A manifold admits a P -structure if and only if its Stiefel-Whitney class w_2 is zero. The groups $\Omega_k(P)$ have no odd torsion, but otherwise I do not know much about them.

3. MISCELLANEOUS COBORDISM THEORIES.

So far we have concentrated on differentiable manifolds. However one could equally well define a cobordism group based on the class \mathcal{T} of all compact topological manifolds. (Compare Brown [3, Theorem 3].) The natural correspondence $\mathcal{D} \rightarrow \mathcal{T}$ induces a homomorphism from the differentiable cobordism group $N_k = H_k(\mathcal{D})$ to the topological cobordism group $H_k(\mathcal{T})$.

Since Thom [16] has shown that Stiefel-Whitney classes can be defined topologically, we have:

THEOREM 3 (Thom). — *The homomorphism $N_k \rightarrow H_k(\mathcal{T})$ has kernel zero.*

Problem: Is this homomorphism onto?

Another possibility would be to consider the class \mathcal{C}_o of all compact, oriented, combinatorial manifolds. Whitehead [20] has shown that each differentiable manifold has a preferred class of triangulations. Hence there is a natural homomorphism from

$\Omega_k = H_k(\mathcal{D}_o)$ to $H_k(\mathcal{C}_o)$. Thom, Rohlin and Švarč have shown that Pontrjagin classes can be defined for combinatorial manifolds. Therefore we have:

THEOREM 3'. — *The homomorphism $\Omega_k \rightarrow H_k(\mathcal{C}_o)$ has kernel zero.*

However examples show that this homomorphism is not onto. The reader is referred to [13, 18].

Another interesting possibility would be to look at the class of compact homology manifolds.

Returning to the differentiable case, interesting cobordism groups can be obtained by restricting the connectivities of the manifolds involved. As an extreme case we can consider only differentiable manifolds which are either homotopy spheres or homotopy cells. The resulting cobordism groups are closely related to the problem of classifying differentiable structures on spheres. The reader is referred to Milnor [8] and Smale [14].

As a final, quite different, example consider differentiable imbeddings of the circle S^1 in the 3-sphere S^3 . Such an object (a knot) is said to *bound* if it can be extended to a differentiable imbedding of the disk D^2 in the disk D^4 . The resulting cobordism group has been studied by Fox and Milnor [5]. This group is not finitely generated.

REFERENCES

- [1] ATIYAH, M., Bordism and cobordism, Proc, Cambridge Phil. Soc., 57 (1961), 200-208.
- [2] AVERBUH, B. G., Algebraic structure of internal homology groups. Doklady Akad. Nauk S.S.S.R., 125 (1959), 11-14.
- [3] BROWN, M., Locally flat embeddings of topological manifolds. *A.M.S. Notices*, 7 (1960), 939-940.
- [4] DOLD, A., Erzeugende der Thomschen Algebra *N. Math. Zeits.*, 65 (1956), 25-35.
- [5] FOX, R. H. and J. MILNOR, Singularities of 2-spheres in 4-space and equivalence of knots. *Bull. Amer. Math. Soc.*, 63 (1957), 406.
- [6] HIRZEBRUCH, F., *Neue topologische Methoden in der algebraischen Geometrie*. Springer (Berlin), 1956.
- [7] ——— Komplexe Mannigfaltigkeiten. *Proceedings Int. Congr. Math.*, 1958 (1960), 119-136.

- [8] MILNOR, J., Sommes de variétés différentiables et structures différentiables des sphères. *Bull. Soc. Math. France*, 87 (1959), 439-444.
- [9] ——— On the cobordism ring Ω^* and a complex analogue, I. *American Journ. of Math.*, 82 (1960), 505-521. [See also *A.M.S. Notices*, 5 (1958), 457.]
- [10] PONTRJAGIN, L. S., Characteristic cycles on differentiable manifolds. *Mat. Sbornik*, 21 (1947), 233-284.
- [11] ——— Smooth manifolds and their applications in homotopy theory. *A.M.S. Translations*, 11 (1959), 1-114.
- [12] ROHLIN, V. A., A 3-dimensional manifold is the boundary of a 4-dimensional manifold. (Russian.) *Doklady Akad. Nauk S.S.S.R.*, 81 (1951), 355. [See also *Doklady*, v. 84, p. 221,, v. 89, p. 789 and v. 119, p. 876.]
- [13] ——— and A. S. ŠVARČ, Combinatorial invariance of the Pontrjagin classes. (Russian.) *Doklady Akad. Nauk S.S.S.R.*, 114 (1957), 490-493.
- [14] SMALE, S., Generalized Poincaré's conjecture in dimensions greater than four. *Annals of Math.*, 74 (1961), 391-406.
- [15] STEENROD, N., *The topology of fibre bundles*. Princeton, 1951.
- [16] THOM, R., Espaces fibrés en sphères et carrés de Steenrod. *Ann. Ecole Norm. Sup.*, 69 (1952), 109-181.
- [17] ——— Quelques propriétés globales des variétés différentiables. *Comment. Math. Helv.*, 28 (1954), 17-86.
- [18] ——— Les classes caractéristiques de Pontrjagin des variétés triangulées. *Symposium Internacional de Topologia Algebraica, Mexico* (1958), pp. 259-272.
- [19] WALL, C. T. C., Determination of the cobordism ring. *Annals of Math.*, 72 (1960), 292-311.
- [20] WHITEHEAD, J. H. C., On C^1 -complexes. *Annals of Math.*, 41 (1940), 809-824.
- [21] CONNER, P. E. and E. E. FLOYD, Differentiable periodic maps, Technical Note 13, Dept. of Math., Univ. of Virginia, 1961.
- [22] NOVIKOV, S. P., Some problems in the topology of manifolds connected with the theory of Thom spaces. *Dokl. Akad. Nauk SSSR* 132 (1960), 1031-1034 (Russian); translated as *Soviet Math. Dokl.* 1, 717-720.

Princeton University.