

2. Manifolds with X-structure.

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **8 (1962)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **20.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

This result is due to Pontrjagin, Thom, Milnor, Averbuh, and Wall. (See [2, 9, 19].) For the definition of the Pontrjagin numbers $p_{i_1} \dots p_{i_n} [V] \in J$ the reader is referred to Hirzebruch [6]. These numbers are defined only if the dimension k is a multiple of 4.

The *oriented cobordism ring* $\Omega_* = H_*(\mathcal{D}_o)$ is defined as follows. For $V \in \mathcal{D}_o$ let $-V$ denote the same manifold V with the opposite orientation. We will say that

$$V \equiv V' \pmod{\partial \mathcal{D}_o}$$

if $(-V) + V'$ is the boundary of some manifold in \mathcal{D}_o . As an example, for any closed manifold V we have $V \equiv V \pmod{\partial \mathcal{D}_o}$ since

$$(-V) + V \approx \partial(V \times I)$$

where I denotes the unit interval. The set of all such congruence classes form the required group Ω_k . Again the cartesian product operation makes $\Omega_* = (\Omega_0, \Omega_1, \dots)$ into a graded ring.

It follows from Theorem 1' that Ω_k is a finitely generated group of the form

$$J \oplus \dots \oplus J \oplus J_2 \oplus \dots \oplus J_2$$

where infinite cyclic summands can occur only if $k \equiv 0 \pmod{4}$.

THEOREM 2'. — *The ring Ω_* , modulo the ideal consisting of 2-torsion elements, is a polynomial ring $J[Y_4, Y_8, Y_{12}, \dots]$ with one generator in each dimension divisible by 4.*

The complex projective space of real dimension $4m$ can be taken as generator for $m = 1, 2, 3$. However a different generator is needed in dimension 16.

For a description of the 2-torsion in Ω_* the reader is referred to Wall's paper.

2. MANIFOLDS WITH X-STRUCTURE.

In this section we will define the concept of an "X-structure" on the tangent bundle of a differentiable manifold; and study the corresponding cobordism theory.

First recall Steenrod's definition of a tensor field [15, § 6.4 and § 9.1 with mild alterations]. Every differentiable k -manifold V can be made Riemannian and hence has a tangent bundle with structural group O_k . Let X be any topological space on which the group O_k acts. Then we can form the weakly associated bundle with base space V and fibre X . This may be called the "tensor bundle of type X " and its cross-sections are "tensor fields". As an example, if $k = 2m$, then O_{2m} acts on the coset space O_{2m}/U_m .

A cross-section of the corresponding bundle is called a *quasi-*(or almost) *complex structure* on V . (See [15, § 41.10].)

We will modify this definition as follows, so that it makes sense for all dimensions simultaneously. Let O denote the union of the orthogonal groups $O_1 \subset O_2 \subset O_3 \subset \dots$ in the fine topology. Then we require that this infinite orthogonal group O act on the space X . It follows that each O_k acts on X . Hence there is a tensor bundle of type X over any manifold $V \in \mathcal{D}$.

Definition: A homotopy class of cross-sections of the tensor bundle with fibre X over V is called an X -structure on V . A manifold $V \in \mathcal{D}$ together with an X -structure on V is called an X -manifold. We will still use the single symbol V to denote this pair.

Now if V is an X -manifold then ∂V is also. Given any closed X -manifold V one can define a second X -manifold $-V$ so that

$$\partial(V \times I) \approx V + (-V).$$

Thus one can define a cobordism group for the class of X -manifolds. The resulting group will be denoted by $N_k(X)$ and called the X -cobordism group. (Following Atiyah [1] this could also be called the k -th "bordism group" of the O -space X .)

Example 1. Let O/U denote the union of the spaces

$$O_2/U_1 \subset O_4/U_2 \subset O_6/U_3 \subset \dots$$

in the fine topology with O acting on O/U in the usual way. Then a manifold with an O/U -structure will be called a *weakly complex manifold*. (Compare Hirzebruch [7].) For example

any complex manifold is quasi-complex and hence weakly complex. Any sphere can be given an O/U -structure although only S^2 and S^6 possess quasi-complex structures.

The following results are due to Milnor and Novikov.

THEOREM 1''. — *A closed weakly complex manifold V is the boundary of a weakly complex manifold if and only if its Chern numbers $c_{i_1} \dots c_{i_n}[V]$ are all zero.*

(Explanation: an O/U -structure on V determines a preferred U -bundle over V . Hence Chern classes are defined.) It follows that $N_k(O/U)$ is zero for k odd and is free abelian for k even.

THEOREM 2''. — *The graded group $N_*(O/U)$ has a natural ring structure, making it into a polynomial ring $J[Y_2, Y_4, Y_6, \dots]$ with one generator in each even dimension.*

As generators one can take certain algebraic varieties with their natural complex structures. (Compare [7]. It is not known whether connected varieties will suffice.)

Example 2. More generally one could use any subgroup G of the infinite orthogonal group in place of U . For example using the infinite symplectic group Sp we would obtain a cobordism ring $N_*(O/Sp)$ which is appropriate for the study of "weakly quaternionic manifolds". The following six groups seem particularly interesting:

$$1 \subset Sp \subset SU \subset U \subset SO \subset 0.$$

Starting from the right, the ring $N_*(O/O)$ is just the non-oriented cobordism ring N_* and $N_*(O/SO)$ is the oriented cobordism ring Ω_* . The rings $N_*(O/SU)$ and $N_*(O/Sp)$ are more or less unknown. (Compare the concluding remarks in [9].)

The ring $N_*(O/1) = N_*(O)$ has essentially been studied by Pontrjagin [11]. An O -structure on V is a trivialization of the tangent O -bundle of V (the "stable" tangent bundle). Manifolds which admit such a structure are called " π -manifolds". It turns out that $N_k(O)$ is isomorphic to the stable homotopy groups $\pi_{k+n}(S^n)$ of the n -sphere, with n large. This fact is the basis for Pontrjagin's method of studying homotopy groups.

Example 3. Let X be a space on which O operates trivially. Then an X -structure on V is just a preferred homotopy class of maps $V \rightarrow X$. As cases of particular interest X might be an Eilenberg-MacLane space or the classifying space of a group. How does one compute the groups $N_k(X)$?

The above definitions can be modified slightly by admitting only oriented manifolds. Thus one obtains groups $\Omega_k(X)$ where X is any space on which the rotation group SO acts. Again I do not know how to compute these groups. (Added in proof: See Conner and Floyd [21].)

Example 4. Let P denote the infinite real projective space, with the infinite rotation group SO acting in the natural way. The cobordism groups $\Omega_k(P)$ for oriented manifolds with P -structure can be called the *spinor cobordism groups*. This name is appropriate since a P -structure is roughly a "lifting" of the structural group of the tangent bundle to the infinite spinor group. A manifold admits a P -structure if and only if its Stiefel-Whitney class ω_2 is zero. The groups $\Omega_k(P)$ have no odd torsion, but otherwise I do not know much about them.

3. MISCELLANEOUS COBORDISM THEORIES.

So far we have concentrated on differentiable manifolds. However one could equally well define a cobordism group based on the class \mathcal{T} of all compact topological manifolds. (Compare Brown [3, Theorem 3].) The natural correspondence $\mathcal{D} \rightarrow \mathcal{T}$ induces a homomorphism from the differentiable cobordism group $N_k = H_k(\mathcal{D})$ to the topological cobordism group $H_k(\mathcal{T})$.

Since Thom [16] has shown that Stiefel-Whitney classes can be defined topologically, we have:

THEOREM 3 (Thom). — *The homomorphism $N_k \rightarrow H_k(\mathcal{T})$ has kernel zero.*

Problem: Is this homomorphism onto?

Another possibility would be to consider the class \mathcal{C}_o of all compact, oriented, combinatorial manifolds. Whitehead [20] has shown that each differentiable manifold has a preferred class of triangulations. Hence there is a natural homomorphism from