

FLATNESS AND PRIVILEGE

Autor(en): **Douady, A.**

Objektyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **14 (1968)**

Heft 1: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **24.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-42343>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

FLATNESS AND PRIVILEGE

by A. DOUADY

I. FLAT MORPHISMS

§ 1. Analytic subspaces of an analytic space

Let Y_1 and Y_2 be closed analytic subspaces of an analytic space X , and let them be defined by the \mathcal{O}_X ideals J_1, J_2 .

Definition 1: We say that Y_1 is *analytically included* in Y_2 , and we write $Y_1 \subset Y_2$, when $J_1 \supset J_2$.

Remark: The analytic inclusion implies the set theoretic inclusion, but the converse is not true.

Example: $X = (\mathbf{C}, \mathcal{O}_{\mathbf{C}})$; $J_1 = (x)$, $J_2 = (x^2)$. The space Y_1 is a simple point, Y_2 is a double point, $Y_1 \not\subset Y_2$, while they have the same underlying set.

Definition 2: The subspace $Y_1 \cup Y_2$ is the smallest subspace of X containing Y_1 and Y_2 , and it is defined by $J_1 \cap J_2$. The subspace $Y_1 \cap Y_2$ is the biggest subspace of X contained in both Y_1 and Y_2 , and it is defined by $J_1 + J_2$.

Remark: The underlying set of $Y_1 \cup Y_2$ (Resp. $Y_1 \cap Y_2$) is the union (Resp. intersection) of the underlying sets of Y_1 and Y_2 . However \cup and \cap of analytic spaces do not satisfy the distributivity laws which hold in set-theory: $(Y_1 \cup Y_2) \cap Y_3$ contains $Y_1 \cap Y_3$ and $Y_2 \cap Y_3$, and therefore their union; similarly $(Y_1 \cap Y_2) \cup Y_3 \subset (Y_1 \cup Y_3) \cap (Y_2 \cup Y_3)$. In general the converse inclusions do not hold.

Example: Let $X = \mathbf{C}^2$ and Y_1, Y_2, Z be given by ideals $(x-y)$, $(x+y)$ and (x) respectively.

$(Y_1 \cup Y_2) \cap Z$ is $\{0\}$ provided with $\mathbf{C}\{y\}/(y^2)$, while $(Y_1 \cap Y_2) \cup (Y_2 \cap Z)$ is the reduced space $\{0\}$. On the other hand: $Y_1 \cap Y_2 \subset Z$, $(Y_1 \cap Y_2) \cup Z = Z$, while $(Y_1 \cup Z) \cap (Y_2 \cup Z)$ is the space defined by the ideal (x^2, xy) . Its local ring at the origin is $\mathbf{C}\{x, y\}/(x^2, xy)$ in which x is nilpotent.

Definition 3 : Let X', X be analytic spaces, Y a closed analytic subspace of X defined by J , and $f = (f_0, f^1) : X' \rightarrow X$ a morphism.

The inverse image of Y by $f, f^{-1}(Y)$, is the analytic subspace Y' of X' defined by the ideal $J' = f^1(J) \mathcal{O}_{X'}$.

The inverse image of a simple point x in X is called the f -fiber over x , and is denoted by $f^{-1}(x)$ or $X'(x)$.

Proposition 1 : If $f = (f_0, f^1) : X' \rightarrow X$ is a morphism of analytic spaces, and Y is a subspace of X , then $f^{-1}(Y) \simeq \underset{X}{Y \times X'}$.

Proof : Let T be any analytic space, and $g : T \rightarrow X'$ a morphism. Then g can be considered as a morphism from T to $f^{-1}(Y)$ if and only if $f \circ g$ can be considered as a morphism from T to Y . Thus $f^{-1}(Y)$ and $\underset{X}{X' \times X}$ are solutions of the same universal problem.

§ 2. Analytic pull-back

In the following we want to generalize the notion of inverse image of a subspace.

We shall first recall the basic properties of the tensor product $E \otimes_A F$, where A is a commutative ring and E, F are two A -modules.

(1°) $E \otimes A^n = E^n$ ($n \in \mathbb{N}$)

(2°) If the sequence of A -modules $F' \rightarrow F \rightarrow F'' \rightarrow 0$ is exact, then also the sequence $E \otimes F' \rightarrow E \otimes F \rightarrow E \otimes F'' \rightarrow 0$ is exact. (Right exactness of the tensor product)

(3°) If $(F_i)_{i \in I}; f_{ij} : F_j \rightarrow F_i$ is an inductive system, then

$$E \otimes \lim_{\rightarrow} F_i = \lim_{\rightarrow} (E \otimes F_i).$$

On the other hand these properties characterize completely the functor \otimes .

Definition 1 : Let $f = (f_0, f^1) : X' \rightarrow X$ be a morphism of analytic spaces, and \mathcal{E} an \mathcal{O}_X -module. Then $f_0^* \mathcal{E}$ is an $f_0^* \mathcal{O}_X$ -module and $\mathcal{O}_{X'}$ is also an $f_0^* \mathcal{O}_X$ -module (by $f^1 : f_0^* \mathcal{O}_X \rightarrow \mathcal{O}_{X'}$).

The analytic pull-back $f^* \mathcal{E}$ of \mathcal{E} by f is defined by scalar extension:

$$f^* \mathcal{E} = f_0^* \mathcal{E} \otimes_{f_0^* \mathcal{O}_X} \mathcal{O}_{X'}$$

Remark : The inverse image is a particular case of the analytic pull-back.

In fact, if Y is a closed analytic subspace of X and $f : X' \rightarrow X$ is a morphism:

$$f^* \mathcal{O}_Y = f_0^* (\mathcal{O}_X / J_Y) \otimes_{f_0^* \mathcal{O}_X} \mathcal{O}_{X'} \simeq f_0^* \mathcal{O}_X / f_0^* J_Y \otimes_{f_0^* \mathcal{O}_X} \mathcal{O}_{X'} \\ \simeq \mathcal{O}_{X'} / f^1(J_Y). \mathcal{O}_{X'} \simeq \mathcal{O}_{f^{-1}(Y)}$$

(The third isomorphism follows from the fact, that $A/I \otimes_A E \simeq E/IE$).

Elementary properties of the analytic pull-back :

- (a) $(f^* \mathcal{E})_{x'} = (f_0^* \mathcal{E})_{x'} \otimes_{(f_0^* \mathcal{O}_X)_{x'}} \mathcal{O}_{X',x'} \simeq \mathcal{E}_x \otimes_{\mathcal{O}_{X,x}} \mathcal{O}_{X',x'}$ where $x = f_0(x')$ (since \otimes commutes with inductive limits).
- (b) $f^* (\mathcal{E} \otimes_{\mathcal{O}_X} \mathcal{F}) = f^* \mathcal{E} \otimes_{\mathcal{O}_{X'}} f^* \mathcal{F}$, where \mathcal{E} and \mathcal{F} are \mathcal{O}_X -modules.
- (c) If \mathcal{E} is a coherent \mathcal{O}_X -module, then $f^* \mathcal{E}$ is a coherent $\mathcal{O}_{X'}$ -module.

In fact, \mathcal{E} has a locally finite presentation:

$$\mathcal{O}_X^q \rightarrow \mathcal{O}_X^p \rightarrow \mathcal{E} \rightarrow 0, \text{ and } f^* \text{ is compatible with cokernels, } f^* (\mathcal{O}_X^r) = \mathcal{O}_{X'}^r.$$

Special case : The pull-back of vector bundle. Let (E, π) be an analytic

$$\begin{array}{ccc} E \times X' & \xrightarrow{\bar{f}} & E \\ \downarrow \pi' & & \downarrow \pi \\ X' & \xrightarrow{f} & X \end{array}$$

vector bundle over the analytic space X , and $f : X' \rightarrow X$ a morphism of analytic spaces. The fiber product carries a unique structure of vector bundle over X' , such that \bar{f} is a bundle morphism. We call this bundle E' .

Proposition 1 : Let \mathcal{E} (Resp. \mathcal{E}') be the sheaf of analytic sections of E (Resp. E'). Then $\mathcal{E}' = f^* \mathcal{E}$.

Proof (Sketch) : We have a $f_0^* \mathcal{O}_X$ linear morphism $f_0^* \mathcal{E} \rightarrow \mathcal{E}'$, which extends to a morphism $f^* \mathcal{E} \rightarrow \mathcal{E}'$. We can prove that this is an isomorphism. Since the question is local with respect to X' , we can suppose that E is a trivial bundle over X with fiber \mathbf{C}^r , then $\mathcal{E} = \mathcal{O}_X^r$. Also $\mathcal{O}_{X'}^r = f^* \mathcal{O}_X^r$. Therefore $f^* \mathcal{E} = \mathcal{E}'$.

§ 3. Introduction to flatness by examples


Let S be an analytic space. By analytic space over s we mean an analytic space X provided with a morphism $\pi : X \rightarrow S$. Let S be a simple point in S , and consider $X(s) = f^{-1}(s)$.

The main purpose of these lectures is to give a precise meaning to the expression:

“ $X(s)$ depends nicely on s ”, and to give a criterion for the “ nice ” behaviour.

We begin with some examples.


Example 1: X is the closed subspace on \mathbf{C}^2 defined by $(y^2 - x)$, $S = \mathbf{C}$ and $\pi = 1st$ projection.



$$X(s) = \begin{cases} 2 \text{ simple points if } s \neq 0 \\ \text{double point if } s = 0. \end{cases}$$

Here the behaviour of $X(s)$ is nice.

Example 2: X is the closed subspace of \mathbf{C}^2 defined by (xy) , $S = \mathbf{C}$ and $\pi = 1st$ projection.



$X(s)$ is given by $(x-s, xy)$, and

$$(x-s, xy) = \begin{cases} (x-s, y) & \text{if } s \neq 0 \\ (x) & \text{if } s = 0. \end{cases}$$

The first case is a simple point, the second one the, y -axis.

A similar example is the map of a point into \mathbf{C} .

In both of these examples the dimension of the fiber makes a jump at one point. We notice, however, that the exceptional point corresponds to an irreducible component of X , and after removing this component π behaves nicely.

This kind of removing is not possible in general, as the following example shows:

Example 3: X is given in \mathbf{C}^3 by $(xz - y)$, and π is the projection on the (x, y) -plane.

If $s = (x_0, y_0)$, then the fiber $X(s)$ is defined by

$$(x-x_0, y-y_0, xz-y) = \begin{cases} \left(x-x_0, y-y_0, z-\frac{y_0}{x_0}\right) & \text{if } x_0 \neq 0 \\ (x, y) & \text{if } x_0 = y_0 = 0 \\ (1) & \text{if } x_0 = 0, y_0 \neq 0. \end{cases}$$

The set of “ nice ” fibers is dense in X , so we cannot remove the z -axis and still get a closed subspace of \mathbf{C}_3 .

§ 4. Algebraic study of flatness

In the following all rings are commutative, with 1, and all modules are unitary.

Definition 1: An A -module E is *flat*, if for every exact sequence of A -modules

$$0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0,$$

the sequence $0 \rightarrow E \otimes F' \rightarrow E \otimes F \rightarrow E \otimes F'' \rightarrow 0$ is also exact. We can also say, because \otimes is right exact, that E is flat, if for every injective homomorphism $F' \rightarrow F$, $E \otimes F' \rightarrow E \otimes F$ is also injective.

Examples of modules which are not flat :

- (1) if $A = \mathbf{Z}$, $E = \mathbf{Z}_2 = \mathbf{Z}/2\mathbf{Z}$, $F = F' = \mathbf{Z}$; then the sequence $0 \rightarrow \mathbf{Z} \xrightarrow{2I} \mathbf{Z} (2I : x' \rightarrow 2x)$ is exact. But now $\mathbf{Z}_2 \otimes \mathbf{Z} = \mathbf{Z}_2$, and the homomorphism $\mathbf{Z}_2 \xrightarrow{2I} \mathbf{Z}_2$ is the zero homomorphism, which is not injective. So \mathbf{Z}_2 is not a flat \mathbf{Z} module.
- (2) If $A = \mathbf{C}\{x\}$, $E = \mathbf{C} = \mathbf{C}\{x\}/(x)$, $F = F' = \mathbf{C}\{x\}$, then the sequence $0 \rightarrow F \xrightarrow{xI} F' (xI : p(x) \rightarrow xp(x))$ is exact. But the homomorphism $E \xrightarrow{xI} E$ is not injective.

Proposition 1: If A is an integral domain and E a flat A -module, then E is torsion-free.

Proof: Let $a \in A$, $a \neq 0$. Because A is an integral domain, the sequence $0 \rightarrow AA \xrightarrow{aI} (aI : x \rightarrow ax)$ is exact. Since E is flat, the sequence $0 \rightarrow E \xrightarrow{aI} E$ is also exact. In other words E has no torsion elements.

Proposition 2: If A is a principal-ideal domain, then E is flat if and only if E is torsionfree.

Proof: See corollary of prop. 6.

Examples of flat modules :

- (1) The inductive limit of flat modules is flat, because the inductive limit preserves exactness, and it commutes with the tensor product.

(2) Every free module is flat. In fact, if E is free and finite type, then $E = A^n$ and $E \otimes F = F^n$. If $F' \rightarrow F$ is injective, so is $F'^n \rightarrow F^n$ too.

If E is an arbitrary free module, then it is an inductive limit of free modules of finite type, and the flatness of E follows from (1).

(3) Let S be a multiplicative system in A . Then the ring of fractions $S^{-1}A$ is a flat A -module. In fact the ring $S^{-1}A$ can be identified with an inductive limit of free modules, so it is flat ((1) (2)). We assume for simplicity that S has only regular elements. We can define in the set S a partial order in the following way:

$$s' \geq s \Leftrightarrow \exists t \in A, \quad ts = s' \quad (\text{such a } t \text{ is then unique}).$$

Let $E_s = A$ for every $s \in S$, and if $s' \geq s$ (i.e. $s' = ts$) then let $f_s^{s'}$ be the homomorphism $t \cdot I_A : E_s \rightarrow E_{s'}$. The family $(E_s)_{s \in S}$ with the homomorphisms $(f_s^{s'})$ is an inductive system.

Let $E = \lim_{\rightarrow} E_s$ be the inductive limit of this system, and φ_s the canonical homomorphism $E_s \rightarrow E$. We shall define an isomorphism $\psi : E \rightarrow S^{-1}A$.

We first define for every s a homomorphism $\psi_s : E_s = A \rightarrow S^{-1}A$; $x \rightarrow x/s$. Now if $s' \geq s$, then

$$(\psi_{s'} \circ f_s^{s'})(x) = \psi_{s'}(tx) = \frac{tx}{s'} = \frac{tx}{ts} = \frac{x}{s} = \psi_s(x).$$

Therefore there exists a homomorphism $\psi : E \rightarrow S^{-1}A$, satisfying $\psi_s = \psi \circ \varphi_s$ for every $s \in S$.

Because every element of $S^{-1}A$ has the form a/s , ψ is surjective. On the other hand if $\psi(\varphi_s(x)) = 0$, then $\psi_s(x) = x/s = 0$. Thus $x = 0$, and ψ is also injective.

The above proof can be extended to the general case, not assuming that the elements of S are regular. The extended proof involves the notion of inductive limit of an inductive system indexed by a category instead of an ordered set.

From (1) and (2) above, any module which is the inductive limit of free modules, is flat. Conversely:

Theorem 1 : (Daniel, Lazard)

Any flat module is a inductive limit of free modules.

For the proof: See *C.R. Acad. Sci. Paris*, 258 (1964), pp. 6313-6316.

Some elementary properties of flat modules :

- (1) If E and F are flat A -modules, then $E \otimes_A F$ is also flat. In fact, if $G' \rightarrow G$ is injective, then $F \otimes_A G' \rightarrow F \otimes_A G$ is injective, and also $E \otimes_A (F \otimes_A G') \rightarrow E \otimes_A (F \otimes_A G)$ is injective. The result follows from the associativity of the tensor product.
- (2) Let $\phi : A \rightarrow B$ be a ring homomorphism, and E a flat A -module. The module $B \otimes_A E$ is a flat B -module.

If F is a B -module, then $F \otimes_B (B \otimes_A E) = (F \otimes_B B) \otimes_A E = F \otimes_A E$ further if F' and F are B -modules, and $F' \rightarrow F$ an injective homomorphism of B -modules, we can consider this homomorphism as an injective homomorphism of A -modules. Because E is A -flat,

$$F' \otimes_A E \rightarrow F \otimes_A E \text{ is injective.}$$

- (3) Let $\phi : A \rightarrow B$ be a ring homomorphism, such that B is a flat A -module. If F is a flat B -module, then F is a flat A -module. In fact: if $E' \rightarrow E$ is injective, then $E' \otimes_A B \rightarrow E \otimes_A B$ is injective, and also $(E \otimes_A B) \otimes_B F' \rightarrow (E \otimes_A B) \otimes_B F$ is injective. But $(E' \otimes_A B) \otimes_B F' = E' \otimes_A F$; $(E \otimes_A B) \otimes_B F = E \otimes_A F$.

If an A -module E is not flat, we want to measure how far it is from being flat. For this purpose we introduce the functor Tor .

Definition 2 : A free resolution of E is an exact sequence: $\dots \rightarrow L_n \rightarrow L_{n-1} \rightarrow \dots \rightarrow L_1 \rightarrow L_0 \rightarrow E \rightarrow 0$, where all L_i are free A -modules.

The complex of the resolution is the sequence

$$(L.) \dots \rightarrow L_n \rightarrow L_{n-1} \rightarrow \dots \rightarrow L_1 \rightarrow L_0 \rightarrow 0.$$

Every module has a free resolution. Two resolutions are algebraically homotopy-equivalent. Forming the tensor products $L_i \otimes F$, we get

$$(L. \otimes F) \dots \rightarrow L_n \otimes F \rightarrow L_{n-1} \otimes F \rightarrow \dots \rightarrow L_1 \otimes F \rightarrow L_0 \otimes F \rightarrow 0.$$

Definition 3 :

$$\text{Tor}_n^A(E, F) = H_n(L. \otimes F) = \frac{\text{Ker}(L_n \otimes F \rightarrow L_{n-1} \otimes F)}{\text{Im}(L_{n+1} \otimes F \rightarrow L_n \otimes F)}$$

if $n \geq 1$, and $\text{Tor}_0^A(E, F) = \text{Coker}(L_1 \otimes F \rightarrow L_0 \otimes F) = E \otimes F$.

Basic properties of Tor :

- (1) $\text{Tor}_n(E, F)$ is independent of the choice of the resolution (up to a canonical isomorphism).

- (2) If we take a free resolution of F , we get $\text{Tor}_n(F, E) = \text{Tor}_n(E, F)$ (Symmetry of the Tor). We can also define $\text{Tor}_n(E, F)$ by taking two free resolutions, one of E and one of F .
- (3) If $0 \rightarrow E' \rightarrow E \rightarrow E'' \rightarrow 0$ is a short exact sequence, then we get a long exact sequence:

$$\begin{array}{ccccccc} \text{Tor}_n(E', F) & \rightarrow & \text{Tor}_n(E, F) & \rightarrow & \text{Tor}_n(E'', F) & \rightarrow & \\ \rightarrow & \text{Tor}_{n-1}(E', F) & \rightarrow & \text{Tor}_{n-1}(E, F) & \rightarrow & \text{Tor}_{n-1}(E'', F) & \rightarrow \\ \rightarrow & \text{---} & \rightarrow & \text{---} & \rightarrow & \text{---} & \rightarrow \\ \rightarrow & \text{Tor}_1(E', F) & \rightarrow & \text{Tor}_1(E, F) & \rightarrow & \text{Tor}_1(E'', F) & \rightarrow \\ \rightarrow & E' \otimes F & \rightarrow & E \otimes F & \rightarrow & E'' \otimes F & \rightarrow 0. \end{array}$$

- (4) Tor is compatible with inductive limit, i.e. if $E = \lim (E_i)$, then
- $$\text{Tor}_n(\lim E_i, F) = \lim (\text{Tor}_n(E_i, F)).$$

- (5) We can define $\text{Tor}_n(E, F)$ by taking a flat resolution of E .

Proposition 3: Let E be an A -module. Then the following conditions are equivalent:

- (a) E is flat.
 (b) For all A -modules F , and for all $n \geq 1$, $\text{Tor}_n(E, F) = 0$.
 (c) For all A -modules F , $\text{Tor}_1(E, F) = 0$.

Proof: (a) \Rightarrow (b). If $\dots \rightarrow L_n \rightarrow L_{n-1} \rightarrow \dots \rightarrow L_1 \rightarrow L_0 \rightarrow F \rightarrow 0$ is a free resolution of F , then the sequence

$$\dots \rightarrow E \otimes L_n \rightarrow E \otimes L_{n-1} \rightarrow \dots \rightarrow E \otimes L_1 \rightarrow E \otimes L_0 \rightarrow E \otimes F \rightarrow 0$$

is exact, thus $\text{Tor}_n(E, F) = 0$ for all $n \geq 1$.

(b) \Rightarrow (c) clear. (c) \Rightarrow (a): If the sequence $0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0$ is exact, so is also (by (3) above) $\text{Tor}_1(E, F'') \rightarrow E \otimes F' \rightarrow E \otimes F \rightarrow E \otimes F'' \rightarrow 0$. Now $\text{Tor}_1(E, F'') = 0$, thus E is flat.

Proposition 4: If I and J are two ideals in A , then $\text{Tor}_1^A(A/I, A/J) = I \cap J / I \cdot J$.

Proof: From the exact sequence $0 \rightarrow I \rightarrow A \rightarrow A/I \rightarrow 0$, we get the exact sequence:

$$\text{Tor}_1(A, A/J) \rightarrow \text{Tor}_1(A/I, A/J) \rightarrow I \otimes A/J \rightarrow A \otimes A/J \rightarrow A/I \otimes A/J \rightarrow 0.$$

But now $\text{Tor}_1(A, A/J) = 0$ (A being A -free), and $I \otimes A/J = I/I \cdot J$; $A \otimes A/J = A/J$. Therefore the sequence $0 \rightarrow \text{Tor}_1(A/I, A/J) \rightarrow I/I \cdot J \rightarrow A/J$ is exact, and $\text{Tor}_1(A/I, A/J) = \text{Ker}(I/I \cdot J \rightarrow A/J) = I \cap J / I \cdot J$.

Example : Let U be an open set in \mathbf{C}^n , and $x \in U$. Further let $X, Y \subset U$ be two hypersurfaces, defined by $I = (f)$ and $J = (g)$. Supposing that f and g do not have common factors: $I_x \cap J_x = I_x J_x$, and

$$\text{Tor}_1(\mathcal{O}_{X,x}, \mathcal{O}_{Y,x}) = \text{Tor}_1(\mathcal{O}_{U,x}/I_x, \mathcal{O}_{U,x}/J_x) = \frac{I_x \cap J_x}{I_x \cdot J_x} = 0.$$

Heuristic remark : The formula $\text{Tor}_1(\mathcal{O}_{X,x}, \mathcal{O}_{Y,x}) = 0$ expresses the fact that X and Y are “in general position”. If for example X and Y are two linear subspaces in \mathbf{C}^n of dimensions p and q , we have $\text{Tor}_1(\mathcal{O}_{X,x}, \mathcal{O}_{Y,x}) = 0$ if $\dim(X \cap Y) = p + q - n$, and $\text{Tor}_1(\mathcal{O}_{X,x}, \mathcal{O}_{Y,x}) \neq 0$ otherwise.

Next we shall prove an elementary flatness criterion.

Proposition 5 : Let E be an A -module. The following conditions are equivalent:

- (a) E is flat.
- (b) For all finitely generated ideals I of A , $\text{Tor}_1(E, A/I) = 0$.
- (c) For all monogenous A -modules F , $\text{Tor}_1(E, F) = 0$.

Proof : (a) \Rightarrow (b), by prop. 3.

(b) \Rightarrow (c): Because Tor is compatible with inductive limit, we can suppose, that $\text{Tor}_1(E, A/I) = 0$ for an arbitrary ideal I of A . But every monogenous A -module F can be represented by A/I .

(c) \Rightarrow (a). By prop. 3 it is sufficient to prove that $\text{Tor}_1(E, F) = 0$ for any A -module F .

First consider the case, where F is finitely generated. We use induction, supposing that $\text{Tor}_1(E, F) = 0$, when F has n generators. Let F have $(n+1)$ generators x_1, \dots, x_n, x_{n+1} . If F' is the submodule generated by $\{x_1, \dots, x_n\}$, then $F' \subset F$ and $F'' = F/F'$ is monogenous. The exact sequence $0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0$ gives the exact sequence $\text{Tor}_1(E, F') \rightarrow \text{Tor}_1(E, F) \rightarrow \text{Tor}_1(E, F'')$. Now $\text{Tor}_1(E, F') = \text{Tor}_1(E, F'') = 0$, thus $\text{Tor}_1(E, F) = 0$. In the general case, F can be considered as an inductive limit of finitely generated modules, and because Tor is compatible with inductive limits, $\text{Tor}_1(E, F) = 0$.

Proposition 6 : Let A be an integral domain, and E an A -module. Then E is torsionfree if and only if $\text{Tor}_1(E, A/(a)) = 0$, for any element $a \in A$.

Proof : If E is A -module, $a \in A$, then the exact sequence $0 \rightarrow A \xrightarrow{aI} A \rightarrow A/(a) \rightarrow 0$ gives the exact sequence $0 \rightarrow \text{Tor}_1(E, A/(a)) \rightarrow E \xrightarrow{aI} E$. In other words $\text{Tor}_1(E, A/(a)) = \{x \in E \mid ax = 0\}$, from which the result follows.

Corollary: Let A be a principal ideal domain. E is flat if and only if E is torsionfree.

Proof: We have already proved that, if E is flat, then it is torsion free. The converse follows from prop. 6 and prop. 5.

The first flatness criterion for noetherian local rings is the following:

Theorem 2: Let A be a noetherian local ring with maximal ideal m ; $k = A/m$, and E a finitely generated A -module. The following conditions are equivalent:

- (a) E is free.
- (b) E is flat.
- (c) $\text{Tor}_1^A(E, k) = 0$.

Proof: We have already proved $(a) \Rightarrow (b) \Rightarrow (c)$.

$(c) \Rightarrow (a)$: We recall first Nakayma's lemma. If A is a local ring with maximal ideal m ; $k = A/m$, and E is a finitely generated A -module, such that $k \otimes_A E = E/mE = 0$, then $E = 0$.

The module $\bar{E} = k \otimes_A E = E/mE$ is a finitely generated vector space over k . Let $\{\bar{x}_1, \dots, \bar{x}_r\}$ be a base of \bar{E} (over k), and $\{x_1, \dots, x_r\}$ E representatives of \bar{x}_i : s . Consider the homomorphism $\phi : A^r \rightarrow E$, $\phi(a_1, \dots, a_r) = \sum a_i x_i$. Denoting by R and Q the kernel and the cokernel of ϕ , we get an exact sequence:

$$(*) \quad 0 \rightarrow R \rightarrow A^r \rightarrow E \rightarrow Q \rightarrow 0$$

and R, Q are finitely generated A -modules. From $(*)$ we get the exact sequence

$$A^r \otimes_A k \rightarrow E \otimes_A k \rightarrow Q \otimes_A k \rightarrow 0.$$

But $\bar{E} = E \otimes_A k \simeq k^r = A^r \otimes_A k$, so $Q \otimes_A k = 0$, and by Nakayama's lemma $Q = 0$.

Therefore we have an exact sequence

$$0 \rightarrow R \rightarrow A^r \rightarrow E \rightarrow 0.$$

From this we get: $\text{Tor}_1(E, k) \rightarrow k \otimes_A R \rightarrow k^r \rightarrow \bar{E} \rightarrow 0$ (exact). Now: $\bar{E} \simeq k^r$, $\text{Tor}_1(E, k) = 0$ (by assumption). Therefore $k \otimes_A R = 0$, and once more by Nakayama's lemma $R = 0$, thus $E \simeq A^r$, i.e. E is free.

Proposition 7: Let $\phi : A \rightarrow B$ be a ring homomorphism, and let B be A -flat. If I is an ideal of A , we write $\bar{A} = A/I$, $\bar{B} = B/IB = \bar{A} \otimes_A B$. Let F be a B -module, then: $\text{Tor}_i^A(\bar{A}, F) = \text{Tor}_i^B(\bar{B}, F)$ ($i \geq 0$).

Proof: We choose first a B -free resolution of F

$$\rightarrow L_{n+1} \rightarrow L_n \rightarrow \dots \rightarrow L_1 \rightarrow L_0 \rightarrow F \rightarrow 0.$$

If $L.$ is the respective complex of resolution, then

$$\bar{B} \otimes_B L. = B/IB \otimes_B L. = \bar{A} \otimes_A (B \otimes_B L.) = \bar{A} \otimes_A L.$$

Because every L_i is B -free, and B is A -flat, every L_i is A -flat (Property 3 after Th. 1). Thus $L.$ is a flat A -resolution, and

$$\text{Tor}_i^A(\bar{A}, F) = H_i(\bar{A} \otimes_A L.) = H_i(\bar{B} \otimes_B L.) = \text{Tor}_i^B(\bar{B}, F).$$

We shall next state the second flatness criterion for noetherian local rings.

Theorem 3: Let A and B be two noetherian local rings, with maximal ideals $\underline{m}, \underline{n}; k = A/\underline{m}$. If $\phi : A \rightarrow B$ is a local homomorphism (i.e. $\phi(\underline{m}) \subset \underline{n}$), and F finitely generated B module then

$$F \text{ is } A\text{-flat} \Leftrightarrow \text{Tor}_1^A(k, F) = 0.$$

The proof of this theorem is much more difficult than that of th. 20 see for example:

Bourbaki: *Algèbre commutative*, Chapter III § 5, th1, (i) \Leftrightarrow (iii), p. 98.

The conditions in Bourbaki's theorem are here fulfilled:

- 1° A finitely generated module F over a noetherian local ring B is idealwise separated for \underline{n} . (*Ibid.*, § 5. 1. Ex. 1, p. 97.)
- 2° If $\phi : A \rightarrow B$ is a local homomorphism, F is also idealwise separated for \underline{m} . (*Ibid.*, § 5, prop. 2, p. 101.)
- 3° Also the flatness condition is fulfilled, because k is a field.

Remark: The main interest of the theorem lies in the fact, that it is true without any assumption of finiteness on B .

Corollary: If the assumptions are the same as in the theorem 3, and if moreover B is A -flat, then

$$F \text{ is } A\text{-flat} \Leftrightarrow \text{Tor}_1^B(\bar{B}, F) = 0,$$

where $\bar{B} = B/\underline{m}B$.

Proof: $\text{Tor}_1^A(k, F) = \text{Tor}_1^B(\bar{B}, F)$, by prop. 7.

§ 5. *Geometric applications of the flatness criterions*

A) *Flatness for finite morphisms*

Proposition 1: Let $\pi: X \rightarrow S$ be a finite morphism (i.e. proper with finite fibres) of analytic spaces. Then $\pi_*(\mathcal{O}_X)$ is a coherent analytic sheaf over S . The following conditions are equivalent:

- (a) π is flat (i.e. for every $x \in X$, $\mathcal{O}_{X,x}$ is a flat $\mathcal{O}_{S,s}$ -module, $s = \pi(x)$).
- (b) For every s , $(\pi_* \mathcal{O}_X)_s$ is a flat $\mathcal{O}_{S,s}$ -module.
- (c) $\pi_* \mathcal{O}_X$ is a locally free sheaf.

Proof: Because π is finite $\pi_*(\mathcal{O}_X)_s = \bigoplus_{x \in \pi^{-1}(s)} \mathcal{O}_{X,x}$, thus the only point to prove is (b) \Rightarrow (c).

Now if $\mathcal{O}_{X,x}$ is a flat $\mathcal{O}_{S,s}$ -module, then (by theorem 2) $\mathcal{O}_{X,x}$ is free, and a coherent sheaf whose fibers are free is a locally free sheaf.

Proposition 2: Let S be a reduced analytic space and \mathcal{E} a coherent \mathcal{O}_S -module. Let $E(s)$ be the finite dimensional vector space (over \mathbb{C}) $\mathcal{E}_s \otimes_{\mathcal{O}_{S,s}} \mathbb{C}_s$. \mathcal{E} is a locally free $\mathcal{O}_{S,s}$ -module if and only if $\dim_{\mathbb{C}} E(s)$ is locally constant.

Proof: If \mathcal{E} is locally free, then $\dim_{\mathbb{C}} E(s)$ is locally constant. Suppose now that $\dim_{\mathbb{C}} E(s)$ is locally constant in an open set $U \subset S$, and that $\mathcal{O}_U^p \xrightarrow{d} \mathcal{O}_U^q \rightarrow \mathcal{E}_U \rightarrow 0$ is exact. d is determined by a $p \times q$ matrix of analytic functions on U , so it gives a morphism $\mathbb{C}_U^p \xrightarrow{d} \mathbb{C}_U^q$ of trivial vector bundles over U .

From the exact sequence $\mathcal{O}_s^p \xrightarrow{d_s} \mathcal{O}_s^q \rightarrow \mathcal{E}_s \rightarrow 0$, we get (by making tensor-products with \mathbb{C}_s) the exact sequence:

$$\mathbb{C}_s^p \xrightarrow{d(s)} \mathbb{C}_s^q \rightarrow E(s) \rightarrow 0,$$

which shows that d has constant rank in U . Thus $\text{Ker } d$ and $\text{Im } d$ are vector bundles, and we can write

$$\mathbb{C}_U^p = F_1 \oplus G_1, \quad \mathbb{C}_U^q = F_0 \oplus G_0,$$

$$d : \begin{cases} F_1 \rightarrow 0 \\ G_1 \simeq F_0. \end{cases}$$

Now $\mathcal{E} \simeq$ the sheaf of analytic sections of G_0 , therefore \mathcal{E} is locally free.

Definition 1: Let $\pi : X \rightarrow S$ be a finite morphism of analytic spaces, and $s \in S$. For each $x \in X(s) = \pi^{-1}(s)$, $\mathcal{O}_{X(s),x} = \mathbf{C} \otimes_{\mathcal{O}_{S,s}} \mathcal{O}_{X,x}$ is finite dimensional vectorspace over \mathbf{C} . Denote its dimension by $v(x)$. Then the degree $v(s)$ of s is defined by $v(s) = \sum_{x \in X(s)} v(x)$.

Theorem 1: Let $\pi : X \rightarrow S$ be a finite morphism of analytic space and let S be a reduced space. Then X is flat over S if and only if $v(s)$ is locally constant function of s .

$$\begin{aligned} \text{Proof: } v(s) &= \sum_{x \in X(s)} \dim_{\mathbf{C}} \mathcal{O}_{X(s),x} = \dim_{\mathbf{C}} \left(\bigoplus_{x \in X(s)} \mathcal{O}_{X(s),x} \right) \\ &= \dim_{\mathbf{C}} \left(\bigoplus_{x \in X(s)} (\mathbf{C} \otimes_{\mathcal{O}_{S,s}} \mathcal{O}_{X,x}) \right) \\ &= \dim_{\mathbf{C}} \mathbf{C} \otimes_{\mathcal{O}_{S,s}} \pi_* (\mathcal{O}_X)_s = \dim_{\mathbf{C}} E(s). \end{aligned}$$

The theorem follows from propositions 1 and 2.

Examples of flat morphisms

Example 1: If $\pi : X \rightarrow S$ is a local isomorphism near x , then π is flat at x .

Example 2: Consider § 2, Ex. 1. Here $v(x) = 1$.

Examples of non-flat morphisms

Examples 1: If $X \subset S$ is a closed subspace, not open, $v(s)$ is not locally constant.

Example 2: Let X be a subspace of \mathbf{C}^4 defined by the ideal intersection of (x_3, x_4) and $(x_1 - x_1, x_4 - x_2)$ (which is equal to the product ideal) and let π be the projection onto the (x_1, x_2) -plane \mathbf{C}^2 . Then X is a union of two 2-planes in \mathbf{C}^4 , whose intersection is (0) . When $s \neq 0$, $X(s)$ consists of two simple points, so $v(s) = 2$. $X(0)$ is given by the ideal $(x_1, x_2, x_3^2, x_3x_4, x_4^2)$, thus $v(0) = 3$.

Example 3: Let $S = \{(u, v, w) \in \mathbf{C}^3 \mid v^2 = uw\}$ and $\pi : \mathbf{C}^2 \rightarrow S$ be the map $(x, y) \rightarrow (x^2, xy, y^2)$. This map identifies S with the quotient of \mathbf{C}^2 by the equivalence relation identifying (x, y) with $(-x, -y)$. However, π is not flat, since for $s \in S$, $v(s) = 2$ if $s \neq 0$ and $v(s) = 3$ if $s = 0$.

B) *Projection of a product of analytic spaces*

Theorem 2: Let S and X be analytic spaces. If $\pi : S \times X \rightarrow S$ is the projection morphism, then π is flat, i.e. $\mathcal{O}_{S \times X, (s,x)}$ is a flat $\mathcal{O}_{S,s}$ module for every $(s, x) \in S \times X$.

To prove this theorem we need first some homological algebra. Then we shall show it in the particular case, when S is a manifold, and finally in the general case.

(a) *Koszul complex*

Let A be a ring, M an A -module and h_1, \dots, h_n homomorphisms $M \rightarrow M$, which commute with each other, i.e. $h_i h_j = h_j h_i$ for every i, j .

If $1 \leq k \leq n$, set $Q_k = M/h_1(M) + \dots + h_k(M)$, and $Q_0 = M$, thus, in particular, $Q_n = Q = M / \sum_{i=1}^n h_i(M)$. Every h_k induces a map $\tilde{h}_k : Q_{k-1} \rightarrow Q_{k-1}$.

Definition 2: The sequence (h_1, \dots, h_n) is called regular if each of the mappings \tilde{h}_k ($1 \leq k \leq n$) is injective.

The Koszul complex of the module M and of the mappings h_k ($1 \leq k \leq n$) $K. = K. [M; h_1, \dots, h_n]$ is defined in the following way:

$$K_i = \wedge^{n+i} A^n \otimes M \simeq M^{\binom{n}{i}}, \quad 0 \leq i \leq n.$$

We define the homomorphisms $d_i : K_i \rightarrow K_{i-1}$ ($i > 0$) by $\lambda \otimes x \rightarrow \sum_i (e_i \wedge \lambda) \otimes \otimes h_i(x)$, where (e_i) is the natural base of A^n . We also define $\varepsilon : K_0 \rightarrow Q$ as the natural map $: K_0 = M \rightarrow M / \sum_{i=1}^n h_i(M) = Q$. Using the fact that h_1, \dots, h_n commute with each other, it is easy to verify that

$$(d_{i-1} \circ d_i)(\lambda \otimes x) = \sum_{i,j} (e_j \wedge e_i \wedge \lambda) \otimes h_j(h_i(x)) = 0;$$

also $\varepsilon d_1 = 0$. Thus $K.$ is really a complex.

Theorem 3 (Poincaré-Koszul).

If (h_1, \dots, h_n) is a regular sequence, then

$$H_i(K.) = \begin{cases} Q & \text{if } i = 0 \\ \cdot & \text{if } i = 1 \\ 0 & \text{if } i > 1 \end{cases}$$

If $h_i \in A$, it defines the map: $A \xrightarrow{h_i I} A$, which we denote also by h_i . We say that (h_1, \dots, h_n) is a regular sequence of elements if $(h_1 I, \dots, h_n I)$ is a regular sequence.

Corollary. If (h_1, \dots, h_n) is a regular sequence of elements, then the Koszul complex $K. = K. [A; h_1, \dots, h_n] = \{ \wedge^{n-1} A^n \simeq A^{(n)} \}$ is a free resolution of $Q = A/(h_i)$ ((h_i) is the ideal generated by h_1, \dots, h_n).

Example: If $A = \mathbf{C} \{x_1, \dots, x_n\}$; $h_i = x_i$, then $Q_k = A/(x_1, \dots, x_k) = \mathbf{C} \{x_{k+1}, \dots, x_n\}$ and $Q = Q_n = \mathbf{C}$. The complex $K. = K. [A; x_1, \dots, x_n]$ is a free resolution of \mathbf{C} .

(b) *Proof of theorem 2, when S is a complex manifold*

In this case we can take $\mathcal{O}_{S,S} = \mathbf{C} \{t_1, \dots, t_m\} = A$ and if $\mathcal{O}_{X,x} = \mathbf{C} \{x_1, \dots, x_n\}/(f_1, \dots, f_p)$, then

$$\mathcal{O}_{S \times X, (s,x)} = \mathbf{C} \{t_1, \dots, t_m, x_1, \dots, x_n\}/(f_1, \dots, f_p) = B.$$

B is an A -module in a natural way.

By the corollary of the Poincaré-Koszul theorem $K. = K. [A; t_1, \dots, t_m]$ in a free resolution of \mathbf{C} . We want to compute the modules $\text{Tor}_i^A(\mathbf{C}, B) = H_i(K. \otimes B)$ ($i > 0$).

It's easily seen, that we can consider the complex $K. \otimes B$ as a Koszul

complex $K'. = K. [B; t_1, \dots, t_m]$ (where $t_i : B \xrightarrow{t_i I} B$). But now the sequence (t_1, \dots, t_m) is regular, thus by the Poincaré-Koszul theorem $H_i[K'] = 0$ if $i > 0$.

In particular: $\text{Tor}_1^A(\mathbf{C}, B) = H_1[K. \otimes B] = H_1[K'] = 0$. By the second flatness criterion B is A -flat.

(c) *The general case*

The question being local, we can suppose that $S \subset W \subset \mathbf{C}^n$, where W is open, and S an analytic subspace of W . Let S be defined by g_1, \dots, g_r . Then $S \times X \subset W \times X$ and $\mathcal{O}_S = \mathcal{O}_W/(g_1, \dots, g_r)$. On the other hand $\mathcal{O}_{S \times X} = \mathcal{O}_{W \times X}/(g_1, \dots, g_r) = \mathcal{O}_S \otimes_{\mathcal{O}_W} \mathcal{O}_{W \times X}$. The last equality follows from

the fact, that if $\pi : X \rightarrow S$ is a morphism, and $S' \subset S$ a subspace, $X' = \pi^{-1}(S')$,

$$\text{then } \mathcal{O}_{X'} = \mathcal{O}_{S'} \otimes_{\mathcal{O}_S} \mathcal{O}_X.$$

Remark : This a particular case of the following proposition: if π and π' are two morphisms of which at least one is finite, then

$$\begin{array}{ccc} X & & Y \\ \pi \searrow & & \swarrow \pi' \\ & S & \end{array} \quad \mathcal{O}_{X \times_S Y} = \mathcal{O}_X \otimes_{\mathcal{O}_S} \mathcal{O}_Y.$$

We have proved that $\mathcal{O}_{W \times X}$ is \mathcal{O}_W -flat, so by scalar extension $\mathcal{O}_{S \times X}$ is \mathcal{O}_S flat.

Corollary : If X and S are two manifolds and $\pi : X \rightarrow S$ is a submersion, then π is flat.

III. PRIVILEGED POLYCYLINDERS

§ 1. Banach vector bundles over an analytic space

Let E be a Banach space and X an analytic space. We denote then by E_X the trivial bundle $X \times E$ over X .

To define bundle morphisms, we first define the sheaf $\mathcal{H}_X(E)$ of germs of analytic morphisms from X to E . If $U \subset \mathbb{C}^n$ is open, then the set $\mathcal{H}(U, E)$ of analytic morphisms from U into E consists of all functions $g : U \rightarrow E$ having at every point $x \in U$ a converging power series expansion.

Let now X' be a local model for X , i.e. X' is the support of the quotient sheaf \mathcal{O}_U/J , where $U \subset \mathbb{C}^n$ is open and J is a coherent sheaf of ideals of \mathcal{O}_U , then $\mathcal{H}_{X'}(E)$ is the sheaf associated to the presheaf $V \rightarrow \mathcal{H}(V, E)/J_V \cdot \mathcal{H}(V, E)$ ($V \subset U$, V -open).

Remark : If X' is reduced, the sections of $\mathcal{H}_{X'}(E)$ are just the functions from X' to E which are locally induced by analytic functions on open sets in U .


The sheaf $\mathcal{H}_X(E)$ is constructed with help of the local models X' of X , i.e. $\mathcal{H}_X(E)|_{X'} = \mathcal{H}_{X'}(E)$, for every local model X' .

Definition 1 : The set of *analytic morphisms* from an analytic space X into a Banach space E is the set $\mathcal{H}(X; E)$ of sections of the sheaf $\mathcal{H}_X(E)$.

Let $\mathcal{L}(E, F)$ be the Banach space of all continuous linear mappings from the Banach space E into the Banach space F .

Definition 2 : An *analytic vector bundle morphism* from E_X into F_X is an analytic morphism from X into $\mathcal{L}(E, F)$.

Let E be a topological space, X an analytic space, and $\pi : E \rightarrow X$ a continuous projection.



Suppose that X has an open covering $(U_\iota)_{\iota \in I}$, and that for every $\iota \in I$ there is given a trivial Banach space bundle $E|_{U_\iota}$ and a homeomorphism ϕ_ι , such that the following diagram is commutative:

We suppose further that for each pair $\iota, \kappa \in I$ there is given an analytic vector bundle morphism $\gamma_{\iota\kappa} : E|_{U_\iota \cap U_\kappa} \rightarrow E|_{U_\iota \cap U_\kappa}$, with the underlying mapping $\phi_\iota \circ \phi_\kappa^{-1}$, such that:

$$\gamma_{\iota\lambda} = \gamma_{\iota\kappa} \gamma_{\kappa\lambda}; \quad \gamma_{\iota\iota} = I, \quad \text{for all } \iota, \kappa, \lambda \in I.$$

This data gives a Banach vector bundle atlas on E and provides E with the structure of a Banach vector bundle over X (two atlases are equivalent if there exists an atlas containing both).

Remark: If X is reduced, the $\gamma_{\iota\kappa}$ are determined by their underlying map and the condition $\gamma_{\iota\lambda} = \gamma_{\iota\kappa} \gamma_{\kappa\lambda}$ is automatically satisfied.

Using local triviality, we can define morphisms for general Banach vector bundles.

Proposition 1: Let $\phi : E \rightarrow F$ be a morphism of two Banach vector



bundles E and F , and $x \in X$.

If $\phi_x \in \mathcal{L}(E(x), F(x))$ is an isomorphism, then there exists an open neighbourhood $U \subset X$ of x , such that $\phi|_U : E|_U \rightarrow F|_U$ is a vector bundle isomorphism.

Proof: First we take a trivialisation $E|_V = E_0|_V, F|_V = F_0|_V$ at $x \in V \subset X$ (V -open).

The set $\text{Isom}(E_0, F_0)$ of isomorphic mappings is an open subset of $\mathcal{L}(E_0, F_0)$ and the mapping $g \rightarrow g^{-1}$ is an analytic isomorphism:


$$\text{Isom}(E_0, F_0) \simeq \text{Isom}(F_0, E_0).$$

So we have in an open neighbourhood $U \subset X$ of x an analytic morphism $y \rightarrow \phi_y^{-1} \in \mathcal{L}(F_0, E_0)$, which defines the inverse morphism $(\phi|_U)^{-1} : F|_U \rightarrow E|_U$.

Definition 3 : Let E and F be two Banach spaces and f a continuous linear mapping from E into F . f is a *split mono-(epi) morphism*, if there exists a mapping $g \in \mathcal{L}(F, E)$ such that $g \circ f = I_E$. (Resp. $f \circ g = I_F$.)

Definirion 4 : Let E_1 and E_2 be two Banach vector bundles over an analytic space X , and f a vector bundle morphism from E_1 into E_2 . f is a *split mono (epi) morphism*, if there exists a vector bundle morphism $g : E_2 \rightarrow E_1$ such that $g \circ f = I_{E_1}$. (Resp. $f \circ g = I_{E_2}$.)

Equivalently, $f : E_1 \rightarrow E_2$ is a split monomorphism if and only if E_2 can




be decomposed in a direct sum $E_2 = F_2 \oplus G_2$ such that

$$f: \begin{cases} E_1 \simeq F_2 \\ 0 \rightarrow G_2 \end{cases}.$$

and f is a split epimorphism if correspondingly

$$E_1 = F_1 \oplus G_1, \quad \text{such that } f: \begin{cases} F_1 \rightarrow 0 \\ G_1 \simeq E_2 \end{cases}.$$

Proposition 2 : Let $E \xrightarrow{\phi} F$ be a bundle morphism and $x \in X$.



If $\phi_x : E(x) \rightarrow F(x)$ is a split epi (mono) morphism, then the point x has an open neighbourhood $U \subset X$, such that $\phi|_U : E|_U \rightarrow F|_U$ is a split vector bundle epi (mono) morphism.

Proof : Suppose that ϕ_x is a split epimorphism. We take first a trivilisation $E|_V = E_{0V}, F|_V = F_{0V}$ at x , so that there exists a mapping $\sigma \in \mathcal{L}(F_0, E_0)$, $\phi_x \circ \sigma = I_{F_0}$. If we define a morphism $\psi : F_{0V} \rightarrow E_{0V}$ by $x \rightarrow \sigma \in \mathcal{L}(F_0, E_0)$, the morphism $\gamma = \phi \circ \psi : F_{0V} \rightarrow E_{0V}$ has an isomorphic fibre mapping $\gamma_x = I_{F_0}$ in x . By proposition 1 we have an isomorphic restriction $\gamma|_U, \phi|_U \circ (\psi|_U \circ (\gamma|_U)^{-1}) = I_{F_{0U}}$.

When ϕ_x is a split monomorphism, the proof is similar.

Definition 5 : Let B_1, B_2, B_3 be Banach spaces, and $j, k : B_1 \xrightarrow{j} B_2 \xrightarrow{k} B_3$ continuous linear mappings. This sequence forms a *complex*, if $k \circ j = 0$. This sequence is *split exact* if the space B_i can be decomposed in direct

sums $B_i = C_i \oplus D_i$ such that

$$j: \begin{cases} C_1 \rightarrow 0 \\ D_1 \simeq C_2 \end{cases} \quad k: \begin{cases} C_2 \rightarrow 0 \\ D_2 \simeq C_3 \end{cases} .$$

Definition 6: A Banach vector bundle morphism sequence

$$\begin{array}{ccccc} E_1 & \xrightarrow{f} & E_2 & \xrightarrow{g} & E_3 \\ & \searrow & \downarrow X & \swarrow & \\ & & X & & \end{array} \quad \text{is a complex if } g \circ f = 0.$$

The sequence is *split exact*, if every E_i can be decomposed $E_i = F_i \oplus G_i$, such that:

$$f: \begin{cases} F_1 \rightarrow 0 \\ G_1 \simeq F_2 \end{cases} \quad g: \begin{cases} F_2 \rightarrow 0 \\ G_2 \simeq F_3 \end{cases} .$$

Theorem 1: Let $\begin{array}{ccccc} E_1 & \xrightarrow{f} & E_2 & \xrightarrow{g} & E_3 \\ & \searrow & \downarrow X & \swarrow & \end{array}$ be a complex of Banach vector

bundles and $x_0 \in X$.

If the sequence of Banach spaces $E_1(x_0) \xrightarrow{f_{x_0}} E_2(x_0) \xrightarrow{g_{x_0}} E_3(x_0)$ is split exact, then there exists an open neighbourhood $U \subset X$ of x_0 , such that $E_1|_U \xrightarrow{f|_U} E_2|_U \xrightarrow{g|_U} E_3|_U$ is a split exact sequence of Banach vector bundles.

Proof: We take a neighbourhood V of x , such that we have a complex $E_{1V} \xrightarrow{f|_V} E_{2V} \xrightarrow{g|_V} E_{3V}$ of trivial bundles. By assumption we have the decompositions $E_{iV}(x_0) = F_i(x_0) \oplus G_i(x_0)$ with

$$f_{x_0}: \begin{cases} F_1(x_0) \rightarrow 0 \\ G_1(x_0) \simeq F_2(x_0) \end{cases} \quad g_{x_0}: \begin{cases} F_2(x_0) \rightarrow 0 \\ G_2(x_0) \simeq F_3(x_0) \end{cases} .$$

By proposition 2, $f|_V : G_{1V} \rightarrow E_{2V}$, $g|_V : G_{2V} \rightarrow E_{3V}$ are both split monomorphisms in a neighbourhood $W \subset V$ of x_0 and the images $F_2 = f(G_{1W})$, $F_3 = g(G_{2W})$ are subbundles of E_{2W} esp. E_{3W} , such that

$$E_{2W} = F_2 \oplus G_{2W}, \quad E_{3W} = F_3 \oplus G_{3W} .$$

By our construction

$$g|_W : \begin{cases} F_2 & \rightarrow 0 \\ G_2 W & \simeq F_3 \end{cases} .$$

If $p: E_{2W} \rightarrow F_2$ is the projection with kernel G_{2W} , the map, $p \circ f: E_{1W} \rightarrow F_2$ is a split epimorphism in x_0 . Again by prop. 2 we have over an open neighbourhood $U \subset W$ of x_0 a decomposition $E_{1U} = F_1 \oplus G_{1U}$ (with $F_1 = \text{Ker } p \circ f$)

$$(p \circ f)|_U : \begin{cases} F_1 & \rightarrow 0 \\ G_{1U} & \xrightarrow{\sim} F_{2U} \end{cases} .$$

The image $f|_U(F_1)$ is contained in G_{2U} . But $g|_U \circ f|_U = 0$ and $g|_{G_{2U}}$ is a monomorphism hence $f|_U: F_1 \rightarrow 0$. We get finally (restricting all our morphisms to U)

$$f|_U : \begin{cases} F_{1U} & \rightarrow 0 \\ G_{1U} & \simeq F_{2U} \end{cases} \qquad g|_U : \begin{cases} F_{2U} & \rightarrow 0 \\ G_{2U} & \xrightarrow{\sim} F_{3U} \end{cases} .$$

§ 2. *Privileged polycylinders*

Definition 1: A polycylinder in \mathbf{C}^n is a compact set K of the form $K = K_1 \times \dots \times K_n$ where each K_i is a compact, convex subset of \mathbf{C} , with nonempty interior. If each K_i is a disc, then K is a polydisc. We first recall the following theorem of Cartan.

Theorem 1: Let K be a polycylinder contained in an open subset U of \mathbf{C}^n . Let \mathcal{F} be a coherent analytic sheaf on U .

(A) There exists an open neighbourhood of K over which \mathcal{F} admits a finite free resolution

$$0 \rightarrow \mathcal{L}_n \rightarrow \dots \rightarrow \mathcal{L}_1 \rightarrow \mathcal{L}_0 \rightarrow \mathcal{F} \rightarrow 0 .$$

(B) $H^q(K, \mathcal{F}) = 0$ for $q > 0$.

(Reference: For instance Gunning and Rossi.)

We have the following consequences of this theorem:

1) Given a finite free resolution

$$0 \rightarrow \mathcal{L}_n \rightarrow \dots \rightarrow \mathcal{L}_1 \rightarrow \mathcal{L}_0 \rightarrow \mathcal{F} \rightarrow 0$$

of a coherent sheaf \mathcal{F} , the sequence

$$0 \rightarrow \mathcal{L}_n(K) \rightarrow \dots \rightarrow \mathcal{L}_0(K) \rightarrow \mathcal{F}(K) \rightarrow 0$$

is an $\mathcal{O}_U(K)$ - free resolution of $\mathcal{F}(K)$.

2) Given a short exact sequence of coherent sheaves

$$0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0,$$

then the sequence

$$0 \rightarrow \mathcal{F}'(K) \rightarrow \mathcal{F}(K) \rightarrow \mathcal{F}''(K) \rightarrow 0 \quad \text{is exact.}$$

Let \mathcal{F} be a coherent analytic sheaf on U , and let $K \subset U$ be a polycylinder. If V is an open neighbourhood of K , then $\mathcal{F}(V)$ can be equipped with a Fréchet-space structure (see: Malgrange).

Hence we can give $\mathcal{F}(K)$ the structure of inductive limit of Fréchet-spaces. It is however essential for certain purposes to have Banach-spaces. This can be obtained by choosing a space slightly different from $\mathcal{F}(K)$ and by choosing K in a "privileged" way.

Let $B(K) = \{f: K \rightarrow \mathbb{C} \mid f \text{ continuous on } K \text{ and analytic on } \overset{\circ}{K}\}$, then $B(K)$ is Banach algebra and $B(K) \subset C(K)$. The sections of \mathcal{O}_U over K are elements of $B(K)$, and $B(K)$ is in fact the uniform closure of $\mathcal{O}_U(K)$ in $C(K)$.

If $\mathcal{L} = \mathcal{O}_U^r$, we define $B(K, \mathcal{L}) = B(K)^r$. Then $B(K; \mathcal{L})$ is a free $B(K)$ -module, and since $\mathcal{L}(K) = \mathcal{O}_U(K)^r$, we have $B(K; \mathcal{L}) = B(K) \otimes_{\mathcal{O}_U(K)} \mathcal{L}(K)$.

We now assume that \mathcal{F} is a coherent sheaf on U , where $U \subset \mathbb{C}^n$ is open. Consider a free resolution

$$(R) \quad 0 \rightarrow \mathcal{L}_n \rightarrow \dots \rightarrow \mathcal{L}_1 \rightarrow \mathcal{L}_0 \rightarrow \mathcal{F} \rightarrow 0 \quad \text{of } \mathcal{F}.$$

From (R) we get an $\mathcal{O}_U(K)$ -free resolution of $\mathcal{F}(K)$

$$(R') \quad 0 \rightarrow \mathcal{L}_n(K) \rightarrow \dots \rightarrow \mathcal{L}_1(K) \rightarrow \mathcal{L}_0(K) \rightarrow \mathcal{F}(K) \rightarrow 0.$$

Taking the tensorproduct $B(K) \otimes_{\mathcal{O}_U(K)}$ we get the complex

$$B(K; \mathcal{L}.): 0 \rightarrow B(K; \mathcal{L}_n) \rightarrow \dots \rightarrow B(K; \mathcal{L}_1) \rightarrow B(K; \mathcal{L}_0).$$

Definition 2: The polycylinder K is called \mathcal{F} -privileged if the complex $B(K; \mathcal{L}.)$ is split-exact in every degree > 0 .

Remark: The property of being \mathcal{F} -privileged is independent of the resolution (R).

The exactness of $B(K; \mathcal{L}.)$ can be expressed by $\text{Tor}_i^{\mathcal{O}_U(K)}(B(K), \mathcal{F}(K)) = 0$, for every $i > 0$, and Tor is independent of the resolution (R). It is a little

more complicated to show, that the splitting property is independent of (R) , and this is omitted.

Since $B(K; \mathcal{L}_i)$ is a Banach space, the image and its complement are thus Banach spaces if K is \mathcal{F} -privileged. In this case we define $B(K; \mathcal{F}) = \text{Coker}(B(K, \mathcal{L}_1) \rightarrow B(K; \mathcal{L}_0)) = B(K) \otimes_{\mathcal{O}_U} \mathcal{F}(K)$ and we get a $B(K)$ -module, which is a Banach-space.

Warning: In the definition of split-exactness, the subspaces are splitting vector spaces, but they are not splitting $B(K)$ -modules in general.

We have the following important theorem about the existence of privileged polycylinders:

Theorem 2: Let U be an open subset of \mathbf{C}^n , and let \mathcal{F} be a coherent analytic sheaf on U . For any $x \in U$ there exists a fundamental system of neighbourhoods of x in U , which are \mathcal{F} -privileged polycylinders.

For the proof, see Douady: § 7, 4, th 1.

Example: (Curves in \mathbf{C}^2) Let $U \subset \mathbf{C}^2$ be an open connected neighbourhood of the origin, and let $h: U \rightarrow \mathbf{C}$ be analytic and $h \neq 0$.

Let X be the curve given by h , that is $X = h^{-1}(0)$, $\mathcal{O}_X = \mathcal{O}_U / (h)$. We have an exact sequence $0 \rightarrow \mathcal{O}_U \xrightarrow{h} \mathcal{O}_U \rightarrow \mathcal{O}_X \rightarrow 0$. Consider a polycylinder $K = K_1 \times K_2 \subset U$. By definition K is \mathcal{O}_X -privileged if and only if $h: B(K) \rightarrow B(K)$ is a split monomorphism.

Let \dot{K}_j denote the boundary of K_j , and define $\ddot{K} = \dot{K}_1 \times \dot{K}_2$ (\ddot{K} is called the Šilov Boundary of K).

Proposition 1: (a) The following conditions are equivalent:

- (i) $h: B(K) \rightarrow B(K)$ is a monomorphism.
- (i') $\exists a > 0$ such that $\|hf\| \geq a\|f\|$, $\forall f \in B(K)$.
- (ii) $X \cap \ddot{K} = \emptyset$.

(b) If $(K_1 \times K_2) \cap X = \emptyset$, then h is a split monomorphism (i.e. K is \mathcal{O}_X privileged).

Proof: (a) (i) \Leftrightarrow (i') is a well known fact from the theory of normed vector spaces.

(ii) \Rightarrow (i'). Assume $X \cap \ddot{K} = \emptyset$. If $f \in B(K)$, then it follows from the maximum principle that $\|f\| = \sup_K |f(x)| = \sup_{\ddot{K}} |f(x)|$. Since $h(x) \neq 0$

whenever $x \in \overset{\circ}{K}$, we get $a = \inf_K |h(x)| > 0$. Hence $\|hf\| = \sup_K |hf(x)| \geq \geq a \sup_K |f(x)| = a \|f\|$.

(i') \Rightarrow (ii). Suppose that $X \cap \overset{\circ}{K} \neq \emptyset$ and $x = (x_1, x_2) \in X \cap \overset{\circ}{K}$. We choose an analytic function $f_1 : U_1 \rightarrow \mathbf{C}$, where $U_1 \supset K_1$, and U_1 is open, such that $f_1(x_1) = 1$, $|f_1(z)| < 1$ if $z \in K_1$, $z \neq x_1$. Similarly we choose an analytic function $f_2 : U_2 \rightarrow \mathbf{C}$, with the same properties. Consider the function $f \in B(K) : (z_1, z_2) \rightarrow f_1(z_1)f_2(z_2)$. Since $h(x) = 0$ it follows that the sequence $\{hf^n\}$ converges pointwise to 0 in K .

Applying Dini's theorem we get $\|hf^n\| \rightarrow 0$. From the inequality $a \|f^n\| \leq \|hf^n\|$ we get $\|f^n\| \rightarrow 0$, which is a contradiction, because for every $n : f^n(x) = 1$.

(b) Use the Weierstrass preparation theorem (extended form).

Question. Does the condition (ii) imply that $h : B(K) \rightarrow B(K)$ is a split monomorphism?

IV. FLATNESS AND PRIVILEGE

§ 1. Morphisms from an analytic space into $B(K)$

Let S be an analytic space and K a polycylinder in an open set $U \subset \mathbf{C}^n$. We want to construct an \mathcal{O}_S -algebra homomorphism $\phi : \mathcal{O}_{S \times U}(S \times U) \rightarrow \mathcal{H}(S; B(K))$.

(a) Consider first $S = U' \subset \mathbf{C}^m$, U' -open. If $h \in \mathcal{O}_{U' \times U}(U' \times U)$ and $s \in U'$, $x \in K$, define $(\phi(h)(s))(x) = h(s, x)$. Using the Cauchy integral, one can show that $\phi(h)$ is analytic. On the other hand it's obvious that ϕ is an $\mathcal{O}_{U'}$ -algebra homomorphism.

(b) Let S have a special model in the polydisc Δ in \mathbf{C}^m , defined by a sheaf \mathcal{I} of ideals of \mathcal{O}_Δ , and let \mathcal{J} be generated by f_1, \dots, f_p , V -a polycylinder neighbourhood of K in U . By Cartan's theorem B for a polycylinder,

the sequence $0 \rightarrow \mathcal{I}(\Delta \times V) \rightarrow \mathcal{O}(\Delta \times V) \xrightarrow{\pi} \mathcal{O}(S \times V) \rightarrow 0$ is exact. If we denote by $\tilde{\pi}$ the projection $\mathcal{H}(\Delta, B(K)) \rightarrow \mathcal{H}(S, B(K))$, $(f_1, \dots, f_p) \cdot \mathcal{H}(\Delta, B(K)) \subset \subset \text{Ker } \tilde{\pi}$. Therefore, because π is surjection, there exists a unique

$\phi : \mathcal{O}(S \times V) \rightarrow \mathcal{H}(S, B(K))$, such that the diagram

$$\mathcal{O}(\Delta \times V) \xrightarrow{\phi} \mathcal{H}(\Delta, B(K))$$

$$\pi \downarrow \qquad \qquad \downarrow \tilde{\pi}$$

$$\mathcal{O}(S \times V) \xrightarrow{\phi} \mathcal{H}(S, B(K))$$

is commutative; ϕ is evidently an \mathcal{O}_S -algebra homomorphism.

§ 2. *The flatness and privilege theorem*

Notation

Let S be an analytic space, U an open set in \mathbb{C}^n , and $\pi : S \times U \rightarrow S$ the first projection.

If \mathcal{F} is an $\mathcal{O}_{S \times U}$ module, then for every $s \in S$ we denote by $\mathcal{F}(s)$ the \mathcal{O}_U -module $i_s^* \mathcal{F}$, where i_s is the injective morphism $x \rightarrow (s, x)$ from U into $S \times U$. If $x \in U$

$$(\mathcal{F}(s))_x \simeq \mathcal{F}_{(s, x)} / m_s \cdot \mathcal{F}_{(s, x)} \simeq \mathcal{F}_{(s, x)} \otimes_{\mathcal{O}_{S, s}} \mathbb{C}_s.$$

Theorem 1: Let \mathcal{E} be a coherent and S -flat $\mathcal{O}_{S \times U}$ -module, and K a poly-cylinder in U .

(a) When K is privileged for $\mathcal{E}(s_0)$, s_0 has a neighbourhood V such that K is $\mathcal{E}(s)$ -privileged for each $s \in V$. In other words: the set $S' = \{s \in S \mid K \text{ is } \mathcal{E}(s)\text{-privileged}\}$ is open in S .

(b) It is possible to define a Banach vector bundle over S' whose fibre at any $s \in S'$ is $B(K, \mathcal{E}(s))$.

To prove the theorem we need:

Lemma 1: Under the conditions of the theorem, we can, for every $s \in S$, find a neighbourhood W of $\{s\} \times K$ and a free resolution of finite length

$$0 \rightarrow \mathcal{L}_p \xrightarrow{d_p} \dots \xrightarrow{d_2} \mathcal{L}_1 \xrightarrow{d_1} \mathcal{L}_0 \xrightarrow{\varepsilon} \mathcal{E} \rightarrow 0 \text{ in } W.$$

Proof: Let (s, x) be a point of $S \times U$ and \mathcal{L}_*^0 a finite resolution of $\mathcal{F}(x)$ in a neighbourhood of x (there exists such one, by the theorem of syzygies). We shall show that there exists a resolution \mathcal{L}^* of \mathcal{F} in a neighbourhood of (s, x) such that $\mathcal{L}^*(s) = \mathcal{L}_*^0$; if $\mathcal{L}_i^0 = \mathcal{O}_x^{r_i}$ define

$$\mathcal{L}_i = \mathcal{O}_{S \times U}^{r_i} \text{ and } \mathcal{K}_i^0 = \text{Ker } d_i^0 : \mathcal{L}_i^0 \rightarrow \mathcal{L}_{i-1}^0.$$

We shall construct by induction (with respect to i) $d_i : \mathcal{L}_1 \rightarrow \mathcal{L}_{i-1}$ in a neighbourhood of (s, x) such that $d_i(s) = d_i^0$, and prove that $\mathcal{K}_i = \text{Ker } d_i$ is S -flat and that $\mathcal{K}_i(s) = \mathcal{K}_i^0$.

$$\begin{array}{ccc} \mathcal{L}_{i+1} & \xrightarrow{d_{i+1}} & \mathcal{K}_i \\ \downarrow & & \downarrow \\ \mathcal{L}_{i+1}^0 & \xrightarrow{d_{i+1}^0} & \mathcal{K}_i^0 \end{array}$$
 Suppose that we have constructed d_i and proved the properties for \mathcal{K}_i . We can construct $d_{i+1} : \mathcal{L}_{i+1} \rightarrow \mathcal{L}_i$ in a neighbourhood of (s, x) such that the diagram is commutative.

Nakayama's lemma shows that $\text{Im } d_{i+1} = \mathcal{K}_i$ at the point (s, x) , therefore in a neighbourhood of that point.

The exact sequence

$$0 \rightarrow \mathcal{K}_{i+1} \rightarrow \mathcal{L}_{i+1} \rightarrow \mathcal{K}_i \rightarrow 0,$$

where \mathcal{K}_i and \mathcal{L}_{i+1} are S -flat, shows that \mathcal{K}_{i+1} is S -flat, and that $\mathcal{K}_{i+1}(s) = \mathcal{K}_{i+1}^0$. The first step of the induction is analogous.

Proof of the theorem: Let $s_0 \in S$ and

$$0 \rightarrow \mathcal{L}_p \xrightarrow{d_p} \dots \xrightarrow{d_1} \mathcal{L}_0 \rightarrow \mathcal{E}|_W \rightarrow 0$$

be a free $\mathcal{O}_{S \times U}$ resolution of \mathcal{E} in a neighbourhood $W = V_1 \times V_2$ of $\{s_0\} \times K$. The sheaf \mathcal{E} is \mathcal{O}_S -flat, so for each $s \in V_1$, the sequence

$$0 \rightarrow \mathcal{L}_p \otimes_{\mathcal{O}_{S|V_1}} \mathbf{C}_s \rightarrow \dots \rightarrow \mathcal{L}_1 \otimes_{\mathcal{O}_{S|V_1}} \mathbf{C}_s \rightarrow \mathcal{L}_0 \otimes_{\mathcal{O}_{S|V_1}} \mathbf{C}_s \rightarrow \mathcal{E}|_W \otimes_{\mathcal{O}_{S|V_1}} \mathbf{C}_s \rightarrow 0$$

is exact. So the sequence

$$(A) \quad 0 \rightarrow \mathcal{L}_p(s) \xrightarrow{d_p(s)} \dots \xrightarrow{d_1(s)} \mathcal{L}_0(s) \rightarrow \mathcal{E}(s)|_{V_2} \rightarrow 0$$

is exact when $s \in V_1$. Now $\mathcal{L}_i(s) \simeq \mathcal{O}_{V_2}^{r_i}$ ($0 \leq i \leq p$) and every $d_i(s)$ induces a continuous linear map:

$B(K, \mathcal{L}_i(s)) \rightarrow B(K, \mathcal{L}_{i-1}(s))$, which we also denote by $d_i(s)$. We can consider $d_i = (d_{ijk})$ as an $r_i \times r_{i-1}$ -matrix with entries from $\mathcal{O}_{S \times U}(W)$.

By § 1 we have a \mathcal{O}_S -algebra homomorphism

$$\mathcal{O}_{S \times W} \rightarrow \mathcal{H}(S, B(K)).$$

From the matrix (d_{ijk}) we get by this homomorphism a morphism \tilde{d}_i :

$$V_0 \rightarrow \mathcal{L}(B(K)^{r_i}, B(K)^{r_{i-1}}) = \mathcal{L}(B(K, \mathcal{L}_i(s)), B(K, \mathcal{L}_{i-1}(s))).$$

(Here V_0 is some neighbourhood of s_0) such that $\tilde{d}_i(s) = d_i(s)$ for each $s \in V_0$. In other words we have a sequence of Banach vector bundle morphisms

$$(B) \quad 0 \rightarrow B(K, \mathcal{L}_p) \xrightarrow{d_p} \dots \xrightarrow{\tilde{d}_1} B(K, \mathcal{L}_0).$$

Using the fact that $\mathcal{O}_{S \times U}(S \times U) \rightarrow \mathcal{H}(S, B(K))$ is an \mathcal{O}_S -algebra homomorphism, it easily follows that (B) is complex of Banach vector bundles over S .

Now K is $\mathcal{E}(s_0)$ -privileged, thus

$$0 \rightarrow B(K, \mathcal{L}_p(s_0)) \xrightarrow{d_p(s_0)} \dots \xrightarrow{d_1(s_0)} B(K, \mathcal{L}_0(s_0))$$

is split exact, so by theorem III.1

$$0 \rightarrow B(K, \mathcal{L}_p)|_V \xrightarrow{\tilde{d}_p|_V} \dots \xrightarrow{\tilde{d}_1|_V} B(K, \mathcal{L}_0)|_V$$

is split exact for some neighbourhood V of s_0 .

Because $\tilde{d}_i(s) = d_i(s)$ and the sequence (A) is exact part (a) of the theorem follows.

(b) $B(K, \mathcal{L}_0)|_V$ splits as the direct sum of $\text{im } \tilde{d}_1$ and a bundle E_V , such that $E_{V,s} \simeq B(K, \mathcal{E}(s))$, for each $s \in V$. We must show that these bundle structures fit together globally.

Suppose therefore that V is open in S' and that

$$\begin{aligned} 0 \rightarrow \mathcal{L}_p \xrightarrow{d_p} \dots \xrightarrow{d_2} \mathcal{L}_1 \xrightarrow{d_1} \mathcal{L}_0 \rightarrow \mathcal{E}|_{V \times V_2} \rightarrow 0 \\ 0 \rightarrow \mathcal{L}'_p \xrightarrow{d'_p} \dots \xrightarrow{d'_2} \mathcal{L}'_1 \xrightarrow{d'_1} \mathcal{L}'_0 \rightarrow \mathcal{E}|_{V \times V_2} \rightarrow 0 \end{aligned}$$

are free resolutions of ξ over $V \times V_2$.

If V_1, V_2 are open polycylinders, we can find an $\mathcal{O}_{S \times U}$ -homomorphism $\phi_0 : \mathcal{L}'_0 \rightarrow \mathcal{L}_0$ such that

$$\begin{array}{ccc} & \varepsilon' & \\ & \mathcal{L}'_0 \rightarrow \mathcal{E}|_{V \times V_2} \rightarrow 0 & \\ \phi_0 \uparrow & \parallel & \\ & \varepsilon & \\ & \mathcal{L}_0 \rightarrow \mathcal{E}|_{V \times V_2} \rightarrow 0 & \end{array}$$

commutes. ϕ_0 determines a bundle morphism $\tilde{\phi}_0: B(K, \mathcal{L}_0) \rightarrow B(K, \mathcal{L}'_0)$. $B(K, \mathcal{L}_0)$ (resp. $B(K, \mathcal{L}'_0)$) splits as $(\text{im } \tilde{d}_1) \otimes E_V$ [Resp. $(\text{im } \tilde{d}'_1) \otimes E'_V$].

Let p' be the projection morphism: $B(K, \mathcal{L}_0) \rightarrow E'_V$ with kernel $\text{im } \tilde{d}'_1$, and put $\tilde{\phi} = p' \circ \phi_0|_{E_V}$.

The commutative diagram

$$\begin{array}{ccc}
 B(K, \mathcal{L}_0(s)) & \xrightarrow{\tilde{\phi}_0} & B(K, \mathcal{L}'_0(s)) \\
 \varepsilon \downarrow & \swarrow & \downarrow \varepsilon' \\
 & E_{V,s} & \xrightarrow{\tilde{\phi}} E'_{V,s} \\
 & \swarrow \varepsilon \simeq \alpha \circ \varepsilon & \searrow \varepsilon' \simeq \alpha' \circ \varepsilon' \\
 B(K, \mathcal{E}(s)) & \xleftarrow{\text{id}} & B(K, \mathcal{E}'(s))
 \end{array}$$

and the open mapping theorem shows that $\tilde{\phi}(s)$ is an isomorphism of Banach spaces for each $s \in V$, so $\tilde{\phi}: E_V \rightarrow E'_V$ is a bundle isomorphism. We also notice that $\tilde{\phi}$ depends only on the choice of splittings in $B(K, \mathcal{L}_0)$ and $B(K, \mathcal{L}'_0)$, and not on the choice of $\tilde{\phi}_0$. This ends the proof of the theorem.

Remark: Consider the general situation where X and S are analytic spaces, and $\pi: X \rightarrow S$ is a morphism, \mathcal{E} an \mathcal{O}_X -module. To study the local dependence of \mathcal{E} on S , one can imbed an open set X' in X in the open set $U \subset \mathbb{C}^n$. The morphism $\phi: X' \rightarrow U, \pi: X' \rightarrow S$ determine the imbedding $\pi \times \phi: X' \rightarrow S \times U$ such that the diagram commutes. \mathcal{E} can be extended by zero into a sheaf \mathcal{E}' over $U \times S$. Obviously this sheaf \mathcal{E}' is S -flat iff \mathcal{E} is S -flat.

Therefore theorem 1 makes clear also this general situation.

Corollary: If $\pi: X \rightarrow S$ is a morphism and \mathcal{E} a coherent \mathcal{O}_X -module. Then $\pi|_{\text{Supp}(\mathcal{E})}$ is an open map.

Proof: Suppose as above that X is imbedded in $S \times U$, and \mathcal{E} is extended by zero to $S \times U$. Let $x_0 \in \text{Supp } \mathcal{E}$, and V be a neighbourhood of x_0 in $S \times U$. Let $s_0 = \pi(x_0)$ and choose an $\mathcal{E}(s_0)$ -privileged polycylinder K in U , such that $\{s_0\} \times K \subset V$, over some neighbourhood W of s_0 . We have the Banach bundle $B(K, \mathcal{E}|_{\pi^{-1}(W)})$, whose fiber over s is $B(K, \mathcal{E}(s))$. Since $x_0 \in \text{Supp } \mathcal{E}(s_0)$ and K is a neighbourhood of x_0 , $B(K; \mathcal{E}(s_0)) \neq 0$. As all the fibers are isomorphic, then for all $s \in U$, $B(K; \mathcal{E}(s)) \neq 0$ and therefore $\{s\} \times K \cap \text{Supp } \mathcal{E} \neq \emptyset$, and $s \in \pi(\text{Supp } \mathcal{E})$. This proves that π is open.

REFERENCES

- GUNNING, R. C. and H. ROSSI, *Analytic functions of several complex variables*. Prentice-Hall, Inc. Englewood Cliffs, N.J. 1965.
- DOUADY, A., Le problème des modules pour les sous-espaces analytiques compacts d'un espace analytique donné. *Annales de Fourier, Université de Grenoble*, tome XVI, 1966, pp. 1-95.
- MALGRANGE, B., *Lecture Notes*. Analytic spaces (see this volume, p. 1-28)