

4. NON-EUCLIDEAN MOTIONS

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **24 (1978)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

4. NON-EUCLIDEAN MOTIONS

The euclidean case was dealt with in [3]. In the present paper we undertake a more detailed study of the hyperbolic case. The unit ball in $\mathbf{R}^n$ is denoted by B , and G is the full group of Möbius transformations mapping B on itself. The Poincaré metric $ds = (1 - |x|^2)^{-1} |dx|$ and the non-euclidean volume element $\rho dx = (1 - |x|^2)^{-n} dx$ are invariant under G .

For $A \in G$ we prefer to denote the Jacobian by $A'(x)$ rather than $DA(x)$. We use $|A'(x)|$ for the linear rate of change, the same in all directions. This notation has the advantage of leading to formulas which are easily recognizable generalizations of the familiar formulas for $n = 2$ in complex notation. $|A'(x)|$ is also the square norm of the matrix $A'(x)$, and $|\det A'(x)| = |A'(x)|^n$.

Reflection in the unit sphere is denoted by $x^* = x/|x|^2$. Its Jacobian is $Dx^* = |x|^{-2} (1_n - 2Q(x))$ with $Q(x)_{ij} = x_i x_j / |x|^2$; note that $(1_n - 2Q(x))^2 = 1_n$.

For every $y \in B$ there is a unique $T_y \in G$ such that $T_y y = 0$ and $T'_y(y) = |T'_y(y)| \cdot 1_n$. The most general $A \in G$ is of the form $A = UT_y$ with $y = A^{-1}(0)$ and $U \in O(n)$.

For $n = 2$, in complex notation,

$$T_y x = \frac{x - y}{1 - \bar{y}x}$$

$$T'_y(x) = \frac{1 - |y|^2}{(1 - \bar{y}x)^2}.$$

The first formula can be rewritten as

$$T_y x = \frac{(x - y)(1 - |y|^2) - |x - y|^2 y}{|y|^2 |x - y^*|^2}.$$

In this form it makes sense for arbitrary n and is in fact the correct formula. The denominator $|y|^2 |x - y^*|^2$ corresponds to $|1 - \bar{y}x|^2$, and it is equal to $1 - 2(xy) + |x|^2 |y|^2$, where (xy) is the inner product. To emphasize the symmetry we shall use the notation $|y| |x - y^*| = |x| |y - x^*| = [x, y]$.

The expression for $T'_y(x)$ is

$$T'_y(x) = \frac{1 - |y|^2}{[x, y]^2} \Delta(x, y)$$

with

$$\Delta(x, y) = (1 - 2Q(y))(1 - 2Q(x - y^*)) = (1 - 2Q(y - x^*))(1 - 2Q(x)).$$

Observe that $\Delta(x, y) = {}^t\Delta(y, x)$ and $\Delta(x, y)^2 = 1_n$ so that $\Delta(x, y) \in O(n)$. The matrix $\Delta(x, y)$ generalizes the angle $\arg(1 - \bar{x}y)/(1 - \bar{y}x)$.

It is useful to note that $|Ax - Ay|^2 = |A'(x)| |A'(y)| |x - y|^2$ for any Möbius transformation A , and $[Ax, Ay]^2 = |A'(x)| |A'(y)| [x, y]^2$ if $A \in G$. There is an important relation between T_yx and T_xy expressed by

$$(4) \quad T_yx = -\Delta(x, y) T_xy.$$

We refer to [2, 3, 4, 5] for the elementary proofs of these formulas.

5. FUNDAMENTAL SOLUTIONS

A continuous mapping $f: B \rightarrow \mathbf{R}^n$ will be called a *deformation*. In this paper we shall assume, mainly for simplicity, that f is continuous on the boundary $S(1)$, and that $x \cdot f(x) = 0$ on $S(1)$; this means that f maps B on itself when regarded as an infinitesimal mapping.

A deformation is *trivial* if $Sf = 0$. There are very few trivial deformations: a complete list is given in [3].

It is customary to say that f is a *quasiconformal* deformation if $\|Sf\| \in L^\infty(B)$; here $\|Sf\|$ is the function whose value at x is the square norm of the matrix $Sf(x)$. More generally, we shall also consider functions with $\|Sf\| \in L^p(B)$; we abbreviate to $Sf \in L^p$, and we denote the L^p -norm of the square norm by $\|Sf\|_p$. The same convention will prevail for all matrix-valued functions.

We shall say that f is *harmonic* if $S^* \rho Sf = 0$, $\rho = (1 - |x|^2)^{-n}$. Because of the invariance, if f is harmonic and $A \in G$, then A^*f is also harmonic. Harmonicity in this sense is not the same as requiring the components to be harmonic with respect to the Poincaré metric.

There are n linearly independent solutions of the equation $S^* \gamma = 0$ which are homogeneous of degree $1 - n$. We denote them by $\gamma_{\dots, k}$, $k = 1, \dots, n$, the elements being

$$\gamma_{ij, k}(x) = |x|^{-n} (\delta_{ik} x_j + \delta_{jk} x_i - \delta_{ij} x_k) + (n - 2) |x|^{-n-2} x_i x_j x_k.$$

There is a unique vector-valued function $g_{\dots, k}(x)$ with components $g_{ik}(x)$ such that $g_{\dots, k}(x) = 0$ for $|x| = 1$ and $\rho Sg_{\dots, k} = \gamma_{\dots, k}$ so that