

3. CO-DIRECTIONS IN PROJECTIVE SPACE

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **25 (1979)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

is one-to-one on an open neighborhood U of x_0 in G/P and $(\exp X) \cdot x_0$ is identifiable as the point $(x_1, \dots, x_n)$ and the incident hyperplane

$$x'_n - x_n = p_1(x'_1 - x_1) + \dots + p_{n-1}(x'_{n-1} - x_{n-1}).$$

Now, $(\exp X) \cdot x_0 \rightarrow (\exp X) \cdot b_0$ is a section of the bundle G/P_1 over U and, via this section, the form ω on G/P_1 pulls down to

$$\omega_0((\exp X)^{-1} d(\exp X))$$

which, when expressed in terms of $x_1, \dots, x_n, p_1, \dots, p_{n-1}$, will be identified with

$$dx_n - p_1 dx_1 - \dots - p_{n-1} dx_{n-1}$$

up to a constant multiple $a \neq 0$. For this latter calculation we will use

$$\begin{aligned} (\exp X)^{-1} d(\exp X) &= \frac{1 - e^{-ad X}}{ad X} (dX) \\ &= dX - \frac{1}{2} [X, dX] + \frac{1}{6} [X, [X, dX]] - \dots \end{aligned}$$

[4, (10.2)], a series which is finite since $\mathfrak{m}$ is nilpotent. In fact, our choice of X will make the series for $\exp X$ themselves finite. The constant $a \neq 0$ could be made unity by using instead the section $(\exp X) \cdot x_0 \rightarrow (\exp X)g^{-1} \cdot b_0$, where g in P is chosen so that $\chi(g) = a$. This amounts to following the original section by R_a^{-1} in the bundle.

3. CO-DIRECTIONS IN PROJECTIVE SPACE

The contact structure on the $(2n-1)$ -dimensional space of co-directions in complex projective space P^n , described in 2.5, is obtained when the construction of 2.10 is carried out for the simple complex Lie algebra of type A_n , $n \geq 1$.

3.1 Let $\mathfrak{g} = \mathfrak{sl}(n+1; \mathbb{C})$, complex $(n+1)$ by $(n+1)$ matrices of trace zero. For Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$ take the diagonal matrices of $\mathfrak{g}$. Let δ_i , $i = 0, 1, \dots, n$ be the linear function on $\mathfrak{h}$ which assigns to $H = \text{diag}(h_1, \dots, h_n)$ in $\mathfrak{h}$ the i^{th} diagonal element: $\delta_i(H) = h_i$. The roots of $\mathfrak{g}$ with respect to $\mathfrak{h}$ are

$$\begin{aligned} \delta_i - \delta_j \quad i, j = 0, 1, \dots, n \\ \text{and } i \neq j \end{aligned}$$

and the root vector E_α corresponding to the root α is

$$E_{\delta_i - \delta_j} = E_{ij},$$

the matrix with 1 in the i^{th} row and j^{th} column and 0s elsewhere [4, (16.2)]. A system of simple roots is

$$\delta_0 - \delta_1, \delta_1 - \delta_2, \dots, \delta_{n-1} - \delta_n,$$

for which the maximal root is

$$\rho = (\delta_0 - \delta_1) + (\delta_1 - \delta_2) + \dots + (\delta_{n-1} - \delta_n) = \delta_0 - \delta_n$$

[4, App., Table E]. The Killing form of $\mathfrak{g}$ is $\langle X, Y \rangle = 2(n+1) \operatorname{tr}(XY)$, but we replace this with $\langle X, Y \rangle = \operatorname{tr}(XY)$ for convenience. Then the H_α in $\mathfrak{h}$ are given by

$$H_{\delta_i - \delta_j} = \operatorname{diag}(0, \dots, 0, 1, 0, \dots, 0, -1, 0, \dots, 0)$$

with 1 and -1 in the i^{th} and j^{th} entry, respectively. Especially,

$$H_\rho = \operatorname{diag}(1, 0, \dots, 0, -1).$$

We have

$$\langle H_\rho, H_{\delta_i - \delta_j} \rangle \begin{cases} < 0 & j = 0 \text{ or } i = n \\ \geq 0 & \text{otherwise} \end{cases},$$

so that $\mathfrak{p}$ in (i) of 2.9 consists of matrices of the form

$$\begin{bmatrix} * & * & \cdots & * \\ & \vdots & & \vdots \\ 0 & * & \cdots & * \\ & \vdots & & \vdots \\ 0 & \cdots & 0 & * \end{bmatrix}$$

of trace zero, where the starred entries are arbitrary.

3.2 The connected centerless simple group $G = PSL(n+1; \mathbf{C}) = SL(n+1; \mathbf{C})/\{\text{center}\}$ is transitive on the space consisting of points x and incident hyperplanes $u, ux = 0$, in P^n , as in 2.5. The isotropy subgroup P of the incident point and hyperplane

$$x_0 = {}^t(1, 0, \dots, 0), \quad u_0 = (0, \dots, 0, 1)$$

has exactly $\mathfrak{p}$ for its Lie algebra. Hence, the homogeneous contact manifold which the construction of 2.10 gives is

$$\begin{aligned} G/P &= \text{space of incident points and hyperplanes in } P^n \\ &= \text{space of co-directions in complex } P^n. \end{aligned}$$

3.3 Let $\mathfrak{m}$ be the $(2n-1)$ -dimensional supplement to $\mathfrak{p}$ in $\mathfrak{g}$ consisting of matrices of the form

$$\begin{bmatrix} 0 & & & \\ & * & & 0 \\ & \vdots & & \\ & * & \cdots & * & 0 \end{bmatrix};$$

cf. 2.12. The product of any two matrices of $\mathfrak{m}$ has a nonzero entry only in the n^{th} row and 0^{th} column; the product of any three is zero. Set

$$X = \begin{bmatrix} 0 & & & \\ & x_1 & & 0 \\ & \vdots & & \\ & x_{n-1} & & \\ x_n - \frac{1}{2} \sum p_i x_i & p_1 \cdots p_{n-1} & 0 \end{bmatrix},$$

where the summation is over $i = 1, 2, \dots, n-1$. X is in $\mathfrak{m}$ and

$$\exp X = 1_{n+1} + X + \frac{1}{2} X^2 =$$

$$\begin{bmatrix} 1 & & & \\ & x_1 & & 0 \\ & \vdots & & \\ & x_{n-1} & & 0 \\ x_n & p_1 \cdots p_{n-1} & 1 \end{bmatrix},$$

$$(\exp X)^{-1} = 1_{n+1} - X + \frac{1}{2} X^2 = \begin{bmatrix} 1 & & & & \\ & -x_1 & & & 0 \\ & \vdots & & & \\ & & -x_{n-1} & & 0 \\ -x_n + \sum p_i x_i & -p_1 & \dots & -p_{n-1} & 1 \end{bmatrix}.$$

The point

$$x = (\exp X) \cdot x_0 = {}^t(1, x_1, \dots, x_n)$$

is incident with the hyperplane

$$u = u_0 \cdot (\exp X)^{-1} = (-x_n + \sum p_i x_i, -p_1, \dots, -p_{n-1}, 1),$$

and the hyperplane $ux' = 0$, $x' = {}^t(1, x'_1, \dots, x'_n)$, is

$$x'_n - x_n = p_1 (x'_1 - x_1) + \dots + p_{n-1} (x'_{n-1} - x_{n-1}).$$

Thus, this choice of X establishes the classically identifiable coordinates $x_1, \dots, x_n, p_1, \dots, p_{n-1}$ on G/P .

3.8 From $\rho = \delta_0 - \delta_n$, we have $W = E_\rho = E_{0n}$ in (iii) of 2.9 and $\omega_0(X) = \langle W, X \rangle$ is the $n0$ -entry of X . The form ω on G/P is obtained as $\omega = \omega_0((\exp X)^{-1} d(\exp X))$ with

$$(\exp X)^{-1} d(\exp X) = dX - \frac{1}{2} [X, dX]$$

as in 2.12. For X as in 3.3, the only nonzero entry in $[X, dX]$ is the $n0^{th}$ and it is $\sum p_i dx_i = -\sum x_i dp_i$. Hence

$$(\exp X)^{-1} d(\exp X) = \begin{bmatrix} 0 & & & & \\ & dx_1 & & & \\ & \vdots & & & \\ & & dx_{n-1} & & \\ dx_n - \sum p_i dx_i & dp_1 & \dots & dp_{n-1} & 0 \end{bmatrix},$$

and the $n0$ -entry is

$$\omega = dx_n - p_1 dx_1 - \dots - p_{n-1} dx_{n-1}.$$

This identifies the contact structure with the classical one as in 2.12.

3.5 The real contact structure on the $(2n-1)$ -dimensional space of co-directions in real projective space P^n is described by viewing all quantities in the foregoing discussion as being real. Especially, G_0 of 2.11 is the connected centerless group $PSL(n+1; \mathbf{R})$ consisting of real contact automorphisms.

4. HIGHER SPHERE GEOMETRY

4.1 In complex Euclidean space E^n , the equation

$$x_1'^2 + \dots + x_n'^2 - 2a_1 x_1' - \dots - 2a_n x_n' + C = 0$$

describes a sphere with center $(a_1, \dots, a_n)$ and complex radius r given by

$$r^2 = a_1^2 + \dots + a_n^2 - C.$$

When $r \neq 0$, the two choices of sign for r is said to give two "orientations" to the sphere. Thus, the $n+2$ coordinates $a_1, \dots, a_n, r, C$, which are related by

$$a_1^2 + \dots + a_n^2 - r^2 - C = 0,$$

describe the space of oriented spheres in E^n [6, §25].

Introduce homogeneous coordinates by

$$a_i = \frac{\alpha_i}{v}, \quad r = \frac{\lambda}{v}, \quad C = \frac{\mu}{v},$$

$i = 1, 2, \dots, n$. Then the oriented spheres of E^n correspond to certain points of the quadric Ψ^{n+1} in P^{n+2} described by

$$\alpha_1^2 + \dots + \alpha_n^2 - \lambda^2 - \mu v = 0.$$

The sphere corresponding to the point $(\alpha_1, \dots, \alpha_n, \lambda, \mu, v)$ of Ψ^{n+1} is

$$v(x_1'^2 + \dots + x_n'^2) - 2\alpha_1 x_1' - \dots - 2\alpha_n x_n' + \mu = 0.$$

Ordinary spheres have finite nonzero radius r , so $v \neq 0$. For $v = 0$, we obtain oriented hyperplanes. For $\lambda = 0$, we obtain point spheres or hyperplanes with isotropic hyperplane coordinate vector; these carry no