

Objektyp: **ReferenceList**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **26 (1980)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **24.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

$$(25.13) \quad 5i \sum_{k=0}^{\infty} \frac{1}{(2k+1)^2 + 10(2k+1)i - 50} = \frac{4689}{11,890},$$

or equivalently that

$$(25.14) \quad 50 \sum_{k=0}^{\infty} \frac{2k+1}{(2k+1)^4 + 2500} = \frac{4689}{11,890},$$

since (25.12) obviously implies that the imaginary part of the left side of (25.13) is zero. To show (25.14), write

$$\begin{aligned} & 50 \sum_{k=0}^{\infty} \frac{2k+1}{(2k+1)^4 + 2500} \\ &= \frac{5}{2} \sum_{k=0}^{\infty} \left\{ \frac{1}{(2k+1)^2 - 10(2k+1) + 50} - \frac{1}{(2k+1)^2 + 10(2k+1) + 50} \right\} \\ &= \frac{5}{2} \sum_{k=0}^{\infty} \left\{ \frac{1}{(2k+1)^2 - 10(2k+1) + 50} \right. \\ & \quad \left. - \sum_{k=5}^{\infty} \frac{1}{(2k+1-10)^2 + 10(2k+1-10) + 50} \right\} \\ &= \frac{5}{2} \sum_{k=0}^4 \frac{1}{(2k+1)^2 - 10(2k+1) + 50} = \frac{4689}{11,890}, \end{aligned}$$

and the proof of (25.14), and hence (25.11), is complete.

REFERENCES

- [1] ABRAMOWITZ, M. and I. A. STEGUN, editors. *Handbook of mathematical functions*. Dover, New York, 1965.
- [2] AIYAR, M. V. On some elliptic function series. *J. Indian Math. Soc.* 18 (1929/30), pp. 46-50.
- [3] AIYAR, S. N. Some theorems in summation. *J. Indian Math. Soc.* 5 (1913), pp. 183-186.
- [4] APOSTOL, T. M. Generalized Dedekind sums and transformation formulae of certain Lambert series. *Duke Math. J.* 17 (1950), pp. 147-157.
- [5] ——— On the Lerch zeta function. *Pacific J. Math.* 1 (1951), pp. 161-167.
- [6] AYOUB, R. *An introduction to the analytic theory of numbers*. American Math. Soc., Providence, 1963.
- [7] BERNDT, B. C. Ramanujan's formula for $\zeta(2n+1)$. *Professor Srinivasa Ramanujan commemoration volume*. Jupiter Press, Madras, 1974, pp. 2-9.
- [8] ——— Periodic Bernoulli numbers, summation formulas and applications. *Theory and application of special functions*. R. A. Askey, editor. Academic Press, New York, 1975, pp. 143-189.

- [9] BERNDT, B. C. Character analogues of the Poisson and Euler-Maclaurin summation formulas with applications. *J. Number Theory* 7 (1975), pp. 413-445.
- [10] — On Eisenstein series with characters and the values of Dirichlet L -functions. *Acta Arith.* 28 (1975), pp. 299-320.
- [11] — Dedekind sums and a paper of G. H. Hardy. *J. London Math. Soc. (2)* 13 (1976), pp. 129-137.
- [12] — Modular transformations and generalizations of several formulae of Ramanujan. *Rocky Mountain J. Math.* 7 (1977), pp. 147-189.
- [13] — Ramanujan's Notebooks. *Math. Mag.* 51 (1978), pp. 147-164.
- [14] — Analytic Eisenstein series, theta-functions, and series relations in the spirit of Ramanujan. *J. Reine Angew. Math.* 303/304 (1978), pp. 332-365.
- [15] BODENDIEK, R. *Über verschiedene Methoden zur Bestimmung der Transformationsformeln der achten Wurzeln der Integralmoduln $k^2(\tau)$ und $k'^2(\tau)$, ihrer Logarithmen sowie gewisser Lambertscher Reihen bei beliebigen Modulusubstitutionen.* Dissertation der Universität Köln, 1968.
- [16] BODENDIEK, R. und U. HALBRITTER. Über die Transformationsformel von $\log \eta(\tau)$ und gewisser Lambertscher Reihen. *Abh. Math. Sem. Univ. Hamburg* 38 (1972), pp. 147-167.
- [17] CAUCHY, A. *Œuvres*. Série II, t. VII, Paris, 1889.
- [18] CHANDRASEKHARAN, K. and R. NARASIMHAN. Hecke's functional equation and arithmetical identities. *Ann. of Math.* 74 (1961), pp. 1-23.
- [19] CHOWLA, S. D. Some infinite series, definite integrals and asymptotic expansions. *J. Indian Math. Soc.* 17 (1927/28), pp. 261-288.
- [20] GLAESKE, H.-J. Eine einheitliche Herleitung einer gewissen Klasse von Transformationsformeln der analytischen Zahlentheorie (I). *Acta Arith.* 20 (1972), pp. 133-145.
- [21] — Eine einheitliche Herleitung einer gewissen Klasse von Transformationsformeln der analytischen Zahlentheorie (II). *Acta Arith.* 20 (1972), pp. 253-265.
- [22] GLAISHER, J. W. L. On the series which represent the twelve elliptic and the four zeta functions. *Mess. Math.* 18 (1889), pp. 1-84.
- [23] GRADSHTEYN, I. S. and I. M. RYZHIK. *Table of integrals, series, and products.* Academic Press, New York, 1965.
- [24] GROSSWALD, E. Die Werte der Riemannschen Zeta-funktion an ungeraden Argumentstellen. *Nachr. Akad. Wiss. Göttingen* (1970), pp. 9-13.
- [25] — Comments on some formulae of Ramanujan. *Acta Arith.* 21 (1972), pp. 25-34.
- [26] GUINAND, A. P. Functional equations and self-reciprocal functions connected with Lambert series. *Quart. J. Math. Oxford ser. 15* (1944), pp. 11-23.
- [27] — Some rapidly convergent series for the Riemann ξ -function. *Quart. J. Math. Oxford ser. (2)* 6 (1955), pp. 156-160.
- [28] HARDY, G. H. A chapter from Ramanujan's notebook. *Proc. Cambridge Philos. Soc.* 21 (1923), pp. 492-503.
- [29] — A formula of Ramanujan. *J. London Math. Soc.* 3 (1928), pp. 238-240.
- [30] — Ramanujan and the theory of Fourier transforms. *Quart. J. Math. Oxford ser. 8* (1937), pp. 245-254.
- [31] — *Ramanujan*. Chelsea, New York, 1959.
- [32] — *Collected papers, vol. IV*. Clarendon Press, Oxford, 1969.
- [33] HENRICI, P. *Applied and computational complex analysis, vol. 2*. John Wiley & Sons, New York, 1977.
- [34] HURWITZ, A. Grundlagen einer independenten Theorie der elliptischen Modul-funktionen und Theorie der Multiplikator-Gleichungen erster Stufe. *Math. Ann.* 18 (1881), pp. 528-592.
- [35] — *Mathematische Werke, Band I*. Emil Birkhäuser & C^{ie}, Basel, 1932.

- [36] SHÔ ISEKI. The transformation formula for the Dedekind modular function and related functional equations. *Duke Math. J.* 24 (1957), pp. 653-662.
- [37] KATAYAMA, K. On Ramanujan's formula for values of Riemann zeta-function at positive odd integers. *Acta Arith.* 22 (1973), pp. 149-155.
- [38] ——— Ramanujan's formulas for L -functions. *J. Math. Soc. Japan* 26 (1974), pp. 234-240.
- [39] ——— Zeta-functions, Lambert series and arithmetic functions analogous to Ramanujan's τ -function. I. *J. Reine Angew. Math.* 268/269 (1974), pp. 251-270.
- [40] ——— Zeta-functions, Lambert series and arithmetic functions analogous to Ramanujan's τ -function. II. *J. Reine Angew. Math.* 282 (1976), pp. 11-34.
- [41] KNOPP, K. *Theory and application of infinite series*. Blackie & Son, Glasgow, 1951.
- [42] KOSHIAKOV, N. S. Application of the theory of sumformulae to the investigation of a class of one-valued analytical functions in the theory of numbers. *Mess. Math.* 58 (1929), pp. 1-23.
- [43] KRISHNAMAGHARI, C. Certain definite integrals and series. *J. Indian Math. Soc.* 12 (1920), pp. 14-31.
- [44] LAGRANGE, J. Une formule sommatoire et ses applications. *Bull. Sci. Math.* (2) 84 (1960), pp. 105-110.
- [45] LERCH, M. Sur la fonction $\zeta(s)$ pour valeurs impaires de l'argument. *J. Sci. Math. Astron. pub. pelo D^r F. Gomes Teixeira, Coimbra* 14 (1901), pp. 65-69.
- [46] LEWITTES, J. Analytic continuation of the series $\sum (m+nz)^{-s}$. *Trans. Amer. Math. Soc.* 159 (1971), pp. 505-509.
- [47] ——— Analytic continuation of Eisenstein series. *Trans. Amer. Soc.* 171 (1972), pp. 469-490.
- [48] LING, C.-B. Generalization of certain summations due to Ramanujan. *SIAM J. Math. Anal.* 9 (1978), pp. 34-48.
- [49] MALURKAR, S. L. On the application of Herr Mellin's integrals to some series. *J. Indian Math. Soc.* 16 (1925/26), pp. 130-138.
- [50] MELLIN, H. Eine Formel für den Logarithmus transcender Funktionen von endlichem Geschlecht. *Acta Soc. Sci. Fennicae* 29 (1902), 49 pp.
- [51] MIKOLÁS, M. Über gewisse Lambertsche Reihen, I: Verallgemeinerung der Modulfunktion $\eta(\tau)$ und ihrer Dedekindschen Transformationsformel. *Math. Z.* 68 (1957), pp. 100-110.
- [52] NANJUNDIAH, T. S. Certain summations due to Ramanujan, and their generalisations. *Proc. Indian Acad. Sci., Sect. A* 34 (1951), pp. 215-228.
- [53] PHILLIPS, E. G. Note on summation of series. *J. London Math. Soc.* 4 (1929), pp. 114-116.
- [54] PREECE, C. T. Theorems stated by Ramanujan (XIII). *J. London Math. Soc.* 6 (1931), pp. 95-99.
- [55] RAMANUJAN, S. Question 358. *J. Indian Math. Soc.* 4 (1912), p. 78.
- [56] ——— Question 387. *J. Indian Math. Soc.* 4 (1912), p. 120.
- [57] ——— Modular equations and approximations to π . *Quart. J. Math.* 45 (1914), pp. 350-372.
- [58] ——— Some definite integrals. *Mess. Math.* 44 (1915), pp. 10-18.
- [59] ——— Some definite integrals connected with Gauss's sums. *Mess. Math.* 44 (1915), pp. 75-85.
- [60] ——— On certain trigonometrical sums and their applications in the theory of numbers. *Trans. Cambridge Philos. Soc.* 22 (1918), pp. 259-276.
- [61] ——— *Collected papers*. Chelsea, New York, 1962.
- [62] ——— *Notebooks of Srinivasa Ramanujan (2 volumes)*. Tata Institute of Fundamental Research, Bombay, 1957.

- [63] RAO, M. B. and M. V. AYYAR. On some infinite series and products, Part I. *J. Indian Math. Soc.* 15 (1923/24), pp. 150-162.
- [64] RAO, M. B. and M. V. AYYAR. On some infinite products and series, Part II. *J. Indian Math. Soc.* 15 (1923/24), pp. 233-247.
- [65] RIESEL, H. Some series related to infinite series given by Ramanujan. *BIT* 13 (1973), pp. 97-113.
- [66] SANDHAM, H. F. Three summations due to Ramanujan. *Quart. J. Math. Oxford ser. (2) 1* (1950), pp. 238-240.
- [67] ——— Some infinite series. *Proc. Amer. Math. Soc.* 5 (1954), pp. 430-436.
- [68] SAYER, F. P. The sums of certain series containing hyperbolic functions. *Fibonacci Quart.* 14 (1976), pp. 215-223.
- [69] SCHLÖMILCH, O. Über einige unendliche Reihen. *Ber. Verh. K. Sächs. Gesell. Wiss. Leipzig* 29 (1877), pp. 101-105.
- [70] ——— *Compendium der höheren Analysis, Zweiter Band, 4th ed.* Friedrich Vieweg und Sohn, Braunschweig, 1895.
- [71] SMART, J. R. On the values of the Epstein zeta function. *Glasgow Math. J.* 14 (1973), pp. 1-12.
- [72] TITCHMARSH, E. C. *Theory of Fourier integrals, second ed.* Clarendon Press, Oxford, 1948.
- [73] WATSON, G. N. Theorems stated by Ramanujan II. *J. London Math. Soc.* 3 (1928), pp. 216-225.
- [74] ——— Ramanujan's note books. *J. London Math. Soc.* 6 (1931), pp. 137-153.
- [75] WHITTAKER, E. T. and G. N. WATSON. *A course of modern analysis, 4th ed.* University Press, Cambridge, 1962.
- [76] ZUCKER, I. J. The summation of series of hyperbolic functions. *SIAM J. Math. Anal.* 10 (1979), pp. 192-206.

(Reçu le 5 janvier 1979)

Bruce C. Berndt

Department of Mathematics
University of Illinois
Urbana, Illinois 61801
U.S.A.

vide-leer-empty