

3.2. The case $SU(1, 1)$

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **28 (1982)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

and

$$\tau(k)v = \tau_0(k)v.$$

(b) \Rightarrow (a): Let $0 \neq v \in \mathcal{H}(\tau_0)$. Let \mathcal{H}_1 be the τ_0 -invariant subspace of $\mathcal{H}(\tau_0)$ generated by v . Then $\phi_\gamma \in \mathcal{H}_1$ for some $\gamma \in \mathcal{M}(\tau_0)$. Now, for some $g \in G$,

$$(\tau_0(g)\phi_\gamma, \phi_\delta) = \tau_{0, \delta, \gamma}(g) = \tau_{\delta, \gamma}(g) \neq 0,$$

so $\tau_0(g)\phi_\gamma$ and ϕ_δ are in \mathcal{H}_1 . For each $\beta \in \mathcal{M}(\tau_0)$ we have $(\tau_0(g)\phi_\delta, \phi_\beta) = \tau_{\beta\delta}(g) \neq 0$ for some $g \in G$. Thus $\phi_\beta \in \mathcal{H}_1$ for all $\beta \in \mathcal{M}(\tau_0)$, so $\mathcal{H}_1 = \mathcal{H}(\tau_0)$.

(a) \Rightarrow (c): Suppose $\tau_{\gamma\delta} = 0$ for some $\gamma, \delta \in \mathcal{M}(\tau_0)$. Then, for all $g \in G$, $(\tau_0(g)\phi_\delta, \phi_\gamma) = 0$. Hence, the τ_0 -invariant subspace of $\mathcal{H}(\tau_0)$ generated by ϕ_δ is orthogonal to ϕ_γ , so τ_0 is not irreducible.

(c) \Rightarrow (b): Clear. □

Let τ be K -multiplicity free, K being compact abelian. Define a relation $<$ on $\mathcal{M}(\tau)$ by: $\gamma < \delta$ iff $\tau_{\gamma, \delta} \neq 0$. Then $\gamma < \delta$ iff ϕ_γ is in the τ -invariant subspace of $\mathcal{H}(\tau)$ generated by ϕ_δ . It follows that

$$\beta < \gamma \text{ and } \gamma < \delta \Rightarrow \beta < \delta$$

Define a relation \sim on $\mathcal{M}(\tau)$ by: $\gamma \sim \delta$ iff $\tau_{\gamma, \delta} \neq 0 \neq \tau_{\delta, \gamma}$. It follows that \sim is an equivalence relation on $\mathcal{M}(\tau)$ and that, if $\tau_{\gamma, \delta} \neq 0, \alpha \sim \gamma, \beta \sim \delta$ then $\tau_{\alpha, \beta} \neq 0$. It follows that, for a given equivalence set, we can partition $\mathcal{M}(\tau)$ into three parts, the first part being the equivalence set, such that the corresponding 3×3 block matrix for $(\tau_{\gamma\delta}(g))$ takes the form (3.3). In view of Theorem 3.2 this proves:

THEOREM 3.3. *Let G be a lcsc. group with compact abelian subgroup K and let τ be a K -multiplicity free representation of G . Then there is a unique orthogonal decomposition of $\mathcal{H}(\tau)$ into subspaces $\mathcal{H}(\tau_i)$, where the τ_i 's are precisely the irreducible subquotient representations of τ .*

3.2. THE CASE $SU(1, 1)$

For $\lambda \in \mathbb{C}$, $\xi = 0$ or $\frac{1}{2}$, the representation $\pi_{\xi, \lambda}$ of $G = SU(1, 1)$ on $L^2_\xi(K)$ (cf. (2.8)) is K -multiplicity free with K -content given by (2.13). By inspecting (2.29) for small but nonzero t and by using (2.24) it follows that

$$(3.4) \quad \pi_{\xi, \lambda, m, n} \neq 0 \Leftrightarrow \pi_{\xi, \lambda, m, n}|_A \neq 0 \Leftrightarrow c_{\xi, \lambda, m, n} \neq 0,$$

where $c_{\xi, \lambda, m, n}$ is given by (2.30). Combination of (3.4) with Theorems 3.2 and 3.3 yields:

THEOREM 3.4. *Depending on ξ and λ , the representation $\pi_{\xi, \lambda}$ of $SU(1, 1)$ has the following irreducible subquotient representations:*

(a) $\lambda + \xi \notin \mathbf{Z} + \frac{1}{2}$:

$\pi_{\xi, \lambda}$ is irreducible itself.

(b) $\lambda = 0, \xi = \frac{1}{2}$:

$\pi_{1/2, 0}^+$ on $\text{Cl Span } \{\phi_{1/2}, \phi_{3/2}, \dots\}$,

$\pi_{1/2, 0}^-$ on $\text{Cl Span } \{\dots, \phi_{-3/2}, \phi_{-1/2}\}$.

These are also subrepresentations.

(c) $\lambda + \xi \in \mathbf{Z} + \frac{1}{2}, \lambda > 0$:

$\pi_{\xi, \lambda}^+$ on $\text{Cl Span } \{\phi_{\lambda+1/2}, \phi_{\lambda+3/2}, \dots\}$,

$\pi_{\xi, \lambda}^-$ on $\text{Cl Span } \{\dots, \phi_{-\lambda-3/2}, \phi_{-\lambda-1/2}\}$,

$\pi_{\xi, \lambda}^0$ on $\text{Span } \{\phi_{-\lambda+1/2}, \phi_{-\lambda+3/2}, \dots, \phi_{\lambda-1/2}\}$.

Among these $\pi_{\xi, \lambda}^+$ and $\pi_{\xi, \lambda}^-$ are subrepresentations.

(d) $\lambda + \xi \in \mathbf{Z} + \frac{1}{2}, \lambda < 0$:

$\pi_{\xi, \lambda}^+$ on $\text{Cl Span } \{\phi_{-\lambda+1/2}, \phi_{-\lambda+3/2}, \dots\}$,

$\pi_{\xi, \lambda}^-$ on $\text{Cl Span } \{\dots, \phi_{\lambda-3/2}, \phi_{\lambda-1/2}\}$,

$\pi_{\xi, \lambda}$ on $\text{Span } \{\phi_{\lambda+1/2}, \phi_{\lambda+3/2}, \dots, \phi_{-\lambda-1/2}\}$.

Among these $\pi_{\xi, \lambda}^0$ is a subrepresentation.

Proof.

(a) $c_{\xi, \lambda, m, n} \neq 0$.

(b) $c_{1/2, 0, m, n} \neq 0 \Leftrightarrow m, n \leq -\frac{1}{2}$ or $m, n \geq \frac{1}{2}$.

(c) $c_{\xi, \lambda, m, n} \neq 0 \Leftrightarrow -\lambda + \frac{1}{2} \leq n \leq \lambda - \frac{1}{2}$
or $m, n \leq -\lambda - \frac{1}{2}$ or $m, n \geq \lambda + \frac{1}{2}$.

Thus $c_{\xi, \lambda, m, n}$ has block matrix

$$\begin{array}{ccc}
 n \leq -\lambda - \frac{1}{2} & -\lambda + \frac{1}{2} \leq n \leq \lambda - \frac{1}{2} & n \geq \lambda + \frac{1}{2} \\
 \begin{array}{l} m \leq -\lambda - \frac{1}{2} \\ -\lambda + \frac{1}{2} \leq m \leq \lambda - \frac{1}{2} \\ m \geq \lambda + \frac{1}{2} \end{array} & \left(\begin{array}{ccc} * & * & 0 \\ 0 & * & 0 \\ 0 & * & * \end{array} \right) &
 \end{array}$$

where each starred block has all entries nonzero.

(d) $c_{\xi, \lambda, m, n} \neq 0 \Leftrightarrow \lambda + \frac{1}{2} \leq m \leq -\lambda - \frac{1}{2}$ or $m, n \leq \lambda - \frac{1}{2}$
 or $m, n > -\lambda + \frac{1}{2}$. □

The finite-dimensional representation occurring in the above classification are the representations $\pi_{\xi, \lambda}^0(\lambda + \xi \in \mathbf{Z} + \frac{1}{2}, \lambda \neq 0)$.

3.3. NOTES

3.3.1. In the case of the unitary principal series (λ imaginary), Theorem 3.4 was first proved by BARGMANN [2, sections 6 and 7]. See van DIJK [9, Theorem 4.1] for the statement and (infinitesimal) proof of our Theorem 3.4 in the general case. A proof of Theorem 3.4 similar to our proof was earlier given by BARUT & PHILLIPS [3, §II (4)].

3.3.2. Theorem 3.4 in the case of imaginary and nonzero λ is contained in a general theorem by BRUHAT [5, Theorem 7; 2]: For $\xi \in \hat{M}$, $\lambda \in ia$, the principal series representation $\pi_{\xi, \lambda}$ of G (cf. (2.2)) is irreducible if $s \cdot \lambda \neq \lambda$ for all $s \neq e$ in the Weyl group for (G, K) .

3.3.3. GELFAND & NAIMARK [18, §5.4, Theorem 1] proved the irreducibility of the unitary principal series for $SL(2, \mathbf{C})$ by a global method different from ours, working in a noncompact realization and calculating the “matrix elements” of the representation with respect to a (continuous) \bar{N} -basis.

3.3.4. Analogues of Theorems 3.2 and 3.3 can be formulated in the case of non-abelian K , cf. [27, Theorem 3.3]. In that case the canonical matrix elements $\tau_{\gamma, \delta}$ are matrix-valued functions. By using this method, NAIMARK [34, Ch. 3, §9, No. 15] examined the irreducibility of the nonunitary principal series for $SL(2, \mathbf{C})$, see also KOSTERS [28].