

2. Minimal left ideals in right topological semigroups

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **35 (1989)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **19.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

2. MINIMAL LEFT IDEALS IN RIGHT TOPOLOGICAL SEMIGROUPS

We present in this section several well known facts which are not usually seen in early graduate courses.

2.1 LEMMA (Ellis [2]). *Let S be a compact Hausdorff right topological semigroup. Then S has an idempotent, that is there exists $x \in S$ with $x + x = x$.*

Proof. Let $\mathcal{A} = \{A \subseteq S : A \neq \emptyset, A \text{ is compact, and } A + A \subseteq A\}$. Now $\mathcal{A} \neq \emptyset$ since $S \in \mathcal{A}$. Let $\mathcal{C}$ be a chain in $\mathcal{A}$. Then $\mathcal{C}$ is a collection of closed subsets of S with the finite intersection property so $\bigcap \mathcal{C} \neq \emptyset$ and $\bigcap \mathcal{C}$ is compact. Trivially $(\bigcap \mathcal{C}) + (\bigcap \mathcal{C}) \subseteq \bigcap \mathcal{C}$. Pick by Zorn's Lemma a minimal member A of $\mathcal{A}$.

Pick $x \in A$ and let $B = A + x$. Now $B = \rho_x[A] (= \{\rho_x(y) : y \in A\})$ so, as the continuous image of a compact set, B is compact (and trivially non-empty). Also $B + B = A + x + A + x \subseteq A + A + A + x \subseteq A + x = B$. Thus $B \in \mathcal{A}$. Since $B = A + x \subseteq A + A \subseteq A$ and A is minimal, $B = A$ so $x \in B = A + x$. That is, there exists $y \in A$ with $x = y + x$.

Let $C = \{y \in A : x = y + x\}$. ρ_x is continuous so $\rho_x^{-1}[\{x\}]$ is closed. Thus C is closed, hence compact. To see that $C + C \subseteq C$, let $y, z \in C$. Then $y + z \in A$ and $(y+z) + x = y + (z+x) = y + x = x$ so $y + z \in C$. Thus $C \in \mathcal{A}$. Since $C \subseteq A$ and A is minimal, $C = A$. Then $x \in C$ and hence $x + x = x$. $\square$

A non-empty subset I of a semigroup S is a left ideal if $S + I \subseteq I$, a right ideal if $I + S \subseteq I$, and a two-sided ideal if it is both a left ideal and a right ideal. It is a fact (which we will not need) that any right ideal in a compact right topological semigroup contains a minimal right ideal, which need not be closed. (For this and other interesting facts see [1].) We do need a similar fact about left ideals.

2.2 LEMMA. *Let S be a compact Hausdorff right topological semigroup. Any left ideal contains a minimal left ideal and minimal left ideals are closed.*

Proof. Let L be a left ideal of S . Let $\mathcal{A} = \{A \subseteq L : A \text{ is a left ideal and } A \text{ is compact}\}$. Choose $x \in L$. Then $S + x = \rho_x[S]$, the continuous image of a compact space. Also $S + (S + x) = (S + S) + x \subseteq S + x$ so $S + x$ is a left ideal. Since $S + x \subseteq S + L \subseteq L$, we have $\mathcal{A} \neq \emptyset$. One easily sees that the intersection of a chain in $\mathcal{A}$ is again in $\mathcal{A}$. Choose by Zorn's Lemma a minimal member A of $\mathcal{A}$.

To see that A is in fact a minimal left ideal, assume we have a left ideal $B \subseteq A$ and pick $x \in B$. Then as above $S + x \in \mathcal{A}$ while $S + x \subseteq S + B \subseteq B \subseteq A$ so $S + x = A$ so $B = A$ $\square$

2.3 Definition. Let S be a semigroup. Then $M(S) = \cup \{L: L \text{ is a minimal left ideal of } S\}$.

It is a fact (which we will not need) that if S is a compact Hausdorff right topological semigroup, then $M(S)$ is a two-sided ideal of S .

2.4 LEMMA. Let S be a compact Hausdorff right topological semigroup and let I be a two-sided ideal of S . Then $M(S) \neq \emptyset$ and $M(S) \subseteq I$.

Proof. Since S is a left ideal of S it contains by Lemma 2.2 a minimal left ideal so $M(S) \neq \emptyset$. So see that $M(S) \subseteq I$, let $x \in M(S)$. There is a minimal left ideal L of S with $x \in L$. Also choose some $y \in I$. Then $y + x \in L \cap I$ (since I is a right ideal) so $L \cap I \neq \emptyset$. Thus $L \cap I$ is a left ideal contained in L so that $L \cap I = L$. $\square$

The proof of the following lemma is an easy exercise and we omit it.

2.5 LEMMA. Let S_1 and S_2 be compact right topological semigroups and let $S_1 \times S_2$ have the product topology and coordinatewise operations. Then $S_1 \times S_2$ is a compact right topological semigroup. Given $x \in S_1$ and $y \in S_2$, λ_x and λ_y may or may not be continuous (where $\lambda_x(t) = x + t$). If $\lambda_x: S_1 \rightarrow S_1$ and $\lambda_y: S_2 \rightarrow S_2$ are continuous, then $\lambda_{(x,y)}: S_1 \times S_2 \rightarrow S_1 \times S_2$ is continuous.

3. VAN DER WAERDEN'S THEOREM

We let $l \in \mathbb{N}$ be fixed throughout and show that given any finite partition of $\mathbb{N}$ some one cell contains a length l arithmetic progression.

3.1 Definition. (a) Let $Y = (\beta\mathbb{N})^l$ with the product topology and coordinatewise operations.

$$(b) \quad E^* = \{(a, a + d, a + 2d, \dots, a + (l-1)d): a \in \mathbb{N} \text{ and } d \in \mathbb{N} \cup \{0\}\}.$$

$$(c) \quad I^* = \{(a, a + d, a + 2d, \dots, a + (l-1)d): a, d \in \mathbb{N}\}.$$

$$(d) \quad E = cl_Y E^*$$

$$(e) \quad I = cl_Y I^*.$$

Note that by Lemmas 1.1 and 2.5, Y is a compact Hausdorff right topological semigroup and whenever $\mathbf{x} = (x_1, x_2, \dots, x_l) \in \mathbb{N}^l$, $\lambda_{\mathbf{x}}$ is continuous.