

3. COROLLARIES OF THE THEOREM OF THE KERNEL

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **36 (1990)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **26.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

- (iii) $(h_{m,1})^{-1}(T) \cap C_m(S)$ is infinite,
- (iv) There exists a finite covering $S_{m,n}$ of S such that the fiber product of $h_{m,n}$ with $S_{m,n}$ is Galois, Abelian and of positive degree.

Let J denote the Jacobian scheme of C over S . Let $a: C \rightarrow J$ be an Albanese morphism. Let p be a prime. Let $\bar{T}$ denote the closure of $a(T)$ in $J(S) \otimes \mathbf{Z}_p$. Since $a(T)$ is infinite it follows from the Mordell-Weil Theorem that there exists a $t \in \bar{T} - a(T)$. Let $t_n \in T$ such that $t - a(t_n) \in p^n J(S)$. Let C_n denote the normalization of the fiber-product of C and J via the map $H_n: x \rightarrow p^n x + t_n$ and $h_{n,1}$ the natural map from C_n to C . It follows that C_n is defined over S and since $H_m(J(S)) \supseteq \{t_n: m \mid n\}$ that $h_{n,1}(C_m(S))$ contains an infinite subset of T .

All that remains is to exhibit the maps $h_{m,n}$. Clearly, $t_m - t_n = p^n r_{m,n}$ for some $r_{m,n} \in J(S)$. Let $H_{m,n}$ denote the map $x: p^{m-n}x + r_{m,n}$. Then $H_{m,k} = H_{n,k} \circ H_{m,n}$. It follows that $H_{m,n}$ pulls back to a morphism $h_{m,n}: C_m \rightarrow C_n$. It is easy to see that this morphism becomes Abelian after adjoining the p^{m-n} -torsion points on J . This proves the proposition. $\square$

Remark. One can also prove the above proposition with the condition $n \leq m$ replaced by $n \mid m$.

3. COROLLARIES OF THE THEOREM OF THE KERNEL

LEMMA 3.3.1. Suppose $g: X' \rightarrow X$ is a morphism of smooth proper schemes with geometrically connected fibers over S . Then if $\mu \in PF(X'/S)$ and $s, t \in X(S)$, $(g^*\mu)(s, t) = \mu(g \circ s, g \circ t)$.

Proof. This follows easily from Lemma 1.3.2. $\square$

Suppose J is the Jacobian of C over S and g is an Albanese morphism, then since $g^*: H_{DR}^1(J/S) \rightarrow H_{DR}^1(C/S)$ is an isomorphism $g^*: PF(J/S) \rightarrow PF(C/S)$ is an isomorphism.

LEMMA 3.3.2. Let μ be a fixed Picard-Fuchs differential equation on C/S . Then $\{\mu(s, t): s, t \in C(S)\}$ lies in a finite dimensional subspace of $K[S]$ over K .

Proof. Suppose $\tilde{\mu} \in PF(J/S)$ such that $g^*\tilde{\mu} = \mu$. The lemma follows from the Mordell-Weil theorem which together with the Theorem of the kernel implies that $J(S)$ modulo the kernel of the homomorphism $s \rightarrow \tilde{\mu}(e, s)$ is a finitely generated Abelian group. $\square$

LEMMA 3.3.3. *Suppose A is an Abelian scheme over S such that $[W_{A/S}] = H_{DR}^1(A/S)$ and $g: C \rightarrow A$ is a non-constant morphism over S . Fix $s \in C(S)$. Then the set $T = \{t \in C(S) : (g^*\mu)(s, t) = 0 \text{ for all } \mu \in PF(A/S)\}$ is of bounded height.*

Proof. Let A' denote the smallest Abelian subscheme of A over S containing $g(C)$. Since the map $g^*: PF(A/S) \rightarrow PF(A'/S)$ is surjective and $[W_{A/S}] = H_{DR}^1(A/S)$, it follows from Proposition 2.1.2 that $g(T)$ is contained in a translation of the group of constant sections of A'/S . Hence, $g(T)$ is a set of bounded height. Finally, since $C \rightarrow g(C)$ is a finite morphism, it follows that T is a set of bounded height. $\square$

In particular,

COROLLARY 3.3.4. *Suppose A is an Abelian scheme over S such that $\kappa_{A/S}$ is an isomorphism and $g: C \rightarrow A$ is a non-constant morphism over S . Fix $s \in C(S)$. Then the set $\{t \in C(S) : (g^*\mu_\omega)(s, t) = 0 \text{ for all } \omega \in \omega_{A/S}\}$ is of bounded height.*

4. PROOF OF MORDELL'S CONJECTURE

PROPOSITION 3.4.1. *Suppose the kernel of the $\kappa_{C/S}$ has rank at least 2 over $K[S]$, then the points of $C(S)$ have bounded height.*

Proof. Suppose $C(S)$ contains points of arbitrarily large height. Fix $s \in C(S)$. By shrinking S , if necessary, we may suppose that there exists a function $z \in K[S]$ such that $\Omega_S^1 = K[S]dz$ and there exists a finite covering $\mathcal{C}$ of C by affine opens U and functions $v_U \in \mathcal{O}_C(U)$ such that $s \in U(S)$, and $\Omega_C^1(U)$ is spanned by dz and dv_U . We may also suppose that $s^*v_U = 0$ by replacing v_U with $v_U - (s \circ f)^*v_U$ if necessary. For $U \in \mathcal{C}$, $u \in \mathcal{O}_C(U)$ we define $\partial_{U,z}u$ and $\partial_{U,v}u$ by the equation

$$du = \partial_{U,z}u dz + \partial_{U,v}u dv_U.$$

Then $\partial_{U,z}$ is a lifting of $\partial = : \partial/\partial z$. We set $\mu(t) = \mu(s, t)$ for

$$\mu \in PF = : PF(C/S)$$

and $t \in C(S)$.

Let ω_1 and ω_2 be two independent elements in the kernel of $\kappa_{C/S}$. It follows that there exist ω'_1 and $\omega'_2 \in \omega_{C/S}$ such that

$$\partial[\omega'_i] = [\omega'_i].$$