

# PERIODIC KNOTS, SMITH THEORY, AND MURASUGI'S CONGRUENCE

Autor(en): **Davis, James F. / Livingston, Charles**

Objektyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **37 (1991)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **24.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-58725>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

P 74 771

PERIODIC KNOTS, SMITH THEORY,  
AND MURASUGI'S CONGRUENCE



by James F. DAVIS and Charles LIVINGSTON

A knot  $K$  in a homology 3-sphere  $\Sigma$  has period  $n$  if it is invariant under a homeomorphism  $h: \Sigma \rightarrow \Sigma$  of order exactly  $n$  with fixed set  $B$ , a circle disjoint from  $K$ . The quotient space  $\bar{\Sigma} = \Sigma/h$  is a homology sphere containing  $\bar{K}$ , the quotient knot. Kunio Murasugi [Mu] discovered the following congruence involving the Alexander polynomials of the two knots. (See also the proof by J. Hillman [H].)

**THEOREM A.** *Let  $K$  be a knot of prime power period  $p^r$  in a homology 3-sphere  $\Sigma$  with fixed set  $B$  and quotient knot  $\bar{K}$ . Let  $\Delta_K(t)$  and  $\Delta_{\bar{K}}(t)$  be their Alexander polynomials and let  $\lambda$  be the linking number of  $K$  and  $B$ . Then*

$$\Delta_K(t) \doteq \Delta_{\bar{K}}(t)^{p^r} (1 + t + \dots + t^{\lambda-1})^{p^r-1} \pmod{p},$$

where  $\doteq$  means congruent up to multiplication by  $ut^i$  where  $u$  and  $i$  are integers and  $u$  is relatively prime to  $p$ .

In another direction it is easily shown that if  $G = \mathbf{Z}/p$  acts cellularly on a finite CW complex  $X$ , then  $\chi(X) + (p-1)\chi(X^G) = p\chi(X/G)$ . Using Smith theory, E. Floyd [F] gave a proof of this when  $X$  is a finite-dimensional CW complex with  $\text{rk } H_*(X; \mathbf{Z}/p) < \infty$ . The proof can be generalized easily to the case of semifree actions of a  $p$ -group  $G$  on  $X$ . (An action is semifree if every point in  $X$  is either freely permuted by  $G$  or fixed by all of  $G$ . An action of  $\mathbf{Z}/p$  is automatically semifree.) We will prove a multiplicative analogue of Floyd's theorem and use it to deduce Murasugi's congruence.

If  $X$  is a space with an action of the infinite cyclic group  $C_\infty = \langle t \rangle$  and  $F$  is a field with  $\text{rk } H_*(X; F) < \infty$ , we define a multiplicative Euler characteristic

$$\chi_m(X; F) \in F(t)^*/F[t, t^{-1}]^*$$

to be the alternating product of the generator of the order ideals of  $H_i(X; F)$ .

(See [Mi] or §1 for definitions). We will be most interested in the case  $F = \mathbf{F}_p$ , the finite field with  $p$  elements.

**THEOREM B.** *Let  $G$  be a  $p$ -group. Suppose  $C_\infty \times G$  act on a finite-dimensional CW complex  $X$  with  $\text{rk } H_*(X; \mathbf{F}_p) < \infty$ , so that  $G$  acts semifreely and cellularly. Then*

$$\chi_m(X; \mathbf{F}_p) \chi_m(X^G; \mathbf{F}_p)^{|G|-1} = \chi_m(X/G; \mathbf{F}_p)^{|G|}.$$

Applying this to the case where  $X$  is the infinite cyclic cover of  $\Sigma - K$  will immediately yield Murasugi's congruence. One advantage of our approach is that it generalizes to the case of high-dimensional periodic knots.

In §1 we prove Theorem B and derive Theorem A. In §2 we discuss the high-dimensional case and in §3 give the following application of Murasugi's congruence to links.

**PROPOSITION C.** *Let  $L$  be a two-component link in a homology 3-sphere. If the  $\mathbf{Z}/2 \times \mathbf{Z}/2$ -cover branched over the link is also a homology 3-sphere, then the linking number of the two components is congruent to  $\pm 1$  modulo 8.*

## §1. MURASUGI'S CONGRUENCE

We will derive Theorem A from Theorem B and then prove Theorem B, but we first give some homological preliminaries. If  $R$  is a commutative Noetherian UFD with quotient field  $K$  and  $M$  is a finitely generated torsion  $R$ -module then we define the *order* of  $M$  to be  $[M] = E^0(M) \in R/R^*$ . Here we take an exact sequence

$$R^k \xrightarrow{A} R^m \rightarrow M \rightarrow 0,$$

and we let  $E^0(M)$  be a greatest common divisor of the determinants of the  $m \times m$ -submatrices of  $A$ . If  $M$  is a torsion f.g.  $R$ -module then  $[M] \neq 0$ , and we consider the order  $[M]$  as an element of  $K^*/R^*$ . If

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

is an exact sequence of torsion f.g.  $R$ -modules, then J. Levine [L, lemma 5] shows  $[M] = [M'] [M'']$ . It follows for formal reasons that if  $C_* = \{C_n \rightarrow \dots \rightarrow C_0\}$  is a chain complex of torsion f.g.  $R$ -modules then

$$\chi_m(C_*) := \prod [C_i]^{(-1)^i}$$

equals  $\chi_m(H_*(C_*))$ . In particular if  $C_*$  is exact, then  $\chi_m(C_*) = 1$ .

Next we turn to Alexander polynomials. By Alexander duality  $H_1(\Sigma - K) \cong \mathbf{Z}$ . Let  $\pi: X \rightarrow \Sigma - K$  be the infinite cyclic cover of the knot complement. The infinite cyclic group  $C_\infty = \langle t \rangle$  acts on  $X$  and  $H_1(X; \mathbf{Z})$  is a f.g. torsion module over the group ring  $\mathbf{Z}[C_\infty] = \mathbf{Z}[t, t^{-1}]$ . The Alexander polynomial  $\Delta_K(t)$  is its associated order. (Note that  $\mathbf{Z}[t, t^{-1}]^*$  consists of  $\pm t^i$  and the quotient field of  $\mathbf{Z}[t, t^{-1}]$  is the field of rational functions  $\mathbf{Q}(t)$ .) As usual we normalize so that  $\Delta_K(t)$  is a polynomial with integer coefficients and non-zero constant term.

If  $K$  has period  $p^r$ , let  $\bar{\pi}: \bar{X} \rightarrow \bar{\Sigma} - \bar{K}$  be the infinite cyclic cover of the quotient knot. The  $G = \mathbf{Z}/p^r$ -action on  $\Sigma - K$  lifts to a  $G$ -action on  $X$  with quotient  $\bar{X}$  and fixed set  $\bar{B} = \pi^{-1}(B)$ . Indeed, let  $g$  be a generator of  $G$ . Then  $g \circ \pi: X \rightarrow \Sigma - K$  induces the trivial map on  $H_1$  and so lifts to  $\tilde{g}: X \rightarrow X$ . Since  $g$  has a non-empty, path-connected fixed-point set there is a unique lift  $\tilde{g}$  with fixed points and the fixed point set is  $\bar{B}$ . Since  $\tilde{g}^{p^r}$  is a lift of the identity which has fixed points, it itself is the identity and hence  $\tilde{g}$  is a map of period  $p^r$ . This gives an action of  $C_\infty \times G$  on  $X$ . It further follows that  $X/G \rightarrow \bar{\Sigma} - \bar{K}$  is an abelian cover inducing the trivial map on  $H_1$ , so that we can identify this cover with  $\bar{\pi}$  and  $X/G$  with  $\bar{X}$ .

The cover  $\pi$  is classified by a map  $c: \Sigma - K \rightarrow S^1 = K(\mathbf{Z}, 1)$  inducing an isomorphism on  $H_1$ . The inclusion map  $B \rightarrow \Sigma - K$  induces multiplication by the linking number  $\lambda$  on  $H_1$ . Thus by considering  $c|_B$  which classifies  $\pi: \bar{B} \rightarrow B$ , we see  $\bar{B}$  is homeomorphic to  $\lambda$  disjoint copies of  $\mathbf{R}$ , cyclically permuted by the action of  $C_\infty$ .

Now  $H_i(X)$  and  $H_i(\bar{X})$  are zero for  $i > 1$  and  $H_0(X)$  and  $H_0(\bar{X})$  are isomorphic to  $\mathbf{F}_p \cong \mathbf{F}_p[t, t^{-1}]/(t-1)\mathbf{F}_p[t, t^{-1}]$ , so  $\chi_m(X) = (t-1)/\Delta_K(t)$  and  $\chi_m(\bar{X}) = (t-1)/\Delta_{\bar{K}}(t)$ . Since  $X^G = \bar{B}$  consists of  $\lambda$  arcs cyclically permuted by  $C_\infty = \langle t \rangle$ ,  $\chi(X^G) = t^\lambda - 1$ . Putting this together with Theorem B we see

$$[(t-1)/\Delta_K(t)] [t^\lambda - 1]^{p^r - 1} = [(t-1)/\Delta_{\bar{K}}(t)]^{p^r}$$

or  $\Delta_K(t) = \Delta_{\bar{K}}(t)^{p^r} (1 + t + \dots + t^{\lambda-1})^{p^r - 1}$  with the equality taking place in  $\mathbf{F}_p(t)/\mathbf{F}_p[t, t^{-1}]^*$ . This gives Murasugi's congruence.

*Proof of Theorem B.* We prove the theorem by induction on the order of  $G$ . Let  $G$  be a group of prime order  $p$  with generator  $g$ . Let



$$\sigma = 1 + g + g^2 + \dots + g^{p-1}$$

$$\delta = 1 - g$$

be elements of the group ring  $\mathbf{F}_p[G]$ . Note that  $\delta\sigma = 0 = \sigma\delta$  and  $\delta^{p-1} = \sigma$ . We consider the following chain complexes of  $\mathbf{F}_p[t, t^{-1}]$ -modules (all homology is with  $\mathbf{F}_p$ -coefficients).

$$\begin{aligned} 0 &\rightarrow C_*(X^G) \rightarrow C_*(\bar{X}) \xrightarrow{\text{tr}} \sigma C_*(X) \rightarrow 0 \\ 0 &\rightarrow \delta C_*(X) \oplus C_*(X^G) \rightarrow C_*(X) \xrightarrow{\sigma} \sigma C_*(X) \rightarrow 0 \\ 0 &\rightarrow \sigma C_*(X) \rightarrow \delta C_*(X) \xrightarrow{\delta} \delta^2 C_*(X) \rightarrow 0 \\ &\quad \vdots \\ 0 &\rightarrow \sigma C_*(X) \rightarrow \delta^{p-2} C_*(X) \xrightarrow{\delta} \delta^{p-1} C_*(X) \rightarrow 0. \end{aligned}$$

These induce long exact sequences in homology. All homology is finitely generated and torsion over the PID  $\mathbf{F}_p[t, t^{-1}]$ . We use shorthand notation – if  $\rho \in \mathbf{F}_p[G]$ , we write  $\chi^\rho(X)$  instead of  $\chi(H_*(\rho C_*(X)))$ . The above homological considerations show

$$\begin{aligned} \chi(\bar{X}) &= \chi(X^G)\chi^\sigma(X) \\ \chi(X) &= \chi^\delta(X)\chi(X^G)\chi^\sigma(X) \\ \chi^\delta(X) &= \chi^\sigma(X)\chi^{\delta^2}(X) \\ &\quad \vdots \\ \chi^{\delta^{p-2}}(X) &= \chi^\sigma(X)\chi^\sigma(X). \end{aligned}$$

Multiplying all equations but the first together and cancelling terms we see

$$\chi(X) = \chi(X^G) \cdot \chi^\sigma(X)^p.$$

Using the first equation to substitute for  $\chi^\sigma(X)$  one finds

$$\chi(X) = \chi(\bar{X})^p / \chi(X^G)^{p-1}.$$

Finally suppose  $G$  has order  $p^r$ . Let  $G_1$  be a normal subgroup of index  $p$ . By the exact sequences above  $\text{rk } H_*(X/G_1; \mathbf{F}_p) < \infty$ . By applying inductively the result for the  $G_1$ -action on  $X$  and the  $G/G_1$  action on  $X/G_1$ , Theorem B follows.

## §2. HIGH-DIMENSIONAL PERIODIC KNOTS

One advantage of our approach to Murasugi's congruence is that it applies equally well to a more general situation. Higher-dimensional periodic knots

were introduced in the thesis of *R. Cruz* [C]. He showed that if there is a semifree  $\mathbf{Z}/q$ -action on  $S^n$  with non-empty fixed set and an invariant knot  $K^{n-2}$  disjoint from the fixed set, then the fixed set is  $S^1$  if  $q \neq 2$ , and is  $S^1$  or  $S^0$  if  $q = 2$ .

For our purposes a knot  $K$  in a homology  $n$ -sphere  $\Sigma$  is an embedded  $(n - 2)$ -dimensional homology sphere. Let  $G$  be a finite group. The knot  $K$  is *G-periodic* if it is invariant under a semifree  $G$ -action on  $\Sigma$  with fixed set  $B \cong S^1$  disjoint from  $K$ . To simplify technicalities we assume the action is smooth. Several complications arise: the group need not be cyclic, the action need not be linear and the quotient  $\bar{\Sigma} = \Sigma/G$  will not be a manifold. (Even in the linear case the quotient looks like a double suspension of a spherical space form.) However we can still make sense of Alexander polynomials.

PROPOSITION 2.1.  $H_*(\bar{\Sigma} - \bar{K}) \cong H_*(S^1)$ .

First we need a lemma.

LEMMA 2.2. *The linking number  $\lambda = \text{lk}(B, K)$  is relatively prime to the order of  $G$ .*

*Proof.* (See also [C, 2.1.1]). By restricting the action to a subgroup  $\mathbf{Z}/p$  of  $G$ , we will assume  $G = \mathbf{Z}/p$ , and show  $(\lambda, p) = 1$ . By applying the Lefschetz Fixed-Point Theorem to a generator  $g$  of  $\mathbf{Z}/p$ , we see that if  $n$  is odd, the action on  $K$  is orientation-preserving, while if  $n$  is even, then  $p = 2$  and the action is orientation-reversing. For local coefficients we will use  $\mathbf{Z}^t$ , the integers with the  $\mathbf{Z}[\mathbf{Z}/p]$ -module structure given by  $(\sum a_i g^i) \cdot k = \sum a_i (-1)^{i(n+1)} k$ .

Let  $\bar{\Sigma} - B \rightarrow K(\mathbf{Z}/p, 1)$  classify the  $G$ -cover. We will consider the commutative diagram:

$$\begin{array}{ccccc}
 H_{n-2}(K; \mathbf{Z}) & \xrightarrow{\alpha} & H_{n-2}(\bar{K}; \mathbf{Z}^t) & \rightarrow & H_{n-2}(K(\mathbf{Z}/p, 1); \mathbf{Z}^t) \\
 (*) & & \downarrow & & \parallel \\
 & & H_{n-2}(\Sigma - B; \mathbf{Z}) & \rightarrow & H_{n-2}(\bar{\Sigma} - B; \mathbf{Z}^t) \rightarrow H_{n-2}(K(\mathbf{Z}/p, 1); \mathbf{Z}^t) .
 \end{array}$$

The two groups on the left are infinite cyclic and the left vertical map is multiplication by  $\lambda$ . A diagram chase shows we will be done if we can show both horizontal exact sequences are isomorphic to the short exact sequence  $0 \rightarrow \mathbf{Z} \rightarrow \mathbf{Z} \rightarrow \mathbf{Z}/p \rightarrow 0$ .

The map  $\alpha$  is isomorphic to  $\mathbf{Z} \xrightarrow{\times p} \mathbf{Z}$  because it comes from a  $p$ -fold cover of  $(n - 2)$ -dimensional closed manifolds. The map

$$H_{n-2}(\bar{K}; \mathbf{Z}^t) \rightarrow H_{n-2}(\mathbf{Z}/p; \mathbf{Z}^t)$$

we compute algebraically by using a free  $\mathbf{Z}G$ -resolution of  $\mathbf{Z}$  as a substitute for the Eilenberg-MacLane space. By lifting a CW structure on  $\bar{K}$  to  $K$ ,

$$C_*(K) = \{C_{n-2} \rightarrow \dots \rightarrow C_0\}$$

with the  $i$ -chains  $C_i$  free  $\mathbf{Z}G$ -modules. By mapping a free  $\mathbf{Z}G$ -module onto  $\ker(C_{n-2} \rightarrow C_{n-3})$  and continuing inductively, one constructs a free  $\mathbf{Z}G$ -resolution of  $\mathbf{Z}$

$$D_* = \{\dots \rightarrow D_n \rightarrow D_{n-1} \rightarrow C_{n-2} \rightarrow \dots \rightarrow C_0\}.$$

It follows that

$$H_{n-2}(\bar{K}; \mathbf{Z}^t) = H_{n-2}(C_*(K) \otimes_{\mathbf{Z}G} \mathbf{Z}^t)$$

maps onto  $H_{n-2}(D_* \otimes_{\mathbf{Z}G} \mathbf{Z}^t) = H_{n-2}(\mathbf{Z}/p; \mathbf{Z}^t)$ . Furthermore by using the standard  $\mathbf{Z}G$ -resolution of  $\mathbf{Z}$  (see e.g. [Mac]), one easily computes that  $H_{n-2}(\mathbf{Z}/p; \mathbf{Z}^t) \cong \mathbf{Z}/p$ .

Choose a  $G$ -invariant normal disk to  $B$  in  $\Sigma$  and let  $S^{n-2}$  be its boundary. Then the inclusion  $S^{n-2} \rightarrow \Sigma - B$  is a homology equivalence. By the comparison theorem applied to the spectral sequence of the  $G$ -coverings (see [Mac]), the bottom row of (\*) is isomorphic to

$$H_{n-2}(S^{n-2}; \mathbf{Z}) \rightarrow H_{n-2}(S^{n-2}/G; \mathbf{Z}^t) \rightarrow H_{n-2}(G; \mathbf{Z}^t),$$

and hence by the previous paragraph to  $0 \rightarrow \mathbf{Z} \rightarrow \mathbf{Z} \rightarrow \mathbf{Z}/p \rightarrow 0$ . Thus  $(\lambda, p) = 1$ .

*Proof of 2.1.* Let  $N$  be an equivariant tubular neighborhood of  $B$ . Then

$$0 = H_*(\Sigma - K, N; \mathbf{Z}[1/\lambda]) = H_*(\Sigma - K - B, N - B; \mathbf{Z}[1/\lambda])$$

where the first equality holds by the definition of linking number and the second by excision. Then

$$\begin{aligned} 0 &= H_*((\Sigma - K - B)/G, (N - B)/G; \mathbf{Z}[1/\lambda]) = H_*((\Sigma - K)/G, N/G; \mathbf{Z}[1/\lambda]) \\ &= H_*((\Sigma - K)/G, B; \mathbf{Z}[1/\lambda]), \end{aligned}$$

where the first equality follows from the spectral sequence of a covering, the second by excision and the third by the homotopy equivalence  $B \rightarrow N/G$ . Thus  $H_*(\bar{\Sigma} - \bar{K})$  looks like  $H_*(S^1)$  except possibly for some  $\lambda$ -torsion. But by 2.1,  $\lambda$  is prime to the order of  $G$ , so for all primes  $q$  dividing  $\lambda$ , the transfer map  $\text{tr}: H_*(\bar{\Sigma} - \bar{K}; \mathbf{Z}/q) \rightarrow H_*(\Sigma - K; \mathbf{Z}/q)$  is injective so there is no extra  $\lambda$ -torsion.

To state Murasugi's congruence in higher dimensions is it necessary to find a substitute for the Alexander polynomial. Let  $X$  and  $\bar{X}$  be the infinite cyclic

covers of  $\Sigma - K$  and  $\bar{\Sigma} - \bar{K}$  respectively. Let  $\Delta_K(t) = \prod_{i>0} [H_i(X)]^{(-1)^{i+1}}$  and  $\Delta_{\bar{K}}(t) = \prod_{i>0} [H_i(\bar{X})]^{(-1)^{i+1}}$ . The Wang sequence shows that multiplication by  $t - 1$  induces an isomorphism on  $H_i(X)$  for  $i > 0$ , so that if we take the polynomial represented by  $[H_i(X)]$  and plug in  $t = 1$  we get  $\pm 1$ . (Indeed if we consider the ring homomorphism  $\varphi: \mathbf{Z}[t, t^{-1}] \rightarrow \mathbf{Z}$  defined by  $\varphi(t) = 1$ , then  $\varphi([H_i(X)])$  is a divisor of  $[H_i(X) \otimes_{\mathbf{Z}[t, t^{-1}]} \mathbf{Z}] = [0] = 1 \in \mathbf{Z}/\mathbf{Z}^*$ .) Thus  $[H_i(X)]$  represented a non-zero element in  $\mathbf{F}_p[t, t^{-1}]$ , and hence  $\Delta_K(t)$  and  $\Delta_{\bar{K}}(t)$  give well-defined elements of  $\mathbf{F}_p(t)^*/\mathbf{F}_p[t, t^{-1}]^*$ . Then the considerations of §1 show:

**THEOREM 2.3.** *Let  $K$  be a  $G$ -periodic knot in a homology  $q$ -sphere  $\Sigma$  with fixed set  $B$ , where  $G$  is a group of prime power order  $p^r$ . Let  $\lambda$  be the linking number of  $K$  and  $B$ . Then*

$$\Delta_K(t) \equiv \Delta_{\bar{K}}(t)^{p^r} (1 + t + \dots + t^{\lambda-1})^{p^r-1} \pmod{p} .$$

### §3. AN APPLICATION OF MURASUGI'S CONGRUENCE

For any  $\lambda \equiv \pm 1 \pmod{8}$ , T. tom Dieck and J. Davis [D-D] constructed a 2-component link with linking number  $\lambda$  in a homology 3-sphere  $\Omega$  whose  $C_2 \times C_2$ -cover branched over the link is a homology 3-sphere  $\Sigma$ . We will show that this congruence condition is necessary. Equivalently, we show

**THEOREM 3.1.** *Suppose the Klein 4-group  $G \times H \cong C_2 \times C_2$  acts on a homology 3-sphere  $\Sigma$  so that the fixed sets  $\Sigma^G$  and  $\Sigma^H$  are disjoint circles. Then their linking number  $\lambda$  is congruent to  $\pm 1$  modulo 8.*

*Proof.* We have

$$\begin{array}{ccc} \Sigma & \rightarrow & \Sigma/G \\ \downarrow & & \downarrow \\ \Sigma/H & \rightarrow & \Sigma/(G \times H) . \end{array}$$

All four of these manifolds are homology 3-spheres and each has two disjoint circles given by the images of the fixed sets. The linking numbers of each pair of circles are all equal.

Let  $K = \Sigma^G/G \subset \Sigma/G$  and  $\bar{K} = K/H \subset \Sigma/(G \times H)$ . Then  $K$  is a knot of period 2. Renormalize  $\Delta_K(t)$  and  $\Delta_{\bar{K}}(t) \in \mathbf{Z}[t, t^{-1}]$  so that  $\Delta_K(t) = \Delta_K(t^{-1})$ ,  $\Delta_{\bar{K}}(t) = \Delta_{\bar{K}}(t^{-1})$ , and  $\Delta_K(1) = 1 = \Delta_{\bar{K}}(1)$ . Murasugi's congruence shows

$$(**) \quad \Delta_K(t) = \Delta_{\bar{K}}(t)^2(t^{(1-\lambda)/2} + \dots + 1 + \dots + t^{(\lambda-1)/2}) + 2f(t),$$

where  $f(t) \in \mathbf{Z}[t, t^{-1}]$  satisfies  $f(t) = f(t^{-1})$ . Writing

$$f(t) = a_n t^{-n} + \dots + a_0 + \dots + a_n t^n,$$

we see  $f(1) \equiv f(-1) \pmod{4}$ . Since  $\Sigma \rightarrow \Sigma/G$  is a 2-fold cover branched over  $K$ ,  $|\Delta_K(-1)| = |H_1(\Sigma)| = 1$ . So  $1 = \Delta_K(1) \equiv \Delta_K(-1) \pmod{4}$ , and we see  $\Delta_K(-1) = 1$ . Take equation (\*\*) and plug in  $t = 1$  and  $t = -1$ :

$$1 = 1 \cdot \lambda + 2 \cdot f(1)$$

$$1 = 1 \cdot (-1)^{(\lambda-1)/2} + 2 \cdot f(-1).$$

Thus  $\lambda \equiv (-1)^{(\lambda-1)/2} \pmod{8}$  so  $\lambda \equiv \pm 1 \pmod{8}$ .

Applying the high-dimensional version of Murasugi's congruence one sees that if  $G \times H \cong C_2 \times C_2$  acts on a homology  $q$ -sphere  $\Sigma$  so that  $\Sigma^G$  is a homology  $q - 2$  sphere and  $\Sigma^H$  is a circle disjoint from  $\Sigma^G$ , then their linking number  $\lambda$  is congruent to  $\pm 1$  modulo 8. This and considerations from  $L$ -theory lead us to conjecture that if  $G \times H \cong C_2 \times C_2$  acts on a homology  $q$ -sphere  $\Sigma$  so that  $\Sigma^G$  is a homology  $k$ -sphere and  $\Sigma^H$  is a homology  $q - k - 1$ -sphere disjoint from  $\Sigma^G$ , then their linking number  $\lambda$  is congruent to  $\pm 1$  modulo 8.

*Acknowledgements.* This work was partially supported by an NSF Postdoctoral Fellowship and an NSF grant. We would like to thank Alejandro Adem for a careful reading of an earlier version of this manuscript.

## REFERENCES

- [C] da CRUZ, R. N. *Periodic knots*. Thesis (New York University 1987).
- [D-D] DAVIS, J. F. and T. TOM DIECK. Some exotic dihedral actions on spheres. *Indiana Univ. Math. J.* 37 (1988), 431-450.
- [F] FLOYD, E. On periodic maps and the Euler characteristics of associated spaces. *Trans. Amer. Math. Soc.* 72 (1952), 138-147.
- [H] HILLMAN, J. A. New proofs of two theorems on periodic knots. *Arch. Math. (Basel)* 37 (1981), 457-461.

- [L] LEVINE, J. A method for generating link polynomials. *Amer. J. Math.* 89 (1967), 69-84.
- [Mac] MAC LANE, S. *Homology* (Springer-Verlag 1963).
- [Mi] MILNOR, J. Infinite cyclic covers. *Conf. on topology of manifolds*, ed. J. G. Hocking (Prindle, Weber, and Schmidt, 1968), 115-133.
- [Mu] MURASUGI, K. On periodic knots. *Comment. Math. Helv.* 46 (1971), 162-174.

*(Reçu le 4 août 1989)*

James F. Davis  
Charles Livingston

Department of Mathematics  
Indiana University  
Bloomington, Indiana 47405

**Vide-leer-empty**