

## 4.2. PROBLEMS

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **39 (1993)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **18.09.2024**

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

### **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

$$(8) \quad c(n, r) \ll_{\varepsilon, k} (m + |D|^{1/2+\varepsilon})^{1/2} \frac{|D|^{k/2-3/4}}{m^{(k-1)/2}} \|\phi\| \quad (\varepsilon > 0)$$

where  $D := r^2 - 4mn$  and the bound in  $\ll$  only depends on  $\varepsilon$  and  $k$ .

For the proof one carries over the method of Poincaré series and Kloosterman sums from the one-variable situation already mentioned above to the case of the Jacobi group. Note that Poincaré series on  $\Gamma_1^J$  were studied in [14, II, §2]. The Kloosterman sums that occur in their Fourier coefficients can be related to Salié sums and therefore can easily be estimated (a similar phenomenon happens in the case of modular forms of half-integral weight, cf. [15]). The proof of (8) for  $D$  a fundamental discriminant (i.e. the discriminant of a quadratic field) is given in [20, §1] and for arbitrary  $D$  is given in [21, §1].

On the other hand, if one applies Landau's theorem to the Dirichlet series  $D_{F,F}(s)$  discussed in §3, one finds that

$$(9) \quad \|\phi_m\| \ll_{\varepsilon, F} m^{k/2-2/9+\varepsilon} \quad (\varepsilon > 0).$$

The estimates (8) and (9) now imply the following

**THEOREM 2** [20, 21]. *One has*

$$(10) \quad a(T) \ll_{\varepsilon, F} (\det T)^{k/2-13/36+\varepsilon} \quad (\varepsilon > 0).$$

In fact, both sides of (10) are invariant under  $T \mapsto U'TU$  ( $U \in GL_2(\mathbf{Z})$ ), hence if in (10) we write  $T = \begin{pmatrix} n & r/2 \\ r/2 & m \end{pmatrix}$ , then we may assume that  $m = \min T$ , where  $\min T$  denotes the least positive integer represented by  $T$ . If we use (8) and (9) together with the fact that  $\min T \ll (\det T)^{1/2}$  which is well-known from reduction theory, we obtain (10).

#### 4.2. PROBLEMS

i) In [15], Iwaniec using some sophisticated arguments for certain sums of Salié sums showed that the Fourier coefficients  $a(n)$  ( $n \in \mathbf{N}$ ) of a cusp form  $f$  of weight  $k - 1/2$  for  $k > 0$  and  $n$  squarefree satisfy

$$a(n) \ll_K \sigma_0(n) (\log 2n)^2 n^{k/2-15/28} \|f\|,$$

where  $\sigma_0(n)$  is the number of positive divisors of  $n$  and  $\|f\|$  is the appropriately normalized Petersson norm of  $f$ . We wonder if it is possible to prove an analogous estimate for the Fourier coefficients  $c(n, r)$  ( $D = r^2 - 4mn$  a fundamental discriminant) of a function  $\phi \in J_{k,m}^{\text{cusp}}$  for  $k > 2$  which also is

analogous to (8) in the sense that an appropriate power of  $m$  appears in the denominator on the right-hand side. This then eventually would lead to some improvement of (10) in the case where  $-4 \det T$  is a fundamental discriminant.

ii) For arbitrary genus  $n$ , the best estimate for Fourier coefficients so far known is due to Böcherer and Raghavan [5] and independently Fomenko [11]. Using Rankin's method they showed that the Fourier coefficients  $a(T)$  ( $T$  a positive definite symmetric half-integral  $(n, n)$ -matrix) of a cusp form  $F$  of integral weight  $k$  on  $\Gamma_n$  satisfy

$$a(T) \ll_{\varepsilon, F} (\det T)^{k/2 - \delta_n + \varepsilon} \quad (\varepsilon > 0)$$

where  $\delta_n := 2n + 2 + 4 \left[ \frac{n}{2} \right] + \frac{2}{n+1}$  and  $[x] =$  integral part of  $x$  (the case  $n = 2$  was discussed above).

It is natural to try to apply the method described in 4.1 also for higher genus  $n$ . For some results in this direction we refer to [7].

iii) Let  $F$  be a non-zero Hecke eigenform in  $S_k(\Gamma_2)$  with eigenvalues  $\lambda_n$  and Fourier coefficients  $a(T)$ .

If  $k$  is even and  $F$  is in the Maass space  $S_k^*(\Gamma_2)$ , then

$$(11) \quad \lambda_n \ll_{\varepsilon} n^{k-1+\varepsilon} \quad (\varepsilon > 0)$$

and this estimate is best possible as follows from Theorem 2 in §2.

On the other hand, if  $k$  is odd or if  $k$  is even and  $F \in S_k^*(\Gamma_2)^\perp$ , then one expects that the generalized Ramanujan-Petersson conjecture holds which predicts that

$$(12) \quad \lambda_n \ll_{\varepsilon} n^{k-3/2+\varepsilon} \quad (\varepsilon > 0).$$

To the author's knowledge, the best estimate proved so far for the numbers  $\lambda_n$  is due to Duke, Howe and Li [8] who showed using representation-theoretic methods that

$$(13) \quad \lambda_n \ll_{\varepsilon} n^{k-1+\varepsilon} \quad (\varepsilon > 0)$$

provided that  $n$  is squarefree (in fact, under the assumption  $n$  squarefree the authors proved that  $\lambda_n \leq \sigma_0(n)^2 n^{k-1}$ ; it is suggestive that their method, in fact, gives (13) for all  $n$ ).

In [1, Chap. 2] Andrianov proved that if  $D$  is a negative fundamental discriminant and  $T_1, \dots, T_h$  ( $h = h(D)$ ) denotes a set of  $\Gamma_1$ -representatives of positive definite symmetric half-integral  $(2, 2)$ -matrices with discriminant  $D$ , then

$$(14) \quad \zeta_{\mathbb{Q}(\sqrt{D})}(s - k + 2) \sum_{v=1}^h \left( \sum_{n \geq 1} a(nT_v) n^{-s} \right) = \left( \sum_{v=1}^h a(T_v) \right) Z_F(s).$$

Thus – roughly speaking – for fixed  $T$  the eigenvalues  $\lambda_n$  are “proportional” to the coefficients  $a(nT)$ .

Suppose that  $F$  is in  $S_k^*(\Gamma_2)$ . Using Theorem 1 in §2 and the estimate (8) with  $m = 1$  one finds that

$$(15) \quad a(T) \ll_{\varepsilon, F} (\det T)^{k/2 - 1/2 + \varepsilon} \quad (\varepsilon > 0),$$

and (11) together with (14) implies that (15), in fact, is best possible.

On the other hand, taking into account (12) and the fact that the Hecke eigenforms form a basis, one may be led to the following

CONJECTURE 1 [11]. *Let  $F$  be a cusp form of integral weight  $k$  on  $\Gamma_2$  and suppose that either  $k$  is odd or that  $k$  is even and  $F$  is in the orthogonal complement of the Maass space. Let  $a(T)$  ( $T$  a positive definite symmetric half-integral  $(2, 2)$ -matrix) be the Fourier coefficients of  $F$ . Then*

$$a(T) \ll_{\varepsilon, F} (\det T)^{k/2 - 3/4 + \varepsilon} \quad (\varepsilon > 0).$$

Concerning norms of Fourier-Jacobi coefficients, one optimistically may hope for the truth of the following

CONJECTURE 2. *Let  $F$  be a cusp form of integral weight  $k$  on  $\Gamma_2$  and denote by  $\phi_m$  ( $m \in \mathbf{N}$ ) its Fourier-Jacobi coefficients. Then*

$$(16) \quad \|\phi_m\| \ll_{\varepsilon, F} m^{k/2 - 1/2 + \varepsilon} \quad (\varepsilon > 0).$$

Note that – in view of Theorem 1 of §3 – (17) would be best possible.

#### NOTE ADDED IN PROOF

The estimates for the Fourier coefficients of cusp forms of arbitrary genus  $n \geq 2$  obtained in [7] improve upon those obtained in [5, 11], cf. 4.2. ii).

#### REFERENCES

- [1] ANDRIANOV, A. N. Euler products corresponding to Siegel modular forms of genus 2. *Russ. Math. Surv.* 29 (1974), 45-116.
- [2] ——— Modular descent and the Saito-Kurokawa conjecture. *Invent. math.* 53 (1979), 267-280.
- [3] BERNDT, R. On automorphic forms for the Jacobi group. To appear in *Jahresbericht der Deutschen Mathematikervereinigung*.
- [4] BÖCHERER, S. *Bemerkungen über die Dirichletreihen von Koecher und Maaß*. Math. Gott. vol. 68, Göttingen 1986.
- [5] BÖCHERER, S. and S. RAGHAVAN. On Fourier coefficients of Siegel modular forms. *J. reine angew. Math.* 384 (1988), 80-101.