

§5. Thompson's group T

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **42 (1996)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **23.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

THEOREM 4.10. *Thompson's group F is not an elementary amenable group.*

Proof. According to (a) of Chou's Proposition 2.2, it suffices to prove that $F \notin EG_\alpha$ for every ordinal α . Since EG_0 consists of finite groups and Abelian groups, it is clear that $F \notin EG_0$, so assume that $\alpha > 0$ and that $F \notin EG_\beta$ for every ordinal $\beta < \alpha$.

If α is a limit ordinal, then there is nothing to prove. Suppose that α is not a limit ordinal. It must be shown that F cannot be constructed from groups in $EG_{\alpha-1}$ as a group extension or as a direct union.

First consider group extensions. Suppose that F contains a normal subgroup N such that $N, F/N \in EG_{\alpha-1}$. Since $F \notin EG_{\alpha-1}$, N is nontrivial. Theorem 4.3 implies that $[F, F] \subset N$. Now Theorem 4.1 and Lemma 4.4 easily imply that N contains a subgroup isomorphic with F . Proposition 2.1 of [C] states that subgroups of groups in $EG_{\alpha-1}$ are also in $EG_{\alpha-1}$. Thus $F \in EG_{\alpha-1}$, contrary to hypothesis. This proves that F cannot be constructed from $EG_{\alpha-1}$ as a group extension.

Second consider direct unions. Suppose that F is a direct union of groups in $EG_{\alpha-1}$. This is clearly impossible because F is finitely generated.

This proves Theorem 4.10. \square

We next show that F is a totally ordered group (this also follows from [BriS]). Define the set of *order positive* elements of F to be the set P of functions $f \in F$ such that there exists a subinterval $[a, b]$ of $[0, 1]$ on which the derivative of f is less than 1 and $f(x) = x$ for $0 \leq x \leq a$. It is easy to see that the positive elements of F are indeed order positive. It is clear that $F = P^{-1} \cup \{1\} \cup P$. It is easy to see that P is closed under multiplication and $f^{-1}Pf \subset P$ for every $f \in F$. This proves Theorem 4.11.

THEOREM 4.11. *Thompson's group F is a totally ordered group.*

§5. THOMPSON'S GROUP T

The material in this section is mainly from unpublished notes of Thompson [T1].

Consider S^1 as the interval $[0, 1]$ with the endpoints identified. Then T is the group of piecewise linear homeomorphisms from S^1 to itself that map images of dyadic rational numbers to images of dyadic rational numbers

and that are differentiable except at finitely many images of dyadic rational numbers and on intervals of differentiability the derivatives are powers of 2. Just as we proved that F is a group, it is easy to see that T is indeed a group.

While T is defined as a group of piecewise linear homeomorphisms of S^1 , Ghys and Sergiescu [GhS] proved that there is a homeomorphism of S^1 that conjugates it to a group of C^∞ diffeomorphisms. (Thurston had proved earlier that T has a representation as a group of C^∞ diffeomorphisms of S^1 .)

EXAMPLE 5.1. The elements A and B of F induce elements of T , which will still be denoted by A and B . A third element of T is the function C defined (on $[0, 1]$) by

$$C(x) = \begin{cases} \frac{x}{2} + \frac{3}{4}, & 0 \leq x \leq \frac{1}{2} \\ 2x - 1, & \frac{1}{2} \leq x \leq \frac{3}{4} \\ x - \frac{1}{4}, & \frac{3}{4} \leq x \leq 1. \end{cases}$$

We can associate tree diagrams and unique reduced tree diagrams to elements of T almost exactly as we did to elements of F . The only difference is the following. Elements of F map leftmost leaves of domain trees to leftmost leaves of range trees. When an element of T does not do this, we denote the image in its range tree of the leftmost leaf of its domain tree with a small circle. For example, the reduced tree diagram for C is in Figure 11.




FIGURE 11

The reduced tree diagram for C

LEMMA 5.2. *The elements A , B , and C generate T and satisfy the following relations:*

- 1) $[AB^{-1}, A^{-1}BA] = 1$,
- 2) $[AB^{-1}, A^{-2}BA^2] = 1$,
- 3) $C = B(A^{-1}CB)$,
- 4) $(A^{-1}CB)(A^{-1}BA) = B(A^{-2}CB^2)$,
- 5) $CA = (A^{-1}CB)^2$, and
- 6) $C^3 = 1$.

Proof. Let H be the subgroup of T generated by $\{A, B, C\}$. Since $\{A, B\}$ is a generating set for F , $F \subset H$. Suppose $f \in T$. Let $[x] = f([0])$. If $[x] = [0]$, then $f \in F$ and hence $f \in H$. If $[x] \neq [0]$, then there is an element $h \in F$ with $h(x) = \frac{3}{4}$ by Lemma 4.2. Then $g = C^{-1}hf$ fixes $[0]$, so $g \in F$. Hence $f = h^{-1}Cg \in H$ and $H = T$. Thus A, B , and C generate T .

Relations 1) and 2) are proved in Section 3.

Consider relation 3). It is equivalent to the relation $CBC^{-1} = AB^{-1}$. The reduced tree diagram for CBC^{-1} is computed in Figure 12, the notation being straightforward.




FIGURE 12

Computing the reduced tree diagram for CBC^{-1}

Referring to Figure 1 shows that AB^{-1} has the same reduced tree diagram as CBC^{-1} , which completes the verification of relation 3).

Consider relation 4). It is equivalent to

$$(B^{-1}C^{-1}A)(AB^{-1})(A^{-1}CB) = BA^{-1}B^{-1}A,$$

where the term $A^{-1}CB$ here corresponds to the same term in relation 4). We compute a tree diagram for the left side of this equation in Figure 13.




FIGURE 13

Computing a tree diagram for $(B^{-1}C^{-1}A)(AB^{-1})(A^{-1}CB)$

Referring to Figure 1 now completes the verification of relation 4).

Relation 6) is easily verified using the reduced tree diagram for C .

Finally consider relation 5). Use relation 6) and then relation 3) to rewrite relation 5): $CA = A^{-1}CBA^{-1}CB \Leftrightarrow CA = A^{-1}C^{-1}(C^{-1}BA^{-1}CB) \Leftrightarrow CA = A^{-1}C^{-1} \Leftrightarrow (AC)^2 = 1$. The reduced tree diagram for AC is computed in Figure 14.




FIGURE 14

Computing the reduced tree diagram for AC

Hence AC acts on S^1 by translation by $[\frac{1}{2}]$, and so $(AC)^2 = 1$, which gives relation 5). \square

Let

$$T_1 = \langle A, B, C : [AB^{-1}, A^{-1}BA], [AB^{-1}, A^{-2}BA^2], C^{-1}B(A^{-1}CB), \\ ((A^{-1}CB)(A^{-1}BA))^{-1}B(A^{-2}CB^2), (CA)^{-1}(A^{-1}CB)^2, C^3 \rangle.$$

LEMMA 5.3. *There is a surjection $T_1 \rightarrow T$ that maps the formal symbols A , B , and C to the functions A , B , and C in T .*

Proof. This follows immediately since the functions A , B , and C satisfy the relations 1) - 6). \square

LEMMA 5.4. *The subgroup of T_1 generated by A and B is isomorphic to F .*

Proof. The results of Section 3 show that there exists a group homomorphism from F to the subgroup of T_1 generated by A and B whose composition with the map from T_1 to T is the identity map on F . This proves Lemma 5.4. \square

It is easier at this point to prove that T is simple than to prove that T_1 is simple. However, it is preferable to prove that T_1 is simple, since then Lemma 5.3 implies that T is isomorphic to T_1 .

Define the elements X_n , $n \geq 0$, of T_1 by $X_0 = A$ and $X_n = A^{-(n-1)}BA^{n-1}$ for $n \geq 1$. It follows from Theorem 3.4 and Lemma 5.4 that $X_nX_k = X_kX_{n+1}$ if $k < n$. Define the elements C_n , $n \geq 1$, of T_1 by $C_n = A^{-(n-1)}CB^{n-1}$. For convenience we define $C_0 = 1$.

To gain some insight into these elements C_n , in Figure 15 we calculate reduced tree diagrams for the corresponding elements, still called C_n , in T . The reduced tree diagram for C_1 is given in Figure 11, and the reduced tree diagram for C_2 is given in Figure 13. This calculation shows that C_n permutes the images of the $n + 2$ intervals

$$[0, 1 - 2^{-1}], [1 - 2^{-1}, 1 - 2^{-2}],$$

$$[1 - 2^{-2}, 1 - 2^{-3}], \dots, [1 - 2^{-n}, 1 - 2^{-(n+1)}], [1 - 2^{-(n+1)}, 1]$$

cyclically.

The rest of this section deals with the group T_1 .




FIGURE 15

Inductively computing the reduced tree diagram for C_n with $n \geq 3$

LEMMA 5.5. *If k, n are positive integers and $k \leq n$, then*

- i) $C_n = X_n C_{n+1}$,
- ii) $C_n X_k = X_{k-1} C_{n+1}$, and
- iii) $C_n A = C_{n+1}^2$.

Proof.

$$C_n = A^{-(n-1)} C B^{n-1} = A^{-(n-1)} B (A^{-1} C B) B^{n-1}$$

$$= (A^{-(n-1)} B A^{n-1}) (A^{-n} C B^n) = X_n C_{n+1},$$

which proves i).

If $k = 1$, ii) follows from the definition. If $k = n = 2$, ii) follows from relation 4). If $k = 2$ and $n > 2$, then by induction on n

$$C_n X_2 \stackrel{i)}{=} X_{n-1}^{-1} C_{n-1} X_2 \stackrel{\text{induct}}{=} X_{n-1}^{-1} X_1 C_n = X_1 X_n^{-1} C_n \stackrel{i)}{=} X_1 C_{n+1}.$$

If $k \geq 3$, then by induction on k

$$C_n X_k = A^{-1} C_{n-1} B X_k$$

$$= A^{-1} C_{n-1} X_{k-1} B \stackrel{\text{induct}}{=} A^{-1} X_{k-2} C_n B = (A^{-1} X_{k-2} A) (A^{-1} C_n B)$$

$$= X_{k-1} C_{n+1}.$$

Equation iii) follows by induction on n . If $n = 1$ then it is relation 5). If $n > 1$, then

$$C_n A = A^{-1} C_{n-1} B A = A^{-1} C_{n-1} A X_2 \\ \stackrel{\text{induct}}{=} A^{-1} C_n^2 X_2 \stackrel{\text{ii)}}{=} A^{-1} C_n B C_{n+1} = C_{n+1}^2. \quad \square$$

LEMMA 5.6. *If n is a positive integer, $m \in \{1, \dots, n+1\}$, and $r, s \in \{0, \dots, n\}$, then*

i)

$$C_n^m X_r = \begin{cases} X_{r-m} C_{n+1}^m, & r \geq m \\ C_{n+1}^{m+1}, & r = m - 1 \\ X_{r+(n+2-m)} C_{n+1}^{m+1}, & r < m - 1; \end{cases}$$

ii)

$$X_s^{-1} C_n^m = \begin{cases} C_{n+1}^{m+1} X_{(s+m)-(n+2)}^{-1}, & s \geq (n+2) - m \\ C_{n+1}^m, & s = (n+1) - m \\ C_{n+1}^m X_{s+m}^{-1}, & s \leq n - m; \end{cases}$$

$$\text{iii)} \quad C_n^m = X_{(n+1)-m} C_{n+1}^m;$$

$$\text{iv)} \quad C_n^m = C_{n+1}^{m+1} X_{m-1}^{-1};$$

$$\text{v)} \quad C_n^{n+2} = 1.$$

Proof. The first line of i) follows from Lemma 5.5.ii). If $r = 0$, the second line is Lemma 5.5.iii); if $r > 0$

$$C_n^m X_r = C_n C_n^r X_r \stackrel{5.5.ii)}{=} C_n A C_{n+1}^r \stackrel{5.5.iii)}{=} C_{n+1}^2 C_{n+1}^r = C_{n+1}^{m+1}.$$

This proves i) if $r \geq m - 1$.

$$C_n^m = C_n^{m-1} C_n \stackrel{5.5.i)}{=} C_n^{m-1} X_n C_{n+1} \stackrel{5.5.ii)}{=} X_{n-(m-1)} C_{n+1}^m,$$

which proves iii). If $r < m - 1$, then

$$C_n^m X_r = C_n^{m-(r+1)} C_n^{r+1} X_r \stackrel{5.5}{=} C_n^{m-(r+1)} C_{n+1}^{r+2} \\ \stackrel{\text{iii)}}{=} X_{n+1-(m-(r+1))} C_{n+1}^{m-(r+1)} C_{n+1}^{r+2} = X_{r+(n+2-m)} C_{n+1}^{m+1},$$

which finishes the proof for i).

The first line of ii) follows from i), with $s = r + (n+2-m)$. The second line of ii) follows from iii). The third line of ii) follows from i), with $s = r - m$. Equation iv) follows from the second line of i). If $t \geq 1$, then

$$C_t^{t+2} = C_t C_t^{t+1} \stackrel{\text{iii)}}{=} C_t A C_{t+1}^{t+1} \stackrel{5.5.iii)}{=} C_{t+1}^2 C_{t+1}^{t+1} = C_{t+1}^{t+3}.$$

Since $C_1^3 = C^3 = 1$, this proves v) and completes the proof of Lemma 5.6. \square

Following the terminology for F , an element of T_1 which is a product of nonnegative powers of the X_i 's will be called *positive* and an inverse of a positive element will be called *negative*.

THEOREM 5.7. *If $g \in T_1$, then $g = pC_n^m q^{-1}$ for some positive elements p, q and nonnegative integers m, n with $m < n + 2$.*

Proof. We first show that if i, j, k , and l are positive integers, then there are positive elements p and q and nonnegative integers m and n such that $C_j^i C_l^k = pC_n^m q^{-1}$. Suppose that i, j, k , and l are positive integers and that $g = C_j^i C_l^k$. Since $C_j^{j+2} = C_l^{l+2} = 1$ by Lemma 5.6.v), we can assume that $i < j + 2$ and $k < l + 2$. Let $n \geq \max\{j, l\}$. By Lemma 5.6.iii) and Lemma 5.6.iv), there is a positive integer r and there are positive elements p and q such that $C_j^i = pC_n^i$ and $C_l^k = C_n^r q^{-1}$. Hence $C_j^i C_l^k = pC_n^{i+r} q^{-1}$.

Let $H = \{g \in T_1 : g = pC_n^m q^{-1} \text{ for some positive elements } p, q, \text{ and nonnegative integers } m, n \text{ with } m < n + 2\}$. Lemma 5.6.v) easily implies that H is closed under inversion. To show that H is closed under multiplication, suppose that $g_1, g_2 \in H$. Then $g_1 = p_1 C_j^i q_1^{-1}$ and $g_2 = p_2 C_l^k q_2^{-1}$ for some positive elements p_1, p_2, q_1 , and q_2 and some nonnegative integers i, j, k , and l with $i < j + 2$ and $k < l + 2$. By Corollary 2.7, there are positive elements p_3 and q_3 such that $q_1^{-1} p_2 = p_3 q_3^{-1}$. Hence $g_1 g_2 = p_1 C_j^i q_1^{-1} p_2 C_l^k q_2^{-1} = p_1 C_j^i p_3 q_3^{-1} C_l^k q_2^{-1}$. Lemma 5.6.iii) and Lemma 2.8, which states that the set of positive elements of F is closed under multiplication, show that if $i > 0$ and $j > 0$, then we may replace C_j^i by C_{j+1}^i . Hence we may assume that if $i > 0, j > 0$, and X_r occurs in p_3 , then $j \geq r$. We may likewise assume that if $k > 0, l > 0$, and X_s occurs in q_3 , then $l \geq s$. Now Lemmas 5.6.i), 5.6.ii), and 2.8 show that there are positive elements p_4 and q_4 and nonnegative integers r, s, t , and u such that $g_1 g_2 = p_4 C_s^r C_u^t q_4^{-1}$. By the previous paragraph and Lemma 2.8, there are positive elements p_5 and q_5 and nonnegative integers m and n such that $g_1 g_2 = p_5 C_n^m q_5^{-1}$. Since we can assume that $m < n + 2$ by Lemma 5.6.v), $g_1 g_2 \in H$. Hence H is a subgroup of T_1 . Since T_1 is generated by $A = X_0, B = X_1$, and $C = C_1$, all of which are in $H, H = T_1. \quad \square$

THEOREM 5.8. *T_1 is simple.*

Proof. Suppose N is a nontrivial normal subgroup of T_1 , and let $\theta: T_1 \rightarrow T_1/N$ be the quotient homomorphism. Then there is an element $g \in T_1$ with $g \neq 1$ and $\theta(g) = 1$. By Theorem 5.7, $g = pC_n^m q^{-1}$ for some positive elements p, q and nonnegative integers m, n with $m < n + 2$. Then

$$\theta(p^{-1}q) = \theta(C_n^m)$$

$$\text{and } \theta((p^{-1}q)^{n+2}) = \theta((C_n^m)^{n+2}) = \theta((C_n^{n+2})^m) \stackrel{5.6.v)}{=} \theta(1) = 1.$$

By Lemma 5.4, there is a homomorphism $\alpha: F \rightarrow T_1/N$ defined on generators by $\alpha(A) = \theta(A)$ and $\alpha(B) = \theta(B)$. If $p^{-1}q \neq 1$, then $(p^{-1}q)^{n+2} \neq 1$, and so $\alpha(F)$ is a proper quotient group of F . Since every proper quotient group of F is Abelian by Theorem 4.3, $\theta(AB) = \theta(BA)$. If $p^{-1}q = 1$, then $m, n > 0$ and $1 = \theta(C_n^m) \stackrel{5.6.iv)}{=} \theta(C_{n+1}^{m+1})\theta(X_{m-1}^{-1})$ and hence $\theta(X_{m-1}^{n+3}) = \theta((C_{n+1}^{m+1})^{n+3}) = \theta((C_{n+1}^{n+3})^{m+1}) = \theta(1) = 1$. It follows as before that $\theta(AB) = \theta(BA)$. Hence $\theta(A^{-1}BA) = \theta(B)$, so $\theta(A^{-1}C) = \theta(BA^{-2}C)$ by relation 4). Hence $\theta(BA^{-1}) = 1$, and so $\theta(B) = 1$ by relation 3). This implies that $\theta(A) = 1$. It now follows from relation 5) that $\theta(C) = 1$. Thus $N = T_1$, and so T_1 is simple. \square

COROLLARY 5.9. T_1 is isomorphic to T .

§6. THOMPSON'S GROUP V

As with the previous section, the material in this section is mainly from unpublished notes of Thompson [T1]; [T1] contains the statements of the lemmas (except for Lemma 6.2) and the statement and proof of Theorem 6.9, but does not contain the proofs of the lemmas.

Let V be the group of right-continuous bijections of S^1 that map images of dyadic rational numbers to images of dyadic rational numbers, that are differentiable except at finitely many images of dyadic rational numbers, and such that, on each maximal interval on which the function is differentiable, the function is linear with derivative a power of 2. As before, it is easy to prove that V is a group.

We can associate tree diagrams with elements of V as we did for F and T , except that now we need to label the leaves of the domain and range trees to indicate the correspondence between the leaves. For example, reduced tree diagrams for A , B , and C are given in Figure 16.

Using the identification of S^1 as the quotient of $[0, 1]$, define $\pi_0: S^1 \rightarrow S^1$ by

$$\pi_0(x) = \begin{cases} \frac{x}{2} + \frac{1}{2}, & 0 \leq x < \frac{1}{2} \\ 2x - 1, & \frac{1}{2} \leq x < \frac{3}{4} \\ x, & \frac{3}{4} \leq x < 1. \end{cases}$$