

3. Uniformly exponential growth and growth of graded algebras

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **43 (1997)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **09.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

3. UNIFORMLY EXPONENTIAL GROWTH AND GROWTH OF GRADED ALGEBRAS

In this section we describe a method of estimating growth functions of a group in terms of its graded Lie, and associative algebras defined via dimension subgroups. We begin by recalling some concepts and notations.

As in [Gri] considerations were given with respect to a Galois field \mathbf{GF}_p , here we modify the arguments for a field of characteristic 0, namely \mathbf{Q} .

Let G be a group; denote by $\mathbf{Q}[G]$ the group algebra of G over \mathbf{Q} , and by $\Delta \subset \mathbf{Q}[G]$ the augmentation ideal, that is the ideal generated by the elements of the form $g - 1$, with $g \in G$. Recall that the *lower central series* of G is the sequence of subgroups $\{\gamma_n(G)\}_{n=1}^{\infty}$ of G defined by $\gamma_1(G) = G$ and, for $n \geq 2$, $\gamma_n(G) = [G, \gamma_{n-1}(G)]$.

The subgroup

$$G_n = \{g \in G : g - 1 \in \Delta^n\}$$

is called the *n-th dimension subgroup* of G over \mathbf{Q} and it has the following characterisation due to Jennings [J] (see also [P: IV, Thm. 1.5] or [Pm: 11, Thm. 1.10])

$$G_n = \sqrt{\gamma_n(G)} := \{g \in G : \exists k \in \mathbf{N}, g^k \in \gamma_n(G)\}.$$

For any group G one defines as usual an associative graded algebra $\mathcal{A}(G)$ and two graded Lie algebras $L(G)$ and $\mathcal{L}(G)$ by

$$\begin{aligned}\mathcal{A}(G) &= \bigoplus_{n=1}^{\infty} \Delta^n / \Delta^{n+1} \\ L(G) &= \bigoplus_{n=1}^{\infty} [(G_n / G_{n+1}) \otimes_{\mathbf{Z}} \mathbf{Q}] \\ \mathcal{L}(G) &= \bigoplus_{n=1}^{\infty} [(\gamma_n(G) / \gamma_{n+1}(G)) \otimes_{\mathbf{Z}} \mathbf{Q}]\end{aligned}$$

(see for instance [P], [Pm]). Quillen's Theorem [Q] states that $\mathcal{A}(G)$ is the universal enveloping algebra of $L(G)$.

Assume now that G is finitely generated and set

$$\begin{aligned}a_n(G) &= \dim(\Delta^n / \Delta^{n+1}) \\ b_n(G) &= \text{rank}(G_n / G_{n+1}) \\ c_n(G) &= \text{rank}(\gamma_n(G) / \gamma_{n+1}(G))\end{aligned}$$

where, by rank, we mean the torsion free rank of the corresponding abelian group. Then the following relations hold

$$\sum_{n=0}^{\infty} a_n(G)z^n = \prod_{n=1}^{\infty} (1 - z^n)^{-b_n(G)} = \prod_{n=1}^{\infty} (1 - z^n)^{-c_n(G)}.$$

The first equality follows easily from Quillen's Theorem [Pm: Thm. 4.10, Chapter 3] and the second one follows from the equality $b_n(G) = c_n(G)$ as proved in [Be].

In [Be] it is also proved that

$$\limsup_{n \rightarrow \infty} \sqrt[n]{a_n} = \limsup_{n \rightarrow \infty} \sqrt[n]{c_n}.$$

3.1. LEMMA. *For any finite system of generators A of a group G the following inequality holds:*

$$a_n(G) \leq \gamma_A^G(n), \quad n \geq 1.$$

Proof. For $x, y \in G$ we have

$$\begin{aligned} xy - 1 &= (x - 1) + (y - 1) + (x - 1)(y - 1) \\ x^{-1} - 1 &= -(x - 1) - (x - 1)(x^{-1} - 1) \end{aligned}$$

so that

$$\begin{aligned} xy - 1 &\equiv (x - 1) + (y - 1) \pmod{\Delta^2} \\ x^{-1} - 1 &\equiv -(x - 1) \pmod{\Delta^2}. \end{aligned}$$

The ideal Δ^n is spanned, over \mathbf{Q} , by the elements of the form

$$y_1(x_1 - 1)y_2(x_2 - 1) \cdots y_n(x_n - 1)y_{n+1},$$

where $x_i \in G$ and $y_j \in \mathbf{Q}[G]$, $1 \leq i \leq n$, $1 \leq j \leq n+1$. Since

$$y = \sum_{g \in G} k_g g \equiv \sum_{g \in G} k_g \pmod{\Delta}, \quad k_g \in \mathbf{Q}$$

a basis for the quotient space Δ^n/Δ^{n+1} can be chosen among the images modulo Δ^{n+1} of the elements of the form

$$(a_{i_1} - 1)(a_{i_2} - 1) \cdots (a_{i_n} - 1),$$

where $a_{i_j} \in A$. But $(a_{i_1} - 1)(a_{i_2} - 1) \cdots (a_{i_n} - 1) = \sum_{g \in G} k'_g g$, where the summation extends over elements g of length at most n with respect to the system of generators A . \square

3.2. COROLLARY. *Let G be a finitely generated group and suppose that the ranks of $\gamma_n(G)/\gamma_{n+1}(G)$ grow exponentially. Then G has uniformly exponential growth and the estimate*

$$\lambda_*(G) \geq \limsup_{n \rightarrow \infty} \sqrt[n]{\text{rank}(\gamma_n(G)/\gamma_{n+1}(G))}$$

holds.

Recall that a group G is *parafree of para-rank m* if it is residually nilpotent and the factors of consecutive groups in its lower central series equal the corresponding ones of a free group of rank m . There are parafree groups which are not isomorphic to free groups [B 2,3].

3.3. PROPOSITION. *A finitely generated parafree group G of para-rank $m \geq 2$ has uniformly exponential growth and*

$$\lambda_*(G) \geq m.$$

Proof. It is known (see for instance [MKS : Thms. 5.11 (Witt's Formulae) and 5.12]) that for a free group \mathbf{F}_m the rank of $(\gamma_n(\mathbf{F}_m)/\gamma_{n+1}(\mathbf{F}_m))$ equals the n -th coefficient of the Maclaurin power series of the function $U(z) = 1/(1-mz)$ and the previous corollary can be applied. \square

Given a parafree group G of para-rank $m \geq 2$ it would be interesting to compare $\lambda_*(G)$ with $\lambda_*(\mathbf{F}_m) = 2m - 1$.

3.4. PROBLEM. *Is it true that, for a finitely generated para-free group G of para-rank $m \geq 2$ which is not free, one has $\lambda_*(G) > 2m - 1$?*

In order to formulate the next statement we recall the following

3.5. DEFINITION. An element $R \in F$ is said to be *primitive with respect to the lower central series* if, for all $n \geq 2$, it is not an n -th power modulo $\gamma_{\omega(R)+1}(F)$ where $\omega(R)$ is the weight of R . (The latter is defined by $R \in \gamma_{\omega(R)}(F)$ but $R \notin \gamma_{\omega(R)+1}(F)$.)

3.6. THEOREM ([L 1,2]). *Let R be an element of the free group F of finite rank m which is primitive with respect to the lower central series. Denote by $k = \omega(R)$ its weight and by $\langle R \rangle$ the normal closure of R in F . Let $G = F/\langle R \rangle$ and let $\mathcal{L}(F)$ and $\mathcal{L}(G)$ be the corresponding Lie algebras. Let then r be the image of R in $\mathcal{L}_k(F)$, the k -th component of $\mathcal{L}(F)$ and denote by I the ideal of $\mathcal{L}(F)$ generated by r .*

Then I is the kernel of the canonical homomorphism of $\mathcal{L}(F)$ onto $\mathcal{L}(G)$, i.e.

$$\mathcal{L}(G) = \mathcal{L}(F)/I.$$

Moreover for all $n \geq 1$ the abelian group $\mathcal{L}_n(G)$ is a torsion free group whose rank is the n -th coefficient of the Maclaurin power series of the function

$$U(z) = \frac{1}{1 - mz + z^k}.$$

4. MORE ON UNIFORMLY EXPONENTIAL GROWTH OF ONE-RELATOR GROUPS

Any two-generated one-relator group G can be presented in the form $G = \langle a, b : a^k w(a, b) = 1 \rangle$ where $k \in \mathbf{Z}$ and $w(a, b)$ belongs to the commutator subgroup $[F, F]$ of the free group $F = F(a, b)$ freely generated by a and b (this follows from Lemma 1.1). Since a and b constitute a basis in $F/\gamma_2(F)$ and $[a, b]$ generates $\gamma_2(F)/\gamma_3(F)$, one can also present G in the form

$$G = \langle a, b : a^k [a, b]^l w(a, b) = 1 \rangle$$

where $k, l \in \mathbf{Z}$ and $w(a, b) \in \gamma_3(F)$.

In this section we shall see that, under suitable assumptions on k, l and $w(a, b)$, the corresponding group has uniformly exponential growth.

As an application of Labute's Theorem we get the following:

4.1. PROPOSITION. *Let $G = \langle a, b : R(a, b) = 1 \rangle$ be such that R is primitive with respect to $\{\gamma_n(F)\}_{n=1}^\infty$ and $R \in \gamma_3(F)$. Then G has uniformly exponential growth.*

Proof. If $\omega(R) \geq 3$, Theorem 3.6 shows that the corresponding function $U(z)$ has a pole z_0 with $0 < z_0 < 1$. It follows that the coefficients $c_n(G)$ grow exponentially. By Corollary 3.2, $\lambda_*(G) > 1$. \square