

# 7. Arithmetic intersection theory

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **43 (1997)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **22.09.2024**

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## 7. ARITHMETIC INTERSECTION THEORY

We recall here the generalization of Arakelov theory to higher dimensions due to Gillet and Soulé. Our main references are [GS1], [GS2] and the exposition in [SABK]. For  $A$  an abelian group,  $A_{\mathbf{Q}}$  denotes  $A \otimes_{\mathbf{Z}} \mathbf{Q}$ . Let  $X$  be an *arithmetic scheme over  $\mathbf{Z}$* , by which we mean a regular scheme, projective and flat over  $\text{Spec } \mathbf{Z}$ . For  $p \geq 0$ , let  $X^{(p)}$  be the set of integral subschemes of  $X$  of codimension  $p$  and  $Z^p(X)$  be the group of codimension  $p$  cycles on  $X$ . The  $p$ -th Chow group of  $X$ :  $CH^p(X) := Z^p(X)/R^p(X)$ , where  $R^p(X)$  is the subgroup of  $Z^p(X)$  generated by the cycles  $\text{div } f$ ,  $f \in k(x)^*$ ,  $x \in X^{(p-1)}$ . Let  $CH(X) = \bigoplus_p CH^p(X)$ . If  $X$  is smooth over  $\text{Spec } \mathbf{Z}$ , then the methods of [F] can be used to give  $CH(X)$  the structure of a commutative ring. In general one has a product structure on  $CH(X)_{\mathbf{Q}}$  after tensoring with  $\mathbf{Q}$ .

Let  $D^{p,p}(X(\mathbf{C}))$  denote the space of complex currents of type  $(p,p)$  on  $X(\mathbf{C})$ , and  $F_{\infty} : X(\mathbf{C}) \rightarrow X(\mathbf{C})$  the involution induced by complex conjugation. Let  $D^{p,p}(X_{\mathbf{R}})$  (resp.  $A^{p,p}(X_{\mathbf{R}})$ ) be the subspace of  $D^{p,p}(X(\mathbf{C}))$  (resp.  $A^{p,p}(X(\mathbf{C}))$ ) generated by real currents (resp. forms)  $T$  such that  $F_{\infty}^* T = (-1)^p T$ ; denote by  $\tilde{D}^{p,p}(X_{\mathbf{R}})$  and  $\tilde{A}^{p,p}(X_{\mathbf{R}})$  the respective images in  $\tilde{D}^{p,p}(X(\mathbf{C}))$  and  $\tilde{A}^{p,p}(X(\mathbf{C}))$ .

An *arithmetic cycle* on  $X$  of codimension  $p$  is a pair  $(Z, g_Z)$  in the group  $Z^p(X) \oplus \tilde{D}^{p-1,p-1}(X_{\mathbf{R}})$ , where  $g_Z$  is a *Green current* for  $Z(\mathbf{C})$ , i.e. a current such that  $dd^c g_Z + \delta_{Z(\mathbf{C})}$  is represented by a smooth form. The group of arithmetic cycles is denoted by  $\widehat{Z}^p(X)$ . If  $x \in X^{(p-1)}$  and  $f \in k(x)^*$ , we let  $\widehat{\text{div}} f$  denote the arithmetic cycle  $(\text{div } f, [-\log |f_{\mathbf{C}}|^2 \cdot \delta_{x(\mathbf{C})}])$ .

The  $p$ -th *arithmetic Chow group* of  $X$ :  $\widehat{CH}^p(X) := \widehat{Z}^p(X)/\widehat{R}^p(X)$ , where  $\widehat{R}^p(X)$  is the subgroup of  $\widehat{Z}^p(X)$  generated by the cycles  $\widehat{\text{div}} f$ ,  $f \in k(x)^*$ ,  $x \in X^{(p-1)}$ . Let  $\widehat{CH}(X) = \bigoplus_p \widehat{CH}^p(X)$ .

We have the following canonical morphisms of abelian groups:

$$\begin{aligned} \zeta : \widehat{CH}^p(X) &\longrightarrow CH^p(X), & [(Z, g_Z)] &\longmapsto [Z], \\ \omega : \widehat{CH}^p(X) &\longrightarrow \text{Ker } d \cap \text{Ker } d^c (\subset A^{p,p}(X_{\mathbf{R}})), & [(Z, g_Z)] &\longmapsto dd^c g_Z + \delta_{Z(\mathbf{C})}, \\ a : \tilde{A}^{p-1,p-1}(X_{\mathbf{R}}) &\longrightarrow \widehat{CH}^p(X), & \eta &\longmapsto [(0, \eta)]. \end{aligned}$$

One can define a pairing  $\widehat{CH}^p(X) \otimes \widehat{CH}^q(X) \rightarrow \widehat{CH}^{p+q}(X)_{\mathbf{Q}}$  which turns  $\widehat{CH}(X)_{\mathbf{Q}}$  into a commutative graded unitary  $\mathbf{Q}$ -algebra. The maps  $\zeta$ ,  $\omega$  are  $\mathbf{Q}$ -algebra homomorphisms. If  $X$  is smooth over  $\mathbf{Z}$  one does not have to tensor with  $\mathbf{Q}$ . The definition of this pairing is difficult; the construction uses the *star product* of Green currents, which in turn relies upon Hironaka's

resolution of singularities to get to the case of divisors. The functor  $\widehat{CH}^p(X)$  is contravariant in  $X$ , and covariant for proper maps which are smooth on the generic fiber.

Choose a Kähler form  $\omega_0$  on  $X(\mathbf{C})$  such that  $F_\infty^* \omega_0 = -\omega_0$  (this is equivalent to requiring that the corresponding Kähler metric is invariant under  $F_\infty$ ). It is natural to utilize the theory of harmonic forms on  $X$  in the study of Green currents on  $X(\mathbf{C})$ . Following [GS1], we call the pair  $\bar{X} = (X, \omega_0)$  an *Arakelov variety*. By the Hodge decomposition theorem, we have  $A^{p,p}(X_{\mathbf{R}}) = \mathcal{H}^{p,p}(X_{\mathbf{R}}) \oplus \text{Im } d \oplus \text{Im } d^*$ , where  $\mathcal{H}^{p,p}(X_{\mathbf{R}}) = \text{Ker } \Delta \subset A^{p,p}(X)$  denotes the space of harmonic (with respect to  $\omega_0$ )  $(p, p)$  forms  $\alpha$  on  $X(\mathbf{C})$  such that  $F_\infty^* \alpha = (-1)^p \alpha$ . The subgroup  $CH^p(\bar{X}) := \omega^{-1}(\mathcal{H}^{p,p}(X_{\mathbf{R}}))$  of  $\widehat{CH}^p(X)$  is called the  $p$ -th *Arakelov Chow group of  $X$* . Let  $CH(\bar{X}) = \bigoplus_{p \geq 0} CH^p(\bar{X})$ .  $CH^p(\bar{X})$  is a direct summand of  $\widehat{CH}^p(X)$ , and there is an exact sequence

$$(11) \quad CH^{p,p-1}(X) \xrightarrow{\rho} \mathcal{H}^{p-1,p-1}(X_{\mathbf{R}}) \xrightarrow{a} CH^p(\bar{X}) \xrightarrow{\zeta} CH^p(X) \longrightarrow 0.$$

In the above sequence the group  $CH^{p,p-1}(X)$  is defined as the  $E_2^{p,1-p}$  term of a certain spectral sequence used by Quillen to calculate the higher algebraic  $K$ -theory of  $X$ , and the map  $\rho$  coincides with the Beilinson regulator map (cf. [G] and [GS1], 3.5).

If  $\mathcal{H}(X_{\mathbf{R}}) = \bigoplus_p \mathcal{H}^{p,p}(X_{\mathbf{R}})$  is a subring of  $\bigoplus_p A^{p,p}(X_{\mathbf{R}})$ , for example if  $X(\mathbf{C})$  is a curve, an abelian variety or a hermitian symmetric space (e.g. a Grassmannian), then  $CH(\bar{X})_{\mathbf{Q}}$  is a subring of  $\widehat{CH}(X)_{\mathbf{Q}}$ . This is not the case in general; for example it fails to be true for the complete flag varieties.

Arakelov [A] introduced the group  $CH^1(\bar{X})$ , where  $\bar{X} = (X, g_0)$  is an arithmetic surface with the metric  $g_0$  on the Riemann surface  $X(\mathbf{C})$  given by  $\frac{i}{2g} \sum \omega_j \wedge \bar{\omega}_j$ . Here  $g$  is the genus of  $X(\mathbf{C})$  and  $\{\omega_j\}$  for  $1 \leq j \leq g$  is an orthonormal basis of the space of holomorphic one forms on  $X(\mathbf{C})$ .

A *hermitian vector bundle*  $\bar{E} = (E, h)$  on an arithmetic scheme  $X$  is an algebraic vector bundle  $E$  on  $X$  such that the induced holomorphic vector bundle  $E(\mathbf{C})$  on  $X(\mathbf{C})$  has a hermitian metric  $h$ , which is invariant under complex conjugation, i.e.  $F_\infty^*(h) = h$ .

To any hermitian vector bundle one can attach characteristic classes  $\widehat{\phi}(\bar{E}) \in \widehat{CH}(X)_{\mathbf{Q}}$ , for any  $\phi \in I(n, \mathbf{Q})$ , where  $n = \text{rk } E$ . For example, we have *arithmetic Chern classes*  $\widehat{c}_k(\bar{E}) \in \widehat{CH}^k(X)$ . Some basic properties of these classes are:

- (1)  $\widehat{c}_0(\bar{E}) = 1$  and  $\widehat{c}_p(\bar{E}) = 0$  for  $k > \text{rk } E$ .
- (2) The form  $\omega(\widehat{c}_k(\bar{E})) = c_k(\bar{E}) \in A^{k,k}(X_{\mathbf{R}})$  is the  $k$ -th Chern form of the hermitian bundle  $\overline{E(\mathbf{C})}$ .

$$(3) \quad \zeta(\widehat{c}_k(\bar{E})) = c_k(E) \in CH^k(X).$$

(4)  $f^*\widehat{c}_k(\bar{E}) = \widehat{c}_k(f^*\bar{E})$ , for every morphism  $f : X \rightarrow Y$  of regular schemes, projective and flat over  $\mathbf{Z}$ .

(5) If  $\bar{L}$  is a hermitian line bundle,  $\widehat{c}_1(\bar{L})$  is the class of  $(\text{div}(s), -\log \|s\|^2)$  for any rational section  $s$  of  $L$ .

Analogous properties are satisfied by  $\widehat{\phi}$  for any  $\phi \in I(n, \mathbf{Q})$  (see [GS2], Th. 4.1). The most relevant property of these characteristic classes is their behaviour in short exact sequences: if

$$\bar{\mathcal{E}} : 0 \rightarrow \bar{S} \rightarrow \bar{E} \rightarrow \bar{Q} \rightarrow 0$$

is such a sequence of hermitian vector bundles over  $X$ , then

$$(12) \quad \widehat{\phi}(\bar{S} \oplus \bar{Q}) - \widehat{\phi}(\bar{E}) = a(\widetilde{\phi}(\bar{\mathcal{E}})).$$

Relation (12) is the main tool for calculating intersection products of classes in  $\widehat{CH}(X)$  that come from characteristic classes of vector bundles. Combining it with the results of §4 and §5 gives immediate consequences for such intersections. For example, we have

**COROLLARY 4.** *Let  $\bar{\mathcal{E}} : 0 \rightarrow \bar{S} \rightarrow \bar{E} \rightarrow \bar{Q} \rightarrow 0$  be a short exact sequence of hermitian vector bundles over an arithmetic scheme  $X$ . Assume that the metrics on  $S(\mathbf{C})$ ,  $Q(\mathbf{C})$  are induced from that on  $E(\mathbf{C})$ .*

(a) *If  $\overline{E(\mathbf{C})}$  is flat, then*

- (1)  $\widehat{p}_\lambda(\bar{S} \oplus \bar{Q}) = \widehat{p}_\lambda(\bar{E})$ , if  $\lambda$  has length  $> 1$ , and
- (2)  $\widehat{p}_k(\bar{S}) + \widehat{p}_k(\bar{Q}) - \widehat{p}_k(\bar{E}) = k\mathcal{H}_{k-1}a(p_{k-1}(\bar{Q}))$ ,  $\forall k \geq 1$ ,

*in the arithmetic Chow group  $\widehat{CH}(X)_{\mathbf{Q}}$ .*

(b) *If  $\bar{E} = \bar{L}^{\oplus n}$  for some hermitian line bundle  $\bar{L}$  and  $\omega = c_1(\overline{L(\mathbf{C})})$ , then*

$$\widehat{c}(\bar{S})\widehat{c}(\bar{Q}) - \widehat{c}(\bar{E}) = \sum_{i,j} (-1)^j \binom{n}{i} (\mathcal{H}_n - \mathcal{H}_{n-i} + \mathcal{H}_j) a(\omega^i p_j(\bar{Q})),$$

*in the arithmetic Chow group  $\widehat{CH}(X)$ .*