

Objektyp: **ReferenceList**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **44 (1998)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **24.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Proof. Let us consider a coding of the rotation by angle α under the left-closed and right-open partition of the unit circle bounded by all the points of the form $\{n\alpha + \gamma_i\}$, for $0 \leq n < n_i$ and $1 \leq i \leq d$; let $\beta_0, \dots, \beta_{p-1}$ denote these consecutive points. The letter associated with the interval $I_k = [\beta_k, \beta_{k+1}[$ has a unique right extension, except when I_k contains points of the form $\{\beta_i - \alpha\}$. Suppose there are $q \geq 2$ points of this form; the associated letter has $q + 1$ right extensions. Since there are at most d points of this type, we obtain $p(2) - p(1) \leq d$. We deduce from Theorem 6 that there are at most $3d$ different frequencies for the letters of the coding, i.e., there are at most $3d$ different lengths for the intervals I_k .

REMARK. The start and finish intervals as introduced by Liang in his proof in [37] correspond exactly to the beginning of the branches in the graph of words. Indeed, Liang shows that any interval is associated either with a start point $\{\gamma_i\}$ (i.e., with one extension of a factor having more than one right extension) or with a finish point $\{(n_i - 1)\alpha + \gamma_i\}$ (i.e., with a factor having more than one left extension). Counting the finish and start points defined in [37] (there are $3d$ such points) is equivalent to counting the number of branches in the graph of words.

As in the remark of the previous section, we can consider a coding of the rotation by irrational angle $1 - \alpha$ under the partition $\{[\gamma_1, \gamma_2[, \dots, [\gamma_d, \gamma_1[\}$. For such a coding, the $3d$ distance theorem can be rephrased as follows.

THEOREM 20. *The frequencies of the factors of given length $n \geq n^{(1)}$ of a coding of a rotation by irrational angle under a partition in d intervals take at most $3d$ values, where $n^{(1)}$ denotes the connectedness index.*

ACKNOWLEDGEMENTS. We would like to thank Arnoux, Berstel, Daudé, Dumont, Liardet and Shallit for their bibliographic help. Furthermore, we are greatly indebted to Allouche for his many useful comments and to Ferenczi and Jenkinson who read carefully a previous version of this paper.

REFERENCES

- [1] ALESSANDRI, P. Codages de rotations et basses complexités. Université Aix-Marseille II. Thèse, 1996.
- [2] ALLOUCHE, J.-P. Sur la complexité des suites infinies. *Bull. Belg. Math. Soc. Simon Stevin* 1 (1994), 133–143.

- [3] BERTHÉ, V. Fréquences des facteurs des suites sturmiennes. *Theoret. Comput. Sci.* 165 (1996), 295–309.
- [4] BERTHÉ, V. and L. VUILLON. Tilings and rotations. Prétirage 97–19, Institut de Mathématiques de Luminy.
- [5] BERTHÉ, V., N. CHEKHOVA, and S. FERENCZI. Covering numbers: arithmetics and dynamics for rotations and interval exchanges. Prétirage 97–20, Institut de Mathématiques de Luminy.
- [6] BERSTEL, J. Recent results in Sturmian words. *Developments in Language Theory II (Magdeburg 1995)*. World Scientific (96), 13–24.
- [7] BESSI, G. and J.-L. NICOLAS. Nombres 2-hautement composés. *J. Math. Pures Appl.* 56 (1977), 307–326.
- [8] BOSHERNITZAN, M. A condition for minimal interval exchange maps to be uniquely ergodic. *Duke Math. J.* 52 (1985), 723–752.
- [9] BROWN, T. C. Descriptions of the characteristic sequence of an irrational. *Canad. Math. Bull.* 36 (1993), 15–21.
- [10] CASSAIGNE, J. Special factors of sequences with linear subword complexity. *Developments in Language Theory II (Magdeburg 1995)*. World Scientific (96), 25–34.
- [11] ———. Complexité et facteurs spéciaux. *Bull. Belg. Math. Soc. Simon Stevin* 4 (1997), 67–88.
- [12] CHEKHOVA, N. Covering numbers of rotations. *Theoret. Comput. Sci.*, to appear.
- [13] CHEKHOVA, N., P. HUBERT AND A. MESSAOUDI. Propriétés combinatoires, ergodiques et arithmétiques de la substitution de Tribonacci. Preprint.
- [14] CHEVALLIER, N. Distances dans la suite des multiples d'un point du tore à deux dimensions. *Acta Arith.* 74 (1996), 47–59.
- [15] ———. Géométrie des suites de Kronecker. *Manuscripta Math.* 94 (1997), 231–241.
- [16] ———. Meilleures approximations d'un élément du tore $\mathbf{T}^2$ et géométrie de la suite des multiples de cet élément. *Acta Arith.* 78 (1996), 19–35.
- [17] CHVÁTAL, V., D. A. KLARNER and D. E. KNUTH. Selected combinatorial research problems. *Stanford CS report* 292 (1972).
- [18] CHUNG, F. R. K. and R. L. GRAHAM. On the set of distances determined by the union of arithmetic progressions. *Ars Combin.* 1 (1976), 57–76.
- [19] COVEN, E. M. and G. A. HEDLUND. Sequences with minimal block growth. *Math. Systems Theory* 7 (1973), 138–153.
- [20] DEKKING, F. M. On the Thue-Morse measure. *Acta Univ. Carolin. Math. Phys.* 33 (1992), 35–40.
- [21] DELÉGLISE, M. Recouvrement optimal du cercle par les multiples d'un intervalle. *Acta Arith.* 59 (1991), 21–35.
- [22] DIDIER, G. Caractérisation des N -écritures et applications à l'étude des suites de complexité ultimement $n + c^{ste}$. *Theoret. Comput. Sci.*, to appear.
- [23] ———. Combinatoire des codages de rotations. *Acta Arith.*, to appear.
- [24] FERENCZI, S. Les transformations de Chacon: combinatoire, structure géométrique, lien avec les systèmes de complexité $2n + 1$. *Bull. Soc. Math. France* 123 (1995), 271–292.
- [25] FLOREK, K. Une remarque sur la répartition des nombres $n\xi \pmod{1}$. *Colloq. Math.* 2 (1951), 323–324.

- [26] FRAENKEL, A. S. and R. HOLZMAN. Gap problems for integer part and fractional part sequences. *J. Number Theory* 50 (1995), 66–86.
- [27] FRAENKEL, A. S., M. MUSHKIN, and U. TASSA. Determination of $[n\theta]$ by its sequence of differences. *Canad. Math. Bull.* 21 (1978), 441–446.
- [28] FRIED, E. and V. T. SÓS. A generalization of the three-distance theorem for groups. *Algebra Universalis* 29 (1992), 136–149.
- [29] GEELLEN, A. S. and R. J. SIMPSON. A two dimensional Steinhaus theorem. *Australas. J. Combin.* 8 (1993), 169–197.
- [30] GOTTSCHALK, W. H. and G. A. HEDLUND. Topological dynamics. *American Math. Soc. Colloquium Publications* 36 (1955).
- [31] HALTON, J. H. The distribution of the sequence $\{n\xi\}$ ($n = 0, 1, 2, \dots$). *Proc. Cambridge Philos. Soc.* 61 (1965), 665–670.
- [32] HARDY, G. H. and E. M. WRIGHT. *An Introduction to the Theory of Numbers*. (Fifth ed.) Oxford University Press (1979).
- [33] HARTMAN, S. Über die Abstände von Punkten $n\xi$ auf der Kreisperipherie. *Ann. Soc. Polon. Math.* 25 (1952), 110–114.
- [34] KNUTH, D. E. *The Art of Computer Programming*. Vol. 3, Ex. 6.4.10. Addison Wesley, New York, 1973.
- [35] LANGEVIN, M. Stimulateur cardiaque et suites de Farey. *Period. Math. Hungar.* 23 (1991), 75–86.
- [36] LEFÈVRE, V. An algorithm that computes a lower bound on the distance between a segment and $\mathbf{Z}^2$. *Rapport de recherche LIP* 18 (1997).
- [37] LIANG, F. M. A short proof of the 3d distance theorem. *Discrete Math.* 28 (1979), 325–326.
- [38] MARZEC, C. and J. KAPPRAFF. Properties of maximal spacing on a circle related to phyllotaxis and to the golden mean. *J. Theor. Biol.* 103 (1983), 201–226.
- [39] MORSE, M. and G. A. HEDLUND. Symbolic dynamics. *Amer. J. Math.* 60 (1938), 815–866.
- [40] MORSE, M. and G. A. HEDLUND. Symbolic dynamics II: Sturmian trajectories. *Amer. J. Math.* 62 (1940), 1–42.
- [41] RAUZY, G. Suites à termes dans un alphabet fini. *Sém. de Théorie des Nombres de Bordeaux* (1983), 25-01–25-16.
- [42] VAN RAVENSTEIN, T. On the discrepancy of the sequence formed from multiples of an irrational number. *Bull. Austral. Math. Soc.* 31 (1985), 329–338.
- [43] ——— Optimal spacing of points on a circle. *Fibonacci Quart.* 27 (1989), 18–24.
- [44] ——— The three gap theorem (Steinhaus conjecture). *J. Austral. Math. Soc. Ser. A* 45 (1988), 360–370.
- [45] VAN RAVENSTEIN, T. G. WINLEY and K. TOGNETTI. Characteristics and the three gap theorem. *Fibonacci Quart.* 28 (1990), 204–214.
- [46] ROTE, G. Sequences with subword complexity $2n$. *J. Number Theory* 46 (1994), 196–213.
- [47] SHALLIT, J. O. Automaticity IV: Sequences, sets, and diversity. *J. Théor. Nombres Bordeaux* 8 (1996), 347–367.
- [48] SIEGEL, A. Théorème des trois longueurs et suites sturmiennes: mots d'agencement des longueurs. Preprint.

- [49] SLATER, N. B. The distribution of the integers N for which $\{\theta N\} < \phi$. *Proc. Cambridge Philos. Soc.* 46 (1950), 525–534.
- [50] ——— Distribution problems and physical applications. *Compositio Math.* 16 (1964), 176–183.
- [51] ——— Gaps and steps for the sequence $n\theta \bmod 1$. *Proc. Cambridge Philos. Soc.* 63 (1967), 1115–1123.
- [52] SÓS, V. T. On the theory of diophantine approximations I. *Acta Math. Acad. Sci. Hungar.* 8 (1957), 461–472.
- [53] ——— On the distribution mod 1 of the sequence $n\alpha$. *Ann. Univ. Sci. Budapest. Eötvös Sect. Math.* 1 (1958), 127–134.
- [54] STOLARSKY, K. Beatty sequences, continued fractions, and certain shift operators. *Canad. Math. Bull.* 19 (1976), 473–482.
- [55] SURÁNYI, J. Über die Anordnung der Vielfachen einer reellen Zahl mod 1. *Ann. Univ. Sci. Budapest, Eötvös Sect. Math.* 1 (1958), 107–111.
- [56] ŚWIERCZKOWSKI, S. On successive settings of an arc on the circumference of a circle. *Fund. Math.* 46 (1958), 187–189.
- [57] VUILLON, L. Contribution à l'étude des pavages et des surfaces discrétisées. *Theoret. Comput. Sci.*, to appear.
- [58] WOLFF, A. and J. PITMAN. Lattice points in strips. Tech. Rep., University of Adelaide, Australia, 1990.

(Reçu le 10 janvier 1997; version révisée reçue le 12 février 1998)

Pascal Alessandri

Valérie Berthé

I. M. L.

Case 907

163, av. de Luminy

F-13288 Marseille Cedex 09

France

e-mail: berthe@iml.univ-mrs.fr