

# Conjectures on finite and p-local groups

Autor(en): **Solomon, Eon / Stancu, Radu**

Objektyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **54 (2008)**

Heft 1-2

PDF erstellt am: **22.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-109929>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## 60

### CONJECTURES ON FINITE AND $p$ -LOCAL GROUPS

by Ron SOLOMON and Radu STANCU

Algebraic topology and group theory are strongly interacting mathematical fields and in the past decade new connections between the  $p$ -completion of the classifying space of a group and its  $p$ -local structure were discovered. A seminal theorem relating the  $p$ -local structure with the mod- $p$  cohomology of a finite group is the following [14].

**THEOREM 60.1** (Mislin 1990). *A subgroup  $H$  of a finite group  $G$  controls the  $p$ -fusion in  $G$  if and only if the canonical inclusion of  $H$  in  $G$  induces an isomorphism in all degrees between the mod- $p$  cohomology rings of  $G$  and  $H$ .*

In other words  $G$  and  $H$  have the same  $p$ -local structure if and only if they have the same mod- $p$  cohomology.

The conjectures and questions below continue in this vein. In the context of group representations and local group theory, a fundamental conjecture is Alperin's Weight Conjecture [1]. Here is a formulation with a topological flavor.

**DEFINITION 60.2.** Let  $G$  be a finite group. A  $p$ -chain  $C$  of  $G$  is a strictly increasing chain

$$C: P_0 < P_1 < \cdots < P_n$$

of  $p$ -subgroups of  $G$ . We let  $C_i$  denote the initial subchain terminating at  $P_i$ . The chain  $C$  is *radical* if  $P_0 = O_p(G)$  and, for each  $i$ ,  $P_i = O_p(N_G(C_i))$ .

We denote by  $\mathcal{R}(G)$  the set of all radical  $p$ -chains of  $G$ , and by  $\mathcal{R}(G)/G$  its orbit space.

DEFINITION 60.3. Fix a prime  $p$ . Let  $\phi$  be an ordinary irreducible character of the finite group  $H$ . The *defect*  $d(\phi)$  of  $\phi$  is the largest non-negative integer  $d$  such that  $p^d$  divides  $\frac{|H|}{\phi(1)}$ .

DEFINITION 60.4. Let  $G$  be a finite group and  $B$  a  $p$ -block of  $G$ . For any  $p$ -chain  $C$  of  $G$  we denote by  $k(C, B, d)$  the number of characters  $\phi \in \text{Irr}(N_G(C))$  having defect  $d(\phi) = d$  and belonging to a  $p$ -block  $B(\phi)$  of  $N_G(C)$  such that the induced  $p$ -block  $B(\phi)^G$  is  $B$ .

CONJECTURE 60.5. Let  $p$  be a prime,  $G$  a finite group with  $O_p(G) = 1$ , and  $B$  a  $p$ -block of  $G$  which is not of defect 0. Then

$$\sum_{C \in \mathcal{R}(G)/G} (-1)^{|C|} k(C, B, d) = 0.$$

This formulation of the conjecture is due to Dade [6], Conjecture 6.3. Using work of Knorr and Robinson [12], Dade proved that it implies the original Alperin Weight Conjecture. Extensions of the Alperin–Dade Conjecture to projective characters are given in [7].

Next, here is a statement about finite groups. In some sense it is not a conjecture because we are fairly certain it can be proved easily as a corollary of the Classification of Finite Simple Groups. The conjecture is that there is an elementary proof, perhaps following from further advances in the  $p$ -modular representation theory of finite groups. (The case  $p = 2$  is a theorem of Goldschmidt [9].)

DEFINITION 60.6. Let  $p$  be a prime and  $P$  a  $p$ -group. Then  $\Omega_1(P)$  is the subgroup of  $P$  generated by all elements of order  $p$ .

CONJECTURE 60.7. Let  $p$  be a prime and let  $G$  be a finite group having an abelian Sylow  $p$ -subgroup  $A$ . Suppose that  $B$  is a strongly closed subgroup of  $A$  with respect to  $G$ , i.e. if  $x \in B$  and  $g \in G$  with  $x^g \in A$ , then  $x^g \in B$ . Then there exists a normal subgroup  $N$  of  $G$  having a Sylow  $p$ -subgroup  $B^*$  such that  $\Omega_1(B) = \Omega_1(B^*)$ . In particular, if  $A = \Omega_1(A)$  and  $B$  is strongly closed in  $A$  with respect to  $G$ , then  $B \in \text{Syl}_p(N)$  for some normal subgroup  $N$  of  $G$ .

The following conjectures are related to the work of Broto, Levi, and Oliver [4] on the theory of so-called  $p$ -local groups. The underlying struc-

ture of a  $p$ -local group is a fusion system on a finite  $p$ -group  $P$ . Fusion systems were introduced by Puig in 1990 [15]. We define them here.

First let us start with a more general definition:

DEFINITION 60.8. A *category  $\mathcal{F}$  on a finite  $p$ -group  $P$*  is a category whose objects are the subgroups of  $P$  and whose set of morphisms between the subgroups  $Q$  and  $R$  of  $P$ , is the set  $\text{Hom}_{\mathcal{F}}(Q, R)$  of injective group homomorphisms from  $Q$  to  $R$ , with the following properties:

- (a) if  $Q \leq R$  then the inclusion of  $Q$  in  $R$  is a morphism in  $\text{Hom}_{\mathcal{F}}(Q, R)$ .
- (b) for any  $\phi \in \text{Hom}_{\mathcal{F}}(Q, R)$  the induced isomorphism  $Q \simeq \phi(Q)$  and its inverse are morphisms in  $\mathcal{F}$ .
- (c) composition of morphisms in  $\mathcal{F}$  is the usual composition of group homomorphisms.

And now the definition of a fusion system:

DEFINITION 60.9. A *fusion system  $\mathcal{F}$  on a finite  $p$ -group  $P$*  is a category on  $P$  satisfying the following properties:

- (1)  $\text{Hom}_P(Q, R) \subset \text{Hom}_{\mathcal{F}}(Q, R)$  for all  $Q, R \leq P$ .
- (2)  $\text{Aut}_P(P)$  is a Sylow  $p$ -subgroup of  $\text{Aut}_{\mathcal{F}}(P)$ .
- (3) Every  $\phi: Q \rightarrow P$  such that  $|N_P(\phi(Q))|$  is maximal in the  $\mathcal{F}$ -isomorphism class of  $Q$ , extends to  $\bar{\phi}: N_{\phi} \rightarrow P$  where

$$N_{\phi} = \{x \in N_P(Q) \mid \exists y \in N_P(\phi(Q)), \phi(xu) = {}^y\phi(u) \forall u \in Q\}.$$

Axiom (2) says that  $P$  is a ‘Sylow  $p$ -subgroup’ of  $\mathcal{F}$  and the extension in Axiom (3) is equivalent to saying that any  $p$ -subgroup can be embedded by conjugation in a Sylow  $p$ -subgroup. In particular, if  $G$  is a finite group,  $p$  is a prime divisor of  $|G|$ , and  $P \in \text{Syl}_p(G)$ , then the morphisms given by conjugation by elements of  $G$  between the subgroups of  $P$  determine a fusion system  $\mathcal{F}_G(P)$  on  $P$ . We say that a fusion system is *exotic* if it does not arise in this way.

The literature on fusion systems contains slightly different definitions of a normal subsystem, and it is as yet unclear which is the correct one. We follow Markus Linckelmann’s approach to normal subsystems [13]. First let’s introduce the notion of *strongly  $\mathcal{F}$ -closed* subgroups.

DEFINITION 60.10. Let  $\mathcal{F}$  be a fusion system on a finite  $p$ -group  $P$  and  $Q$  a subgroup of  $P$ . We say that  $Q$  is *strongly  $\mathcal{F}$ -closed* if for any subgroup  $R$  of  $Q$  and any morphism  $\phi \in \text{Hom}_{\mathcal{F}}(R, P)$  we have  $\phi(R) \leq Q$ .

And now the notion of a normal fusion subsystem.

DEFINITION 60.11. Let  $\mathcal{F}$  be a fusion system on a finite  $p$ -group  $P$  and  $\mathcal{F}'$  a fusion subsystem of  $\mathcal{F}$  on a subgroup  $P'$  of  $P$ . We say that  $\mathcal{F}'$  is *normal* in  $\mathcal{F}$  if  $P'$  is strongly  $\mathcal{F}$ -closed and if for every isomorphism  $\phi: Q \rightarrow Q'$  in  $\mathcal{F}$  and any two subgroups  $R, R'$  of  $Q \cap P'$  we have

$$\phi \circ \text{Hom}_{\mathcal{F}'}(R, R') \circ \phi^{-1} \subseteq \text{Hom}_{\mathcal{F}'}(\phi(R), \phi(R')).$$

Here is a conjecture on normal fusion subsystems:

CONJECTURE 60.12. *Let  $\mathcal{F}$  be a fusion system on  $P$  and  $P'$  a strongly  $\mathcal{F}$ -closed subgroup of  $P$ . Then there exists a normal fusion subsystem  $\mathcal{F}'$  of  $\mathcal{F}$  on  $P'$ .*

A *simple fusion system* is a fusion system that has no non-trivial normal subsystems. In particular, if  $G$  is a finite simple group which does not have a proper strongly  $p$ -embedded subgroup, then  $\mathcal{F}_G(P)$  is a simple fusion system.

When  $p$  is odd, it seems to be fairly easy to construct examples of exotic simple  $p$ -local groups. On the other hand when  $p = 2$ , the only known exotic examples live in a single infinite family,  $\mathcal{F}_{Sol}(q)$ , which may be regarded as the analogue of finite Chevalley groups with respect to the exotic 2-compact group of Dwyer and Wilkerson.

CONJECTURE 60.13. *The family  $\mathcal{F}_{Sol}(q)$  contains all of the exotic simple 2-local groups.*

This conjecture is hard to believe. On the other hand, it has thus far been impossible to dream up other examples. It is conceivable that it could be proved by a lengthy and elaborate analysis in the vein of the traditional 2-local analysis of finite simple groups used in the proof of the Classification Theorem. It would be much more interesting if homotopy-theoretic tools could be brought to bear to prove this result. That would really give hope for new and exciting applications of topology to finite group theory.

Simple Lie groups are of course linked to real reflection groups. Similarly,  $p$ -compact groups are linked to  $p$ -adic reflection groups. This suggests the following question.

QUESTION 60.14. *Is there a family of interesting topological objects connected to quaternionic reflection groups? (See [5].)*

As the Dwyer–Wilkerson 2-compact group [8] is, in some sense, a 45-dimensional object, it is a plausible conjecture that it might have some connection to the 45-dimensional algebra  $SH(3, \mathfrak{D})$  of  $3 \times 3$  skew hermitian matrices over the octonions (with bracket multiplication).

QUESTION 60.15. *What is the automorphism group of this algebra and of related rings, e.g. defined over rings of integral octonions?*

Finally, we note that another source of examples for fusion systems are the fusion systems coming from  $p$ -blocks of group algebras, given by the conjugations between the Brauer pairs in a maximal Brauer pair of the  $p$ -block. Such examples are called Brauer categories [2]. Here is a natural question one can ask.

QUESTION 60.16. *Are there Brauer categories which are exotic fusion systems?*

It is pretty hard to check that a given fusion system is not a Brauer category. Up to now, the only known way to do it is by reduction to Brauer categories of quasisimple groups and then by using the classification of finite simple groups [10], [11]. There are examples of embedded fusion systems where the minimal and the maximal ones come from finite groups and the intermediate ones are exotic. It is not yet known whether these exotic fusion systems are Brauer categories.

#### REFERENCES

- [1] ALPERIN, J.L. Weights for finite groups. In: *The Arcata Conference on Representations of Finite Groups*, 369–379. Proc. Sympos. Pure Math. 47. Amer. Math. Soc., Providence, R.I., 1987.
- [2] ALPERIN J.L. and M. BROUÉ. Local methods in block theory. *Ann. of Math.* 110 (1979), 143–157.

- [3] ASCHBACHER, M. and A. CHERMAK. A group-theoretic approach to a family of 2-local finite groups constructed by Levi and Oliver. (Preprint, 2005.)
- [4] BROTO, C., R. LEVI and B. OLIVER. The homotopy theory of fusion systems. *J. Amer. Math. Soc.* 16 (2003), 779–856.
- [5] COHEN, A.M. Finite quaternionic reflection groups. *J. Algebra* 64 (1980), 293–324.
- [6] DADE, E.C. Counting characters in blocks, I. *Invent. Math.* 109 (1992), 187–210.
- [7] ——— Counting characters in blocks, II. *J. Reine Angew. Math.* 448 (1994), 97–190.
- [8] DWYER, W.G. and C.W. WILKERSON. A new finite loop space at the prime two. *J. Amer. Math. Soc.* 6 (1993), 37–64.
- [9] GOLDSCHMIDT, D.M. 2-fusion in finite groups. *Ann. of Math.* 99 (1974), 70–117.
- [10] KESSAR, R. The Solomon system  $\mathcal{F}_{\text{Sol}}(3)$  does not occur as fusion system of a 2-block. *J. Algebra* 296 (2006), 409–425.
- [11] KESSAR, R. and R. STANCU. A reduction theorem for fusion systems of blocks. *J. Algebra* 319 (2008), 806–823.
- [12] KNORR, R. and G.R. ROBINSON. Some remarks on a conjecture of Alperin. *J. London Math. Soc.* (2) 39 (1989), 48–60.
- [13] LINCKELMANN, M. Simple fusion systems and the Solomon 2-local groups. *J. Algebra* 296 (2006), 385–401.
- [14] MISLIN, G. On group homomorphisms inducing mod- $p$  cohomology isomorphisms. *Comment. Math. Helv.* 65 (1990), 454–461.
- [15] PUIG, L. Frobenius categories. *J. Algebra* 303 (2006), 309–357. (Preprint since 1990.)
- [16] SOLOMON, R. Finite groups with Sylow 2-subgroups of type .3. *J. Algebra* 28 (1974), 182–198.

R. Solomon

The Ohio State University  
 231 West 18th Avenue  
 Columbus, OH 43210  
 USA  
*e-mail*: solomon@math.ohio-state.edu

R. Stancu

The Ohio State University  
 231 West 18th Avenue  
 Columbus, OH 43210  
 USA  
*e-mail*: stancu@math.ohio-state.edu