

Alternating quotients of free groups

Autor(en): **Wilton, Henry**

Objektyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **58 (2012)**

PDF erstellt am: **25.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-323246>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

ALTERNATING QUOTIENTS OF FREE GROUPS

by Henry WILTON*)

ABSTRACT. We strengthen Marshall Hall's theorem to show that free groups are locally extended residually alternating. Let F be any free group of rank at least two, let H be a finitely generated subgroup of infinite index in F and let $\{\gamma_1, \dots, \gamma_n\} \subseteq F \setminus H$ be a finite subset. Then there is a surjection f from F to a finite alternating group such that $f(\gamma_i) \notin f(H)$ for any i . The techniques of this paper can also provide symmetric quotients.

DEFINITION 1. A group Γ is *residually* \mathcal{P} (for some class of groups \mathcal{P}) if, for any $\gamma \in \Gamma \setminus 1$, there is a surjection $f: \Gamma \rightarrow P \in \mathcal{P}$ with $f(\gamma) \neq 1$.

Many groups are known to be residually finite, and it is natural to ask whether one can restrict attention to smaller classes of finite groups. Katz and Magnus proved that free groups are residually alternating, hence residually simple [6] (see also [8], [12] and [11] *et seq.*); in the topological context, Long and Reid showed that many hyperbolic 3-manifold groups are residually $PSL_2(\mathbf{F}_p)$ [7].

One of the most obvious generalisations of Definition 1 replaces the trivial subgroup with an arbitrary finitely generated subgroup H . (In many cases, it is too much to expect infinitely generated subgroups to satisfy this property — see Remark 3.) This consideration leads to the following definition.

DEFINITION 2. A group Γ is said to be *locally extended residually finite* (*LERF*, also often called *subgroup separable*) if, for any finitely generated subgroup $H \subseteq \Gamma$ and any $\gamma \in \Gamma \setminus H$, there is a surjection f from Γ onto a finite group such that $f(\gamma) \notin f(H)$.

*) Partially supported by NSF grant number DMS-0906276 and by an EPSRC Career Acceleration Fellowship.

REMARK 3. Without the requirement that the subgroup H be finitely generated, such a group Γ is called *extended residually finite* or *ERF*. It is well known that the $(2, 3)$ -Baumslag–Solitar group

$$BS(2, 3) = \langle a, b \mid b^{-1}a^{-2}ba^3 \rangle$$

is not residually finite. By considering the kernel of the natural map $F_2 \cong \langle a, b \rangle \rightarrow BS(2, 3)$, it follows easily that non-abelian free groups are not ERF.

Marshall Hall Jr. proved that free groups are locally extended residually finite [4]. His theorem has been reinterpreted topologically and generalised to much larger classes of groups [9, 10, 13, 3]. However, to the best of the author's knowledge, no results have been proved that restrict the finite quotients that arise to a more specific class. The aim of this note is to begin to fill this gap by proving that free groups are what one might call locally extended residually alternating.

THEOREM A. *Let F be a free group of rank greater than one. Let H be a finitely generated subgroup of infinite index in F and let $\{\gamma_1, \dots, \gamma_n\}$ be a finite subset of $F \setminus H$. There is a surjection f from F onto some finite alternating group A_k such that $f(\gamma_i) \notin f(H)$ for all i .*

A small modification of the argument gives symmetric, rather than alternating, quotients.

THEOREM B. *Let F be a free group of rank greater than one. Let H be a finitely generated subgroup of infinite index in F and let $\{\gamma_1, \dots, \gamma_n\}$ be a finite subset of $F \setminus H$. There is a surjection f from F onto some finite symmetric group S_k such that $f(\gamma_i) \notin f(H)$ for all i .*

REMARK 4. In the case when \mathcal{P} is the class of all finite groups, it is equivalent to state the theorem for a single element $\gamma \notin H$ instead of for a finite subset $\{\gamma_1, \dots, \gamma_n\} \subseteq \Gamma \setminus H$: to deduce the latter from the former, simply take the product of the quotients across all γ_i . This works because a product of finite groups is finite. As a product of alternating groups is not alternating and a product of symmetric groups is not symmetric, we give the stronger statements.

REMARK 5. The hypothesis that H is of infinite index in F is necessary. For instance, suppose that H is a normal subgroup of finite index in F . For any surjection $f: F \rightarrow A_n$ with $n \geq 5$, $f(H)$ is a normal subgroup of A_n and is therefore either the whole of A_n or trivial. In the latter case, it follows that F/H maps onto A_n . But F has many finite quotients that do not map onto A_n .

Stallings interpreted Hall's original proof that free groups are LERF using the topology of graphs, reducing it to the topological fact that any immersion of finite graphs can be completed to a covering map [10]. Wiegold's proof that free groups are residually even alternating [12] uses a classical theorem of Jordan, which asserts that if the minimal degree of a primitive permutation group is small enough, then that group must be symmetric or alternating [5]. The proofs of Theorems A and B combine Jordan's theorem with the covering theory of graphs. The key technical result is Lemma 18. In the proof, we show how to complete an immersion of a finite graph into the rose to a covering map in such a way that the resulting permutation action of F on the vertices is primitive and satisfies Jordan's condition. This proves that free groups are, so to speak, locally extended residually symmetric-or-alternating. A small modification of this construction then forces the action to be alternating; a slightly different modification forces the action to be symmetric.

In order to apply Jordan's theorem, we must ensure that the action of F on the vertices of the cover we construct is primitive. In this paper we do so in the simplest possible way, by requiring the number of vertices of the cover to be prime. Thus, we actually prove that free groups are locally extended residually alternating-of-prime-degree. Alternatively, one could ensure that the action is primitive by forcing it to be 2-transitive. This is possible, using a more complicated construction, of which we do not give the details here. Via this more complicated construction, one can obtain different restrictions on the degrees of the alternating quotients.

After free groups, the fundamental groups of surfaces are the next examples to consider.

CONJECTURE 6. *Let Σ be a closed, orientable, hyperbolic surface and let H be a finitely generated subgroup of infinite index in $\pi_1\Sigma$. For any $\gamma \in \pi_1\Sigma \setminus H$ there is a surjection f from Γ onto a finite alternating group with $f(\gamma) \notin f(H)$.*

In Section 1, we recall the very well known facts that subgroups of free groups can be viewed in terms of either permutation representations or coverings of graphs, and observe that it is easy to pass from one point of view to the other. In Section 2 we revisit Stallings's proof of Marshall Hall's theorem. Finally, in Section 3 we bring in Jordan's theorem and prove Theorems A and B.

1. PERMUTATIONS AND COVERINGS OF GRAPHS

Let F be a free group of rank $r > 1$, with generators $\{\alpha_1, \dots, \alpha_r\}$. It is convenient to take r to be finite, although our results apply just as well to the infinite-rank case. Let H be a subgroup of finite index. The action of F by left multiplication on F/H can be thought of as a homomorphism from F to the symmetric group $\text{Sym}(F/H)$. Of course, F acts transitively, and the subgroup H can be recovered as the stabiliser of the trivial coset. We summarise this paragraph as follows.

REMARK 7. Subgroups of F of index d correspond canonically to transitive actions of F on pointed sets of cardinality d .

We now switch to a topological point of view. Let X be a rose with r petals, that is, a graph with precisely one vertex x_0 and r edges. (All our results can be generalised to the situation in which X is an arbitrary finite graph. However, the case in which X is a rose is sufficient for our purposes.) We fix once and for all an isomorphism $F \cong \pi_1(X, x_0)$ by orienting each edge of X and labelling it with a generator of F .

By standard covering space theory, a subgroup H of F corresponds to a connected, pointed covering space $(Y, y_0) \rightarrow (X, x_0)$ with the covering map inducing an isomorphism $\pi_1(Y, y_0) \cong H$.

The orientation and labelling of the edges of X pulls back to an orientation and labelling of the edges of Y . Conversely, any orientation and labelling of the edges of Y determines a combinatorial map $Y \rightarrow X$ that sends vertices to vertices and edges to edges. We will only consider such maps, and we will usually think of them in terms of the corresponding labelled and oriented graph Y .

Covering maps to X are easily characterised in terms of the labelling and orientation on Y .

DEFINITION 8. A labelling and orientation on a graph Y are said to satisfy the *covering condition* if, for each vertex y of Y and each label α_i , there is exactly one incoming and one outgoing edge labelled α_i at y .

The proof of the following lemma is a simple exercise.

LEMMA 9. *A combinatorial map of non-empty, connected graphs $Y \rightarrow X$ is a covering map if and only if it is bijective on the links of vertices. Equivalently, $Y \rightarrow X$ is a covering map if and only if the corresponding labelling and orientation on the graph Y satisfy the covering condition.*

When a covering map is restricted to a subgraph, we obtain an immersion.

DEFINITION 10. A combinatorial map $Y \rightarrow X$ is an *immersion* if it is injective on links of vertices.

Again, this can be characterised in terms of the labelling and orientation on Y .

DEFINITION 11. A labelling and orientation on a graph Y are said to satisfy the *immersion condition* if, for each vertex y of Y and each label α_i , there is at most one incoming and one outgoing edge labelled α_i at y .

Just as before, the proof of the following lemma is an easy exercise.

LEMMA 12. *A combinatorial map of non-empty, connected graphs $Y \rightarrow X$ is an immersion if and only if the corresponding labelling and orientation on the graph Y satisfy the immersion condition.*

The immersion condition and the covering condition are illustrated in Figure 1.




FIGURE 1

The graph on the left satisfies the covering condition.
The graph on the right does not satisfy the immersion condition.

It follows from standard covering space theory that F acts on the vertex set of Y by path lifting. To be precise, any element $\gamma \in F$ can be thought of as a loop in X , represented by a continuous map $\gamma: [0, 1] \rightarrow X$ with $\gamma(0) = \gamma(1) = x_0$. Covering space theory asserts that, for each vertex $y \in Y$, there is a unique lift $\tilde{\gamma}: [0, 1] \rightarrow Y$ with $\tilde{\gamma}(0) = y$. The action of γ on the vertices of Y is defined by

$$\gamma \cdot y = \tilde{\gamma}(1) .$$

Furthermore, the map that sends $\gamma \mapsto \gamma \cdot y_0$ induces an F -equivariant bijection between F/H and the vertices of Y . We can summarise this discussion in the following remark.

REMARK 13. Subgroups of F correspond to connected, based, labelled, oriented graphs that satisfy the covering condition. The index of the subgroup is equal to the number of vertices of the graph.

It is apparent from Remarks 7 and 13 that connected, based, labelled, oriented graphs satisfying the covering condition are in bijection with transitive permutation actions of F on based sets. Indeed, given such a graph Y , as observed above the group F acts by path lifting on the vertices. The action of the generator α_i can be seen by restricting attention to the edges of Y labelled by α_i : by the covering condition, these edges form a union of topological circles; each circle corresponds to a cycle under the action of α_i ; taken together, these circles give the cycle type of α_i .

Conversely, given an action of F by permutations on a set V , we can build a covering graph Y . Take V to be the vertex set of Y . There is an oriented edge labelled α_i from u to v if $\alpha_i \cdot u = v$. By construction, this graph Y satisfies the covering condition.

2. MARSHALL HALL'S THEOREM

In this section, we recall Stallings's proof of Marshall Hall's Theorem [10]. The key observation is that one can complete any immersion to a covering map without increasing the number of vertices.

LEMMA 14 (Stallings [10]). *Let Z be a finite graph, let X be a rose and let $Z \rightarrow X$ be an immersion. The immersion factors as $Z \hookrightarrow Y \rightarrow X$ where Z is a subgraph of Y , every vertex of Y is a vertex of Z , and $Y \rightarrow X$ is a covering map.*

Proof. As above, we can think of Z as an oriented, labelled graph that satisfies the immersion condition. For each i , let $A^+(i)$ be the set of vertices of Z that adjoin an outgoing edge labelled α_i , and let $A^-(i)$ be the set of vertices of Z that adjoin an incoming edge labelled α_i . Counting the edges labelled α_i , we see that

$$\#A^+(i) = \#A^-(i)$$

whence

$$\#(A^+(i))^c = \#(A^-(i))^c.$$

Choose any bijection between $(A^+(i))^c$ and $(A^-(i))^c$. We can use this additional bijection to add new oriented edges labelled α_i to Z . Let Y be the result of carrying this out for each i . By construction, Y satisfies the covering condition. \square

The proof is illustrated in Figure 2.




FIGURE 2

The union of the black edges, Z , satisfies the immersion condition.
 After adding the grey edges, the resulting graph Y satisfies the covering condition.

THEOREM 15 (Marshall Hall Jr. [4]). *Let F be a finitely generated free group, let $H \subseteq F$ be a finitely generated subgroup, and let $\gamma_1, \dots, \gamma_n \in F \setminus H$. There is a homomorphism f from F to a finite group such that $f(\gamma_i) \notin f(H)$ for all i .*

Proof. We identify F with $\pi_1(X, x_0)$ for X a suitable rose. Let $(X', x'_0) \rightarrow (X, x_0)$ be the covering map corresponding to H . Each γ_i lifts at x'_0 to a path $\gamma'_i: [0, 1] \rightarrow X'$; by assumption, $\gamma'_i(1) \neq x'_0$. We now consider the topology of the graph X' . Let $T \subseteq X'$ be a maximal tree. Because H is finitely generated, it follows from the Seifert–van Kampen Theorem that $X' \setminus T$ is a finite union of edges $e'_1 \cup \dots \cup e'_k$.

For each j , let μ'_j be an edge path from x'_0 to the initial vertex of ϵ'_j and let ν'_j be an edge path from x'_0 to the terminal vertex of ϵ'_j . Let

$$Z = \bigcup_{i=1}^n \gamma'_i \cup \bigcup_{j=1}^k (\mu'_j \cup \epsilon'_j \cup \nu'_j) .$$

That is, Z is a finite, connected subgraph of X' that contains the paths γ'_i and that also carries the fundamental group of X' . The labelling of X' restricted to Z satisfies the immersion condition, so we may apply Lemma 14 to construct a finite-sheeted covering space Y .

Consider the action of F on the vertices of Y . Every element of H is represented by a loop in Z . Therefore, H is contained in the stabiliser of x'_0 . On the other hand, $\gamma'_i(1) \neq x'_0$ for each i , and so γ_i does not stabilise x'_0 . \square

3. ALTERNATING QUOTIENTS

In this section, we show how to modify Stallings's construction to force the action of F on the vertices of Y to be alternating or symmetric.

Recall some basic terminology from the theory of symmetric group actions. Consider a transitive action by a group Γ on a finite set of order n , in other words a homomorphism $\Gamma \rightarrow S_n$ with transitive image; the integer n is called the *degree* of the action. The smallest degree of a non-trivial cyclic subgroup of the image of Γ is called the *minimal degree* of the action. The action is *primitive* if it does not preserve any proper partition of the finite set.

REMARK 16. Because the action is transitive, the cardinality of any partition preserved by Γ divides n . Therefore, if n is prime then the action is primitive.

We will use a classical theorem of Jordan, which gives a criterion for an action to be symmetric or alternating (in other words, for the homomorphism $\Gamma \rightarrow S_n$ to have image of index at most two) [5, 1]. See Theorem 3.3D of [2] for a modern treatment.

THEOREM 17 (Jordan). *There is a function $J: \mathbf{N} \rightarrow \mathbf{N}$ with the following properties :*

1. $J(n) \rightarrow \infty$ as $n \rightarrow \infty$;
2. *if $\Gamma \rightarrow S_n$ is primitive with minimal degree at most $J(n)$ then the image of Γ is the symmetric group S_n or the alternating group A_n .*

We can now provide the alternating analogue of Lemma 14.

LEMMA 18. *Assume that $r \geq 2$. Let Z be a finite graph and let X be a rose with r petals. Any immersion $Z \rightarrow X$ that is not a covering map factors as $Z \hookrightarrow Y \rightarrow X$ where $Y \rightarrow X$ is a finite-sheeted covering map and the action of $F \cong \pi_1(X)$ on the vertex set of Y is alternating.*

Proof. As before, we think of Z as a labelled, oriented graph. Because the immersion $Z \rightarrow X$ is not a covering map, the labelling and orientation on the graph Z do not satisfy the covering condition, so for some i there are (not necessarily distinct) vertices a, b of Z such that a does not adjoin an outgoing edge labelled α_i and b does not adjoin an incoming edge labelled α_i . Without loss of generality, we may take $i = 1$. By adding edges to Z as in the proof of Lemma 14, we may assume that a and b are the only vertices that adjoin fewer than $2r$ half-edges. Let d be the number of vertices of Z .

Before we proceed with the details of the proof, we will give an outline. The labelled, oriented graph Y is constructed from Z by attaching a large labelled, oriented graph W_n on which α_2 acts trivially. We need to ensure that Y has the following properties:

1. the labelling and orientation on Y satisfy the covering condition;
2. the number of vertices of Y is prime (which ensures that the action of F on the vertices of Y is primitive);
3. the generator α_2 fixes a large number of the vertices of Y (this ensures that the action of F on the vertices of Y has small minimal degree).

At this point it will follow from Theorem 17 that the action of F on the vertices of Y is either alternating or symmetric. In order to force the action to be alternating, we will also attach a carefully chosen labelled, oriented graph V_s . A schematic diagram of the construction of Y is given in Figure 5.

We will now give the details of the construction of W_n . Take a graph homeomorphic to an interval with $n + 1$ vertices, denoted w_0, \dots, w_n . Label each edge α_1 and orient the edges consistently, so that the immersion condition is satisfied. Attach an oriented edge labelled α_i from w_j to itself for each $i \neq 1$ and for each j . The result is a labelled, oriented graph W_n , with n vertices, that satisfies the immersion condition. Only two vertices of W_n adjoin fewer than $2r$ half edges: one does not adjoin an incoming edge labelled α_1 and the other does not adjoin an outgoing edge labelled α_1 . An example is shown in Figure 3.

Next, consider any sequence $(s_i) \in \{\pm 1\}$ (for $1 \leq i \leq r$). We will define a labelled, oriented graph V_s , with four vertices, denoted v_1, v_2, v_3, v_4 . By changing the sequence (s_i) , we will be able to change the signs of the




FIGURE 3
The graph W_4

permutations defined by the generators of F . The construction is as follows. There is an oriented edge labelled α_1 from v_1 to v_2 ; for each $i > 2$, there are oriented loops of length one attached to v_1 and v_2 . For each $i \neq 2$, if $s_i = +1$ then there are oriented loops of length one labelled α_i attached to v_3 and v_4 ; if $s_i = -1$ then v_3 and v_4 are the two vertices of an oriented loop of length two labelled α_i . If $s_2 = +1$ then v_1 and v_3 are the two vertices of an oriented loop of length two labelled α_2 , and likewise v_2 and v_4 are the two vertices of such a loop; if $s_2 = -1$ then v_1, v_2, v_3, v_4 are the vertices of an oriented loop of length four labelled α_2 . Two examples are shown in Figure 4.




FIGURE 4
The graph $V_{+1,-1}$ is shown on the left, and $V_{-1,+1}$ on the right.

We are nearly ready to construct Y from the pieces Z , W_n and V_s . However, we first construct a similar labelled, oriented graph Y' , in which we will not try to control the parity of the action of F . Let p be the smallest prime such that $J(p) \geq d+4$ and let $n = p-d-4$. Let $t_i = +1$ for $1 \leq i \leq r$. Construct Y' from $V_i \sqcup Y \sqcup W_n$ by attaching four oriented edges labelled α_1 in any way that makes the result connected and that satisfies the covering condition. So F acts on the vertex set of Y' , and we can read off the cycle types by looking at the edges. Because Y' is connected, the action is transitive, and because the number of vertices p is prime, it follows that the action is primitive. By construction, $\alpha'_2.w_j = w_j$ for any j , and so we see that the minimal degree of the action is at most $d+4$. It follows from Jordan's theorem that the action is alternating or symmetric.




FIGURE 5

The construction of Y from Z is illustrated. The original edges of Z are marked in black, and the added edges in grey. The labels have been suppressed for clarity.

Finally, we will modify Y' to force the action to be alternating. For each i , consider the sign $s_i \in \{\pm 1\}$ of the action of α_i on the vertex set of Y' . Now construct Y from Y' by replacing V_i by V_{s_i} . As before, we see that the action of F on the vertex set of Y is symmetric or alternating, but the modification ensures that each generator α_i acts as an even permutation. Therefore, the action of F on the vertex set of Y is alternating. \square

The construction of Y is illustrated in Figure 5. Note that, in order to ensure that the resulting action is symmetric rather than alternating, one can simply change the sequence s in one place. We therefore also have a symmetric version of the previous lemma.

LEMMA 19. *Assume that $r \geq 2$. Let Z be a finite graph. Any immersion $Z \rightarrow X$ that is not a covering map factors as $Z \hookrightarrow Y \rightarrow X$ where $Y \rightarrow X$ is a finite-sheeted covering map and the action of $F \cong \pi_1(X)$ on the vertex set of Y is symmetric.*

Theorem A follows from Lemma 18 in exactly the same way that Theorem 15 follows from Lemma 14. Likewise, Theorem B follows from Lemma 19.

REFERENCES

- [1] DIEUDONNÉ, J. *Œuvres de Camille Jordan. Tomes I, II*. Publiées sous la direction de M. Gaston Julia. Gauthier-Villars & Cie, Paris, 1961.
- [2] DIXON, J.D. and B. MORTIMER. *Permutation Groups*. Graduate Texts in Mathematics 163. Springer-Verlag, New York, 1996.
- [3] HAGLUND, F. Finite index subgroups of graph products. *Geom. Dedicata* 135 (2008), 167–209.
- [4] HALL, M., JR. Subgroups of finite index in free groups. *Canad. J. Math.* 1 (1949), 187–190.
- [5] JORDAN, C. Théorèmes sur les groupes primitifs. *J. Math. Pures Appl. (2)* 16 (1871), 383–408.
- [6] KATZ, R.A. and W. MAGNUS. Residual properties of free groups. *Comm. Pure Appl. Math.* 22 (1968), 1–13.
- [7] LONG, D.D. and A. W. REID. Simple quotients of hyperbolic 3-manifold groups. *Proc. Amer. Math. Soc.* 126 (1998), 877–880.
- [8] PRIDE, S.J. Residual properties of free groups. *Pacific J. Math.* 43 (1972), 725–733.
- [9] SCOTT, P. Subgroups of surface groups are almost geometric. *J. London Math. Soc. (2)* 17 (1978), 555–565.
- [10] STALLINGS, J.R. Topology of finite graphs. *Invent. Math.* 71 (1983), 551–565.
- [11] WEIGEL, T.S. Residual properties of free groups. *J. Algebra* 160 (1993), 16–41.
- [12] WIEGOLD, J. Free groups are residually alternating of even degree. *Archiv der Math.* 28 (1977), 337–339.
- [13] WILTON, H. Hall’s theorem for limit groups. *Geom. Funct. Anal.* 18 (2008), 271–303.

(Reçu le 6 mai 2010)

Henry Wilton

Department of Mathematics
University College London
Gower Street
London WC1E 6BT
United Kingdom
e-mail: hwilton@math.ucl.ac.uk