Zeitschrift: Veröffentlichungen des Geobotanischen Institutes Rübel in Zürich

Herausgeber: Geobotanisches Institut Rübel (Zürich)

Band: 8 (1932)

Artikel: The beech forests of Czechoslovakia

Autor: Domin, Karel

Kapitel: XIV: Spore plants

DOI: https://doi.org/10.5169/seals-307034

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 02.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

a Quercetum lanuginosae (with scattered Fagus, Quercus cerris, Quercus sessilis). Analogous examples Mikyška gives from the eruptive rocks (mostly andesites) in the Štiavnické Středohoří Mts. where, however, Quercus robur is the dominant tree.

From a sociological standpoint, we may distinguish in this category besides the sociations already described (as for instance *Oxalis-Galeobdolon*, *Luzula nemorosa* sociations) especially the following:

- 1. Majanthemum sociation.
- 2. Festuca ovina Luzula nemorosa sociation (see p. 141).
- 3. Myrtillus-Homogyne sociation, to the Sudetic-Hercynian facies of which two variants belong, namely Struthiopteris spicant and Calamagrostis villosa described by Zlatník from the Krkonoše Mts. and besides also Western Carpathian and Eastern Carpathian facies, regionally specific species of the latter are Aposeris foetida, Hieracium transsilvanicum, Campanula abietina.
- 4. Calamagrostis villosa sociation, as for instance described from the Krkonoše Mts. by Zlatník.

XIV. Spore plants.

In typical beech forests, the ground is wholly or practically destitute of mosses. We find them, however, on the roots, trunks, and stumps of beeches and not seldom even on stones where the soil is stony. Likewise the humid to damp beech communities are sometimes mossy (see p. 117) as may also be the «spruce» types of spurious beech forests. The epiphytic vegetation of mosses, lichens, and algae however, is usually very interesting and has been in some regions of Bohemia thoroughly studied by A. Hilitzer (2) who deals in great detail also with the ecological factors and distinguishes many sociations, some of which are specific for the beech. As far as the local distribution of these sociations on the trunk itself is concerned, we find on the beech usually on the trunk base the sociations Pyrenula nitida, Thelotrema lepadinum or Pertusaria amara, in the middle part of the trunk Parmelia saxatilis sociation, and on the upper part the Evernia prunastri sociation. Sometimes we notice on the base a differentiation of moss and lichen sociations, for instance on beech roots the sociation Isothecium myurum or Pteriginandrum filiforme, on the trunk base Thelotrema lepadinum or Graphis scripta sociation, in the middle part of the trunk Parmelia saxatilis or Cetraria glauca sociation, in the upper part Alectoria jubata sociation.

In the most simple case, there is on the trunk only one epiphytic sociation and that of indifferent sociations Parmelia physodes or Protococcus viridis, of sociations characteristic for the beech Lecanora subfusca + Phlyctis or Parmelia saxatilis. Sometimes the epiphytic vegetation is restricted to only one side of the trunk, in other instances it is on both sides, in which case at the same height the following differentiation may be observed.

Fagus

Exposed side:

Protected side:

sociation

sociation

Parmelia physodes

Algae

Parmelia saxatilis Cetraria glauca Lobaria pulmonaria

Lecidea parasema Lecanora subfasca Parmelia sulcata

Pyrenula nitida

Trente pohlia.

As examples of differentiation of the epiphytic sociations on beech Hilitzer gives the following:

1. Beeches in the virgin forest of Boubín in the Šumava Mts.

base: Isothecium myurum sociation;

lower part of trunk and protected side of the middle part: Thelotrema lepadinum sociation;

middle part of trunk, exposed side: Lobaria pulmonaria sociation; upper part of trunk: Parmelia saxatilis sociation;

branches: Alectoria jubata sociation.

2. Beeches in an old beech forest near Kdyně in Český Les Mts. base: Dicranum longifolium sociation;

lower part of trunk: Pyrenula nitida sociation;

upper part of trunk, exposed side: Parmelia saxatilis sociation; upper part of trunk, protected side: Lecanora subfusca.

As a typical succession of the epiphytic sociations in the pure beech forests, Hilitzer gives this scheme:

Base
Parmeliopsis ambigua

mosses

Exposed side

Lecanora subfuscaPhlyctis

Parmelia saxatilis

Protected side

Lecanora-Phlyctis

Parmelia saxatilis

mosses and Lobaria

In his paper on the beech forests in the neighbourhood of Kdyně, Hilitzer (¹) describes in detail the epiphytic vegetation of mosses and lichens and distinguishes 21 sociations as accompanying sociations of beech forests.

Very characteristic and different from that of the spruce forests is the mycoflora of the beech forests, especially as regards *Hymenomycetineae*. Numerous contributions on the fungi of our beech forests have been published, but notwithstanding, it is not possible at present to make a sociological analysis of the beech forest mycoflora of the whole state.

XV. Exclusive species of beech forests.

Beech forests are one of the rather exceptional communities in which we may perhaps distinguish faithful (exclusive) species, although even here the number of the absolutely exclusive species is insignificant when taking the whole Czechoslovak republic into consideration. Under special conditions, many beech forest species go over also into other sociations, however, avoid spruce forests with acid soils; many are at home in mixed spruce forests with fir, maple, and beech. Many species, and even whole communities, especially Caricetum pilosae and Melicetum uniflorae, penetrate beyond the limits of beech forests into mixed oak and hornbeam forests. Nevertheless, a rather great number of species is more or less confined to beech forests and these species may by therefore designated as beech forest species. With regard to the whole territory of our state, we can classify these species into three categories according to their more or less frequent occurence outside of the beech forests. The exclusive species of the beech forests are given in the first group.

A. Especially characteristic beech forest species.

Asperula odorata (generally in our beech forests, on all kinds of geological substrata, from foothills up to the mountain region).