Zeitschrift: Veröffentlichungen des Geobotanischen Institutes Rübel in Zürich

Herausgeber: Geobotanisches Institut Rübel (Zürich)

Band: 34 (1958)

Artikel: Some radio-carbon dates in the postglacial vegetation history of the

northern Netherlands

Autor: Zeist, W. van

DOI: https://doi.org/10.5169/seals-308102

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 28.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Some radio-carbon dates in the postglacial vegetation history of the northern Netherlands

By W. VAN ZEIST, Institute for Biological Archaeology, State University, Groningen

About a century ago the greater part of south-eastern Drenthe (Fig. 1) was covered by raised bog. However, as a consequence of intensive peat-digging only a comparatively small part of these peat deposits have been left.



Fig. 1

During the last few years a fair amount of palynological research has been carried out in this area. In this connection the radio-carbon dating of some palynological levels of more than local importance appeared to be desirable. The ¹⁴C-measurements were carried out by professor H. de Vries (Physical Laboratory, State University, Groningen).

As the results have been published elsewhere (Van Zeist 1955, 1956), they will not be discussed here in detail.

Fig. 2 shows a diagram from the Emmen region with the radio-carbon dates inserted on the right. The Preboreal/Boreal border is here placed at the increase of *Corylus*. For this border an age of about 6700 B. C. was obtained.

As there is some difference of opinion on the Boreal/Atlantic border we have submitted a sample from the middle point of the increase in Alnus and one from the intersection of the lines for Alnus and Pinus. Although the distance between both samples is less than 10 cm, the difference in age is rather great.

The level at which the curve for *Ulmus* begins to fall was dated at about 3000 B. C. At a depth of 120 cm *Plantago lanceolata* shows a conspicuous rise. On the ground of palynological investigations of burial monuments we are inclined to ascribe this increase of plantain to the invasion of a tumulus-burying people (beakers with protuding foot, cf. Van der Waals and Glasbergen 1955). In contrast with the settled farmers of the funnel beaker culture, these new immigrants were nomads, clearing large parts of the forest in order to get suitable grazing land for their cattle. The dating of about 2200 B. C. for the increase of plantain agrees well with the results of radio-carbon measurements from burial monuments of the tumulus people.

At a depth of about 60 cm Fagus reaches somewhat higher percentages. This level corresponds with the beginning of the continuous Fagus-curve in most diagrams from north-western Germany. On the evidence of the position of archaeological objects in raised bogs it could be established that in north-western Europe the first increase of Fagus corresponds approximately with the transition from the Neolithic to the Bronze Age. The sample dated at 1400 B. C. lies just above the increase of Fagus, so that according to this result the transition from the Neolithic to the Bronze Age must have taken place shortly before 1400 B. C. This is in good agreement with the dating of about 1500 B. C. for this transition arrived at on archaeological grounds.

The beginning of a more considerable increase of Fagus, constituting the Subboreal/Subatlantic border, could be dated at about 800 B. C.

In consequence of buckwheat cultivation the upper part of the peat has vanished, so that practically the whole Subatlantic period is lacking here.

From another site in the same raised bog, where less peat has vanished, the upper part of the peat profile was sampled. The most characteristic features of the diagram from this profile (Fig. 3) are the gradual decline of *Tilia* and the rise of *Fagus* and particularly of *Car*-




Fig. 2




Fig. 3

pinus. The first increase of Carpinus could be dated at about 70 B. C., whereas at about 500 A. D. this tree must have been very common in this region.

In connection with these radio-carbon dates some facts which may be of interest for the discussion of the post-glacial climate will be mentioned here briefly.

At the end of the Atlantic time Fagus arrived in the northern Netherlands. Then, for a long time, this tree was present here in small numbers. Evidently, conditions for an increase of Fagus were not favourable until the level where this tree attains a value of about 1 per cent.

There is reason to suppose that in the Netherlands it was an increase in precipitation which favoured the beech. From about the level where Fagus shows an increase the formation of Sphagnum cuspidatum peat started in various places in the raised bog of south-eastern Drenthe. The formation of this type of peat must have been the effect of an increase of rainfall, on account of which pools were formed on the surface of the bog. In these pools the hydrophilous Sphagnum cuspidatum could develop. The fact that the Sphagnum cuspidatum peat is highly fissile suggests that the pools dried up during the summer.




Fig. 4

The further increase of Fagus is preceded by another indication of a changing climate. Just before the rise of the Fagus curve the conditions for the formation of fresh Sphagnum imbricatum peat became favourable in this region. From that time on, the general humidity must have increased in such a way that in the large raised bogs the surface became sufficiently moist to establish conditions favourable for the growth of Sphagnum imbricatum.

Finally the behaviour of *Hedera* in this part of the Netherlands will be mentioned. The decline of *Ulmus* is not accompanied here by a similar fall of *Hedera* (cf. IVERSEN 1941, 1944). It is not until the first increase of *Fagus* that the *Hedera* values become considerably lower than in the preceding periods. This is shown in Fig. 4, in which the average frequency of *Hedera* is expressed as a percentage of a tree pollen sum of *Quercus*, *Tilia*, *Ulmus*, *Fraxinus*, *Fagus* and *Carpinus*. It is clear that there is a marked fall of *Hedera* after about 1500 B. C. On the ground of what has been said before it is highly improbable that this decrease of *Hedera* is to be ascribed to a more continental climate.

References

- J. IVERSEN, 1941: Landnam i Danmarks Stenalder. Danm. Geol. Unders. II. Raekke. Nr. 66.
- J. IVERSEN, 1944: Viscum, Hedera and Ilex as climate indicators. Geol. Fören. Förhandl. 66 (463—83).
- Förhandl. 66 (463-83).

 J. D. van der Waals and W. Glasbergen, 1955: Beaker types and their distribution in the Netherlands. Palaeohistoria 4 (5-46).
- in the Netherlands. Palaeohistoria 4 (5—46).

 W. van Zeist, 1955: Some radio-carbon dates from the raised bog near Emmen (Netherlands). Palaeohistoria 4 (113—8).
- (Netherlands). Palaeohistoria 4 (113—8). W. van Zeist, 1956: Die palynologische Bearbeitung des Münzfundes von Bargercompascuum. — Palaeohistoria 5 (93—9).