

Zeitschrift: Veröffentlichungen des Geobotanischen Institutes Rübel in Zürich
Herausgeber: Geobotanisches Institut Rübel (Zürich)
Band: 34 (1958)

Artikel: Radiocarbon-dated pollen sequences in eastern North America
Autor: Deevey, Edward S.
DOI: <https://doi.org/10.5169/seals-308072>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 10.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Radiocarbon-dated pollen sequences in eastern North America

By EDWARD S. DEEVEY, Osborn Zoological Laboratory,
Yale University, New Haven, Connecticut, U. S. A.

The main outlines of the vegetational history of western Europe are now quite clear, thanks mainly to the extensive and detailed work of pollen stratigraphers. Radiocarbon dating (BARENDSEN et al., 1957; GODWIN et al., 1957) tends to confirm what was already known in its broader features, and its principal value for European pollen stratigraphy is in strengthening regional correlation of pollen dates — for absolute synchrony can never be assumed in a chronology that is ultimately based on migration of plants. By contrast, the pollen chronology of northeastern North America is not only less well known but is harder to study. Postglacial time probably began 10 000 years earlier in southern New England than in the Lake Mistassini district of central Quebec, 1000 km to the north, and throughout this distance the invasion of plants, though it may have been zonally regular, obviously can not have been synchronous. Before the advent of radiocarbon dating little progress could be made in establishing regional correlation of pollen zones, and transatlantic correlations, with their important implications for the earth's climatic history, were entirely speculative.

Despite the difficulties, it has been evident since 1939 or earlier (SEARS 1932, 1948; DEEVEY 1939, 1943) that postglacial forests on both sides of the Atlantic underwent strikingly parallel changes; a BLYTT-SERNANDER-sequence, including Pre-Boreal, Boreal, Atlantic, and Sub-Atlantic climatic phases, can be discerned in most American pollen diagrams, and DEEVEY and FLINT (1957) have urged the use of CHIARUGI's term *hypsihermal* for the Boreal-through-Atlantic time of maximum warmth. Closer attention to nonarboreal pollen and to pre-Boreal deposits, inspired by the leadership of IVERSEN and FIRBAS, began in northern Maine (DEEVEY 1951), and it was immediately clear that North America's late-glacial history was also closely similar to Europe's. The sections near Presque Isle, Aroostook County, Maine, gave the first clear evidence of dominantly nonarboreal vegetation, and the tripartite or Alleröd sequence that they also showed, though not wholly convincing at the time, seems now to have been firmly established. Evidence of open-country vegetation, doubtfully identical to tundra because of the low latitude and the proximity to unbroken forest, but certainly different from taiga, has since been found at all of the few places where it has been looked for by modern methods; hence it is not true, as has been maintained by several workers, that forests grew

so close to the margin of the North American ice sheet that they left no room for tundra.

Substages of the Wisconsin glacial stage are recognized with difficulty in the New England sector of the glaciated territory, and their correlation with substages in the Middle West is uncertain, owing to absence or discontinuity of end moraines in this relatively rugged country of resistant rocks. The outer limits reached by Wisconsin ice are believed to be marked by the complex of moraines on Long Island and on Cape Cod and its offshore islands (FLINT 1953; see Fig. 1). If this complex is a correlative of the outermost Wisconsin (Farmdale — Iowan — Tazewell) complex in Ohio, Indiana, and Illinois, as Flint believed, it should have been deposited between about 21 000 and about 18 000 years ago (FLINT 1955; FLINT and RUBIN 1955). No important readvance is known north of Cape Cod, but in the long retreat across New England the ice margin paused for appreciable lengths of time at several places. One pause, in the vicinity of Middletown, Connecticut, corresponds to pollen zone T-3 (LEOPOLD 1956) and is dated by radiocarbon at about 12 800 B. P.; it is correlated with the Port Huron moraine in the Lake Michigan basin, which in turn is correlated with the Mankato maximum south of Lake Superior (WRIGHT and RUBIN 1956; FLINT 1956). Farther north in the Connecticut valley another pause is known near St. Johnsbury, Vermont; from the pollen stratigraphy at several sites in Aroostook County, Maine (DEEVEY 1951), it is inferred that this pause corresponds to zone L-3, and occurred about 10 600 B. P., the time of the Valdres maximum (formerly correlated with Mankato; see LEIGHTON 1957a, 1957b; ELSON 1957; and WRIGHT 1955, 1957) in the Lake Michigan basin.

The evidence on which these New England correlations rest is scanty, consisting as it does in pollen-stratigraphic sections from about ten localities, only six of which are actually in New England. Radiocarbon dates, though also few in number, are remarkably helpful in clarifying points that otherwise would be ambiguous. Many ambiguities remain, and the correlation chart of Figure 2, which attempts to cover an area 1200 miles wide and 700 miles from north to south, for postglacial as well as for late-glacial time, contains as much of hope and prophecy as it does of factual basis. Two important conclusions that emerge from it seem likely to be firm; both are to be credited to Miss LEOPOLD: (1) southern and central New England were forested before, during, and since Two Creeks (= Alleröd) time, and the tundra-like vegetation of Valdres (= Younger Dryas) time was confined to a belt a few miles wide; (2) at least one climatic oscillation, comparable both in character and in age to Bölling, occurred in southern New England and was

recorded outside the line of the Middletown pause. These conclusions are documented and amplified below.




Fig. 1. Sketch map of New England and adjacent Canada, showing location of pollen-stratigraphic sections outside the inferred drift border of Valdres time; ice-marginal positions according to FLINT (1953).

The suggestion that the Aroostook County climatic oscillation corresponded to that of Alleröd was made in 1951, and the zone designations (L-1, L-2, L-3) were given in the belief that these zones are late-glacial in the European sense. Just as the Younger Dryas zone represents the Fennoscandian glaciation at localities near but outside the Fennoscandian ice-margin, the younger park-tundra zone in Aroostook County is believed to represent the nearby Valdres glaciation (thought of as Mankato in 1951). This correlation (implying L-2 = Two Creeks = Alleröd zone II) rested and still rests on inference from the pollen stratigraphy, not on detailed mapping or on radiocarbon dates, for there are no end-moraines in the vicinity that can be correlated with Valdres, and the older sediments of the Aroostook County lakes are poor in organic carbon. The inference is supported, however, by the pollen diagram from Gillis Lake, Cape Breton Island, Nova Scotia (D. and B. LIVINGSTONE 1958). The Livingstones' section agrees with

the sequence in northern Maine, approximately on the same latitude, but is more convincing in its demonstration of park-tundra; moreover, it has yielded a radiocarbon date (Y-524, $10\,160 \pm 160$) that shows the upper part of zone L-3 to be of Valders age.

If the Valders ice sheet approached but did not override the Maine and Nova Scotia localities, one would not expect it to have had much influence on the climate of southern Connecticut, 450 miles from Presque Isle and 120 miles from the nearest place (Fort Edward, New York) where Valders ice is supposed to have lingered or readvanced. Miss LEOPOLD's work at Durham Meadows and Totoket Bog, in the vicinity of New Haven, Connecticut, confirmed this deduction and broke entirely new ground (LEOPOLD 1956, 1958 in press). At Durham, on outwash related to the Middletown pause, her diagrams show a brief tundra phase (T-3), followed by a long spruceforest period with two maxima of spruce pollen. The two maxima (A-2 and A-4 in the Durham spruce zone) bracket a zone (A-3) containing much pine and some oak pollen; low values of NAP were found above zone T-3. The sequence implies early postglacial afforestation and development of deciduous forest throughout Two Creeks time, interrupted by a climatic deterioration that brought a resurgence of spruce but not a reversion to tundra during Valders time. That zone A-2 was older than Two Creeks was shown by the date of W-46, $12\,700 \pm 280$ years. The Durham stratigraphy was encountered at three localities in northern Massachusetts (DAVIS 1958 in press), where Mrs. DAVIS' beautiful diagrams show characteristic differences related to the more northerly location — zone A-4 is relatively richer in spruce in Massachusetts than in Connecticut — but establish the essential correctness of Miss LEOPOLD's interpretation. Radiocarbon dates for a log in her zone A-3 at Pleasant Street Bog, Athol (W-361, $10\,800 \pm 250$; M-413, $10\,700 \pm 800$), support the inferred Valders age of the resurgence of spruce. Despite the differences in detail and the probable slight differences in age, the Durham stratigraphy has been combined with that of the Athol district to represent central New England in Figure 2.

At Totoket Bog, 10 miles south of Durham, Miss LEOPOLD also found the Durham sequence, but underneath were two older zones, the oldest (T-1) having much more abundant NAP, whereas the next-oldest (T-2) contained a «pre-Durham» spruce maximum and less NAP than the zones above and below. Evidently the pause of the ice margin at Middletown was represented by the younger NAP zone (T-3), which was complete at Totoket beyond the ice, but incomplete downward at Durham where the outwash was spread. Radiocarbon date Y-285 ($13\,550 \pm 460$), though it now seems slightly too old, verified the correlation of zone A-2 at Totoket with that at Durham; two attempts have been

made since 1955 to date the older zones. The 1956 studies (BARENDSEN et al., 1957) gave results which are systematically too old, probably because of inexperience with a new boring instrument, with resulting disturbance of the stratification. More careful studies in 1957 (DEEVEY, unpublished) give acceptable ages for zones T-2 (Y-502, $13\,280 \pm 420$) and T-3 (Y-505, $12\,350 \pm 400$), as well as confirmatory dates, listed below, for zones A-2 and A-4.

AGE B.P.	AGE AD/BC	MICHIGAN Potzger, Andersen	SOUTHERN CONNECTICUT Leopold	SOUTH-CENTRAL NEW ENGLAND Leopold, Davis	NORTHERN MAINE AND NOVA SCOTIA Deevey, Livingstone	CENTRAL QUEBEC AND ONTARIO Ignatius	GLACIAL SUBSTAGES	PROBABLE EUROPEAN EQUIV. Dates after Godwin et al.	AGE AD/BC
1000	1000	C3b SPRUCE, PINE RISE	C3b SPRUCE, PINE RISE	C3b SPRUCE, PINE RISE	C3b BEECH DECLINES	3b PINE, SPRUCE			1000
2000	0	C3a OAK, ELM, MAPLE, HEMLOCK	C3a OAK, HEMLOCK, CHESTNUT ● C37, C119	C3a OAK, HEMLOCK	C3a BIRCH, HEMLOCK, BEECH	3a SPRUCE, PINE		IX SUB-ATLANTIC	0
3000	1000	2nd beech maximum	2nd beech maximum	2nd beech maximum					1000
4000	2000	C2 OAK, HICKORY ● M291, C349	C2 OAK, HICKORY	C2 OAK, HICKORY	C2 PINE, BIRCH, OAK	2c PINE, BIRCH		VIII SUB-BOREAL	2000
5000	3000	1st beech maximum	1st beech maximum ● C38, C120	1st beech maximum					3000
6000	4000	C1 OAK, WHITE PINE, HEMLOCK ● Y1698	C1 OAK, HEMLOCK	C1 OAK, HEMLOCK	C1 BIRCH, HEMLOCK ● C335	2b SPRUCE, WHITE PINE, HEMLOCK (tr)		VII ATLANTIC	4000
7000	5000	B2 PINE		B2 PINE, OAK	B PINE	2A BIRCH, SPRUCE ● Y222 ● Y223			5000
8000	6000	B1 JACK PINE, ELM ● M286	B PINE ● C39	B1 PINE, SPRUCE ● Y282	A2 SPRUCE, BIRCH	1 ALDER, PARK-TUNDRA		VI TRANSITION	6000
9000	7000							V BOREAL	7000
10000	8000		A4 SPRUCE RETURNS ● Y4470 ● Y504	A4 SPRUCE RETURNS ● W361, M413	A1 BIRCH, SPRUCE ● Y324			IV PRE-BOREAL	8000
11000	9000	A SPRUCE, FIR (undifferentiated) ● M288A ● Y293B		A3 PINE, SPRUCE	L3 PARK-TUNDRA	(GLACIATED)	VALDERS	III YOUNGER DRYAS	9000
12000	10000		A3 PINE, SPRUCE, OAK ● Y503		L2 SPRUCE, PARK-TUNDRA		TWO CREEKS INTERVAL	II ALLERÖD	10000
13000	11000	T3 BIRCH PARK-TUNDRA T2 SPRUCE, PARK-TUNDRA	A2 BIRCH, SPRUCE ● Y505	A2 BIRCH, SPRUCE A1 ● W45	L1 TUNDRA		PORT HURON = MANKATO?	Ic OLDER DRYAS	11000
			A1 ● Y4474 ● Y502 ● Y285	T3 TUNDRA ● W45	(GLACIATED)			Ib BÖLLING	
			T1 TUNDRA	(GLACIATED)				Ia OLDEST DRYAS	
	12000								12000

Fig. 2. Correlation of pollen zones in eastern North America, with time scale provided by radiocarbon dates. The zone boundaries are drawn on the basis of averages and estimates, hence the radiocarbon-dated samples do not necessarily fall in their proper zones, but are plotted at their correct chronologic positions as published.

Totoket Bog thus provides the clearest evidence for the present view, that zone T-3 records a time of glacial pause or readvance of Port Huron age, following a period of ameliorating climate — represented

in the Lake Huron basin by the Arkona retreat — that is similar to that of Bölling. Because of the large standard errors of single radiocarbon dates, and perhaps because the carbon in a late-glacial clay-gyttja sample is not invariably contemporary with the pollen grains, this evidence is not decisive. It is supported by other stratigraphic data, however:

1. At Red Maple Swamp, near New London, Connecticut, on the latitude of Totoket, an unpublished pollen diagram by Nellie BEETHAM shows the Totoket sequence, beginning in zone T-2. Radiocarbon dates for zones A-2 and A-4 substantiate the correlation, but the older zones are poor in carbon.

2. On Martha's Vineyard, off Cape Cod, unpublished work by J. G. OGDEN strongly suggests that the late-glacial sequence there belongs to an older cycle of amelioration and glacial readvance than the one at Totoket — as it should if the ice-marginal correlations shown in Figure 1 are correct. The Martha's Vineyard data are too new to have been included in Figure 2.

3. Southern Michigan, like southern and central Connecticut, seems to have been forested during Two Creeks time, and climatic changes during Valders time disturbed but did not destroy the forest. This inference was made by S. T. ANDERSEN (1954), but the sequence at South Haven, Michigan (ZUMBERGE and POTZGER 1956), included in Figure 2 because it yielded several radiocarbon dates, is more problematical in its late-glacial portion than is ANDERSEN's sequence from the George Reserve pond.

Little can be said as yet about glacial events subsequent to the Valders maximum; the problem of a Cochrane substage (KARLSTROM 1956) is still unresolved. Unpublished work by Heikki IGNATIUS in the Lake Mistassini and Chibougamau districts of central Quebec, and in the region of Cochrane, Ontario, shows that the pollen sequences began some time after the Valders maximum and before about 7000 B. P.; the date of W-36 (RUBIN and SUESS 1955) is relevant here, as are those of Y-222 (6730 ± 200) and Y-223 (6960 ± 90), which came from IGNATIUS' sections. It is a reasonable guess that the Cochrane readvance is correlated with the Boreal-Atlantic transition, and is recorded far to the south in the United States by the mesic conditions of zone C-1.

In summary, Figure 2 reports a number of local sequences and their probable correlations, made by the classic methods of pollen stratigraphy. Its absolute time scale is only approximate, and is arrived at by considering (but not by blindly relying on) the following radiocarbon dates and their means:

Zone	Locality	Sample	Date	Average
C—3 (bottom)	Upper Linsley, Conn.	C—37, C—110	1970 \pm 360	1970
C—2 (bottom)	Upper Linsley, Conn.	C—38, C—120	5230 \pm 180	5230
B (top)	Plissey Pond, Maine	C—335	5960 \pm 320	5960
B (middle?)	Upper Linsley, Conn.	C—39	8320 \pm 400	c. 8020
	Durham, Conn.	W—45	7570 \pm 250	
	Durham, Conn.	Y—282	8155 \pm 410	
	[Totoket, Conn., 1956]	[Y—446g]	[9650 \pm 90]	
	(cf. So. Haven, Mich.)	(M—288)	(7925 \pm 400)	
L—3 (top)	Gillis Lake, N. S.	Y—524	10,160 \pm 160	10,160
A—4 (middle?)	Red Maple Swamp, Conn.	Y—447c	10,480 \pm 140	10,460
	Totoket, Conn., 1957	Y—504	10,440 \pm 200	
	[Totoket, Conn., 1956]	[Y—446f]	[12,080 \pm 300]	
A—3 (middle?)	Athol, Mass.	W—361 (M—413)	10,800 \pm 250	10,800
A—2 (middle?)	Durham, Conn.	W—46	12,700 \pm 280	c. 12,780
	Red Maple Swamp, Conn.	Y—447d	13,290 \pm 120	
	Totoket, Conn., 1955	Y—285	13,550 \pm 460	
	Totoket, Conn., 1957	Y—503	11,590 \pm 200	
	[Totoket, Conn., 1956]	[Y—446d]	[14,790 \pm 160]	
T—3	Totoket, Conn., 1957	Y—505	12,350 \pm 400	c. 12,800
T—2	Totoket, Conn., 1957	Y—502	13,280 \pm 420	c. 13,500

References

- ANDERSEN, S. T., 1954: A late glacial pollen diagram from southern Michigan, USA. — *Danm. Geol. Unders.*, (II), no. 80:140—155, pl. 8, 13.
- BARENDSEN, G. W., DEEVEY, E. S., and GRALENSKI, L. J., 1957: Yale natural radiocarbon measurements III. — *Science*, 126:908—919.
- DAVIS, M. B., 1958 in press.: Three pollen diagrams from central Massachusetts — *Amer. J. Sci.*
- DEEVEY, E. S., 1939: Studies on Connecticut lake sediments. I. A postglacial climatic chronology for southern New England. — *Amer. J. Sci.*, 237:691—724.
- 1943: Additional pollen analyses from southern New England. — *Amer. J. Sci.*, 241:717—752.
- 1951: Late-glacial and postglacial pollen diagrams from Maine. — *Amer. J. Sci.*, 249:177—207.
- and FLINT, R. F., 1957: Postglacial hypsithermal interval. — *Science*, 125:182—184.
- ELSON, J. A., 1957: Lake Agassiz and the Mankato-Valders problem. — *Science*, 162:999—1002.
- FLINT, R. F., 1953: Probable Wisconsin substages and late-Wisconsin events in northeastern United States and southeastern Canada. — *Geol. Soc. Amer., Bull.*, 64:897—919, 3 pl.
- 1955: Rates of advance and retreat of the margin of the late-Wisconsin ice sheet. — *Amer. J. Sci.*, 253:249—255.
- 1956: New radiocarbon dates and late-Pleistocene stratigraphy. — *Amer. J. Sci.*, 254:265—287.
- and RUBIN, Meyer, 1955: Radiocarbon dates of pre-Mankato events in eastern and central North America. — *Science*, 121:649—658.
- GODWIN, H., WALKER, D., and WILLIS, E. H., 1957: Radiocarbon dating and post-glacial vegetational history: Scaleby Moss. — *Roy. Soc. London, Proc.*, (B), 147: 352—366.
- KARLSTROM, T. N. V., 1956: The problem of the Cochrane in late Pleistocene chronology. — *U. S. Geol. Surv., Bull.* 1021—J:303—331, pl. 31.

- LEIGHTON, M. M., 1957a: The Cary-Mankato-Valders problem. — *J. Geol.*, 65:108—111.
— 1957b: Radiocarbon dates of Mankato drift in Minnesota. — *Science*, 125: 1037—1038.
- LEOPOLD, E. B., 1956: Two late-glacial deposits in southern Connecticut. — *Nat. Acad. Sci., Proc.*, 42:863—867.
— 1958: in press.
- LIVINGSTONE, D. A., and LIVINGSTONE, B. G. R., 1958: Late-glacial and post-glacial vegetation from Gillis Lake in Richmond County, Cape Breton Island, Nova Scotia. — *Amer. J. Sci.*, 256:341—359.
- RUBIN, Meyer, and SUESS, H. E., 1955: U. S. Geological Survey radiocarbon dates II. — *Science*, 121:481—488.
- SEARS, P. B., 1932: Postglacial climate in eastern North America. — *Ecology*, 13:1—6.
— 1948: Forest sequence and climatic change in northeastern North America since early Wisconsin time. — *Ecology*, 29:326—333.
- WRIGHT, H. E., 1955: Valders drift in Minnesota. — *J. Geol.*, 63:403—411.
— 1957: Radiocarbon dates of Mankato drift in Minnesota. — *Science*, 125:1038—1039.
— and RUBIN, Meyer, 1956: Radiocarbon dates of Mankato drift in Minnesota. — *Science*, 124:625—626.
- ZUMBERGE, J. H., and POTZGER, J. E., 1956: Late Wisconsin chronology of the Lake Michigan basin correlated with pollen studies. — *Geol. Soc. Amer., Bull.*, 67: 271—288, pl. 1.