

Zeitschrift: Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübel, in Zürich

Herausgeber: Geobotanisches Institut, Stiftung Rübel (Zürich)

Band: 41 (1968)

Artikel: The nitrogen relations and other ecological investigations on wet fertilised meadows

Autor: Williams, John Trevor

Kapitel: F: Summary

DOI: <https://doi.org/10.5169/seals-308317>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 12.10.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

F. Summary

The nitrogen relations and other ecological investigations on wet fertilised meadows

Section A—*Systematics*

The present sociological nomenclature was examined. The wet fertilised meadows investigated are described as the associations *Polygono-Cirsietum oleracei* (TÜXEN and PREISING, 1951) and *Bromo-Senecionetum*. There appears to be an edaphic difference between the two associations. Both are subdivided by some authors. For experimental purposes it was considered more meaningful to regard the wet fertilised meadows as a single group: the *Cirsietum oleracei medioeuropaeum*.

An ordination of vegetation tables revealed that the *Cirsietum* was not clearly distinct from the wetter forms of the *Arrhenatheretum*. There were also montane varieties of the community, but it is probably better to regard these simply as variants of the community. The subdivision of the *Molinio-Arrhenatheretea* is clearly too artificial. Probably in nature a multidimensional continuum exists, the named associations are representing parts of the continuum which are distinguishable floristically.

The distribution of the *Cirsietum oleracei medioeuropaeum* was considered. It seemed to be of subatlantic origin, and the present distribution must be correlated with management practices. Various facets of the physiognomy of the community were described.

Section B—*Soil factors*

Three experimental areas were selected, i.e. (a) Swiss Midlands, (b) Alsace, (c) Baden-Württemberg. These offered contrasting soil conditions.

The two major controlling factors were water and nitrogen. Ground water and soil moisture content were examined, and the latter was correlated with the soil water tension. Studies of the "available water" by determining the yearly variations in suction force showed that water was rarely a limiting factor.

The nitrogen relations were studied by measuring the seasonal pattern of nitrate- and ammonium-contents in the field and by measuring the seasonal pattern of potential and actual nitrification. There were similarities in the patterns of both potential and actual nitrification in the three areas, although they were modified locally in response to climate.

Various factors affecting the nitrogen budget were investigated in the laboratory, e.g. aeration, water content, calcium carbonate content, ammonium supply and the distribution of nitrifying activity horizontally and vertically in the soil. Populations of the nitrifying bacteria (nitrite and nitrate producers) were cultured and the fluctuations in the population size was related to the pattern of nitrification. These experiments were discussed in relation to the nitrogen budget of the community, and it appeared that along with water, nitrogen was a major factor controlling the stability of the community.

Soil phosphorus was also determined and the content showed a characteristic seasonal pattern as did pH and water content. Phosphorus may be a factor likely to cause the differentiation into some vegetation variants.

Section C—*Productivity*

The productivity of the wet fertilised meadows was measured both quantitatively and qualitatively. From sample cuts the hay yield was estimated and the hay was fractionated into three portions important from the agricultural point of view: grasses, legumes and herbs. Protein and fibre contents were measured.

Since mineral nutrition was found to be important (Section B), the growth of several species, typical and constant of the community, was measured in sand culture. Their responses to varying levels of nitrogen, phosphorus and calcium were tested. The results were discussed in relation to the role played by these macronutrients in the light of the results in Section B.