

Zeitschrift: Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübel, in Zürich

Herausgeber: Geobotanisches Institut, Stiftung Rübel (Zürich)

Band: 59 (1976)

Artikel: Quercus pubescens : Wälder und ihre ökologischen Grenzen im Wallis (Zentralalpen)

Autor: Burnand, Jacques

Kapitel: Summary

DOI: <https://doi.org/10.5169/seals-308483>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Summary

=====

The Quercus pubescens forests in Valais (Swiss Central Alps) and their ecological limits

On the background of the general problem of the limits between deciduous (and mediterranean type) and coniferous forests, ecological investigations were made from 1970 to 1974 in the Quercus pubescens and Pinus silvestris forests in Valais, a region with continental climate where these two species are the only important forest trees in the lower parts. The aim was the determination of the factors producing the forest pattern. The factors investigated were the human influence on the stands, the climate (especially the irradiation), the soil factors and the distribution of the two trees. Besides, a phytosociological investigation of the oak woods was carried out after the method of Braun-Blanquet.

Results

On the base of a model, it has been shown that the distribution is mainly determined by irradiation and altitude. The oaks are restricted to the sites with strong irradiation: the minimum irradiation required increases with increasing altitude. The oaks further avoid the depressions where late spring frosts occur. The pines grow on all sites where the oaks cannot exist (Chapter 3.15).

Parent rock conditions do not play an important role in the distribution of the two species, which are both equally drought resistant. The nutrient supply in the soils is never low (3.13).

As far as they are not destroyed, the forests suffered the influence of human activities (fire, pasture, coppice selection system) for centuries. Some oak woods were transformed into pine stands or into composite forests with oaks in the scrub layer and pines in the tree layer. Today, as every

human activity has stopped in these forests, they are regenerating towards a more natural composition (3.11).

Other mountain regions exist with relatively continental climate and where, during the growth period, there is intense irradiation, but also sufficient quantities of precipitations. The comparison shows that there, too, the distribution of oaks and conifers is determined by the same factors as in Valais. These regions are: the other inner valleys of the central and western Alps (with the same species as in Valais), the mountains of the south-western United States (Quercus and Pinus), the inner valleys of the north-western Himalayas (Quercus, Abies, Pinus) and some Caucasus valleys (Quercus, Conifers). This model, therefore, seems to fit as well to deciduous as to other oak species (3.23, 3.24).

From the phytosociological point of view, the Valais oak woods can be divided into two associations according to climate and parent rock conditions: into the Campanulo trachelii-Quercetum pubescentis prov. under less continental climate at the limit towards the outer Alps, on silicate soils, and into the Saponario-Quercetum pubescentis prov. in the centre of Valais. The subdivisions also follow a complex water gradient influenced by climate and soils (4.1). The comparison of these associations with other existing units is made difficult by the present development of the stands and by the different dates of the relevés and must therefore be considered only provisional (4.2.).

In the last chapter, the conservation of the most outstanding places with oak woods is suggested.