

Zeitschrift: Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübel, in Zürich

Herausgeber: Geobotanisches Institut, Stiftung Rübel (Zürich)

Band: 66 (1979)

Artikel: Soziologisch-ökologische Untersuchungen im Grenzbereich *Fagus silvatica* L. / *Pinus silvestris* L. in der nördlichen Schweiz

Autor: Roth, Christian

Kapitel: Summary

DOI: <https://doi.org/10.5169/seals-308568>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 28.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Summary

The present paper deals with phytosociological and pedological investigations in natural as well as anthropogenous forests of the *Molinio-Pinetum* from Swiss Midlands. The author's results having been compared with the previous data, the following classification of the *Molinio-Pinetum* is suggested:

- a) subassociation with
 - 1) *Epipactis palustris* - variant
 - 2) *Listera ovata* - variant
 - 3) *Genista tinctoria* - variant
 - 4) *Sanguisorba minor* - variant
- b) subassociation with *Laserpitium latifolium*
 - 5) typical variant
 - 6) *Geranium sanguineum* - variant
 - 7) *Parnassia palustris* - variant

The variants described under 1 - 3 represent open forests with well-developed herb layer containing numerous *Graminae* and *Cyperaceae*. The variant No. 4 cannot be fully compared to them, for a different method has been used for the resp. relevés. The variants No. 5 - 7 are natural forests occurring on steep slopes of northwestern Swiss Midlands; the variant No. 6 appears on drier soils than the typical variant, but its exposure is most frequently the same, i.e. southern or western. The variant No. 7 occurs on N-exposed steep slopes within the ridge of Albis, in sites that are permanently wet due to seeping soil water.

All the studied variants are characterized by the absence of *Fagus silvatica* in the tree layer, in spite of the fact that the beech represents the climax forest species of colline and submontane zone within the studied area. The present investigations show that the *Molinia-Pinus* forests inhabit the marly rendzina that is younger and shallower than soils of the neighbouring forests of *Fagus*. Marly rendzina develops very slowly; its uppermost layer may sometimes be removed by landslides which results in recurrent successions. *Fagus silvatica* is able to germinate in the *Molinio-Pinetum*, its life span being about 30 years; however, only exceptionally grow the individuals taller than 50 cm and so they nearly always remain within the herb layer. The abnormally reduced growth is apparently caused by properties of the marl soil that may remain quite wet for some time, but rapidly manifests a strong water deficiency in dry periods. The shallowness of marly rendzina as well as the above mentioned qualities of the marl may accordingly be considered as principal factors responsible for the absence of *Fagus silvatica* in the *Molinia-Pinus* forests occurring within the climax zone of this species.