

Zeitschrift: Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübel, in Zürich

Herausgeber: Geobotanisches Institut, Stiftung Rübel (Zürich)

Band: 69 (1980)

Artikel: "luncus trifidus" L. subsp. "carolinianus" Hämet-Ahti, n. subsp., in eastern North America

Autor: Hämet-Ahti, Leena

DOI: <https://doi.org/10.5169/seals-308595>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 26.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Juncus trifidus L. subsp. *carolinianus* Hämet-Ahti,
n. subsp., in eastern North America

by

Leena HÄMET-AHTI

Juncus trifidus L. subsp. *carolinianus* Hämet-Ahti, n. subsp.

Type: U.S.A. North Carolina. Buncombe Co.: Craggy Gardens, 1897 Biltmore 5546 (NCU, holotype; GH, NY, S, isotypes).

Dense caespitosus; caules (13-)16-30(-37) cm alti; folia basilaria sine lamine vel breviter laminigera; folia caulina 3; flores 1-2(-3), tepala 3.5-4 mm longa; fructus perigonum aequans vel subaequans.

Densely caespitose perennial; fruiting stem (13-)16-30(-37) cm tall; basal sheaths 3-4 cm long; lamina up to 1 cm long or missing; the upper leaves, usually three, much exceeding the inflorescence, the base of the lowermost situated 1-3 cm below the inflorescence, the two uppermost usually subtending the inflorescence; flowers 1-2(-3), not usually densely crowded; perianth segments 3.5-4 mm long, equal or outer ones slightly longer; capsule equal to or slightly shorter than the perianth, with ca. 0.7 mm long mucro; seeds 1.0-1.3 mm long including the appendages.

Juncus trifidus is an arctic-oroarctic ("arctic-alpine") amphi-atlantic species occurring in westernmost Eurasia and northeasternmost North America (HULTÉN 1958). It deviates to some extent from the other members of the genus *Juncus* by having, for instance, lacerate auricles and bracteoles and serrulate leaf margins, which also occur in some other genera of Juncaceae (cf. NILSSON & SNOGERUP 1972). It has been usually regarded as a fairly uniform species and the only commonly accepted race is subsp. *monanthos* (Jacq.)

Asch. & Graebner, occurring in limestone areas in the Alps, Apennines and Balkans (e.g. JANCHEN 1960, HESS et al. 1967).

The Appalachian specimens of *J. trifidus* differ from *J. trifidus* s. str. in being taller, producing usually only 1-2 flowers and in bearing a perianth often equal to slightly exceeding the capsule (Tab. 1, Fig. 1,2). They have been usually included in subsp. (or var.) *monanthos* (e.g. FERNALD 1950, RADFORD et al. 1968) though, for instance, LöVE and LöVE (1965, 1966) and BÖCHER (1972) did accept this treatment. However, the Appalachian material is really distinct from subsp. *monanthos* which is taller, has longer leaves, especially basal leaves and sometimes a stem leaf, and which bears its lowest summit leaf far below the inflorescence. Also the second leaf of subsp. *monanthos* may be distinctly separated from the one-flowered inflorescence, the perianth is up to 5 mm long, and the seeds 1.6 - 2.0 mm long (Tab. 1, Fig. 1). (The measurements were very difficult to obtain because most of the herbarium specimens were too young or too old or damaged, etc.).

The first report of the occurrence of subsp. *monanthos* in North America was probably published by ROBINSON & FERNALD (1908), being apparently based on BUCHENAU's (1906) note that in North America the intermediates between *J. trifidus* s.str. and subsp. *monanthos* are common. Obviously the Appalachian specimens of *J. trifidus* belong to a distinct taxon rather than to subsp. *monanthos* or subsp. *trifidus*. The best rank for it seems to be the subspecific level.

Also ecologically subsp. *carolinianus* and subsp. *monanthos* are different: subsp. *monanthos* occurs only in oroboreal limestone areas (cf. JANCHEN 1960, HESS et al. 1967) whereas subsp. *carolinianus*, like subsp. *trifidus*, prefers oligotrophic schistose rock crevices. The northernmost localities of subsp. *carolinianus* seem to be in the oroboreal zone above the timberline, but in the southernmost part of its range there is no oroboreal area and it occurs in the forested oroboreal zone (obviously hemioroboreal or lower oroboreal) on some tops of the Blue Ridge Mountains.

The number of flowers has often been regarded as a distinguishing feature between subsp. *monanthos* and subsp. *trifidus*. However, the number of flowers in subsp. *trifidus* may vary: NILSSON & SNOGERUP (1972) reported one-flowered specimens from the limestone areas at lower altitudes in southern Norway, and




Figure 1. Schematic diagrams of *Juncus trifidus* subsp. *trifidus* (1), subsp. *carolinianus* (2) and subsp. *monanthos* (3).

a = the basal sheaths, b = the lamina of the basal sheaths, c = the lowermost summit leaf, d = the middlemost summit leaf, e = the uppermost summit leaf.

Table 1. A comparison of some characters of *Juncus trifidus* subsp. *trifidus* (material from northernmost Fennoscandia), subsp. *carolinianus* (from the Appalachian Mts.) and subsp. *monanthos* (from Switzerland).

	subsp. <i>trifidus</i>	subsp. <i>carolinianus</i>	subsp. <i>monanthos</i>
length of plant cm	19 ± 5.2 (n 25)	23 ± 7.5 (n 30)	30 ± 5.8 (n 24)
length of basal sheaths cm	3.0 ± 0.6 (n 25)	2.9 ± 1.3 (n 27)	4.0 ± 1.2 (n 24)
length of lowermost summit leaf cm	6.5 ± 1.8 (n 24)	7.7 ± 2.4 (n 18)	9.0 ± 1.9 (n 20)
distance between base of lowermost summit leaf and inflorescence cm	1.8 ± 1.3 (n 24)	2.1 ± 1.3 (n 25)	6.0 ± 3.3 (n 21)
length of middlemost summit leaf cm	5.2 ± 1.0 (n 24)	6.5 ± 1.7 (n 35)	7.0 ± 2.2 (n 21)
distance between base of middlemost summit leaf and inflorescence cm	0	(0 - 0.5)	1.3 ± 0.5 (n 21)
length of uppermost summit leaf cm	2.6 ± 1.1 (n 24)	2.6 ± 1.7 (n 36)	3.3 ± 2.9 (n 21)
number of flowers per culm	2.3 (n 78)	1.5 (n 74)	1.1 (n 146)
length of perianth segments mm	3.7 ± 0.3 (n 11)	4.0 ± 0.3 (n 23)	4.8 ± 0.2 (n 21)
mucro of capsule mm	0.7 - 0.8	0.7 - 0.8	1.0 - 1.5

LÖVE & LÖVE (1966) doubted the taxonomic value of this character. Also, in northern Fennoscandia there are often solitary one-flowered stems side by side with three-flowered stems, but the northern Finnish colonies growing on serpentine rocks as well as on river banks below the timberline are mainly one-flowered. Both these habitats are wet seepages, while, above the timberline, the usual sites of *J. trifidus* are very dry. However, these specimens do not represent subsp. *monanthos* or subsp. *carolinianus*, but perhaps instead some local ecotypes occurring in ecologically special sites. Also, in the Canadian material of *J. trifidus* the one- and two-flowered specimens are not very rare.

J. trifidus s. str. has been reported from New York (e.g. FERNALD 1950), but

all the examined specimens can be included in subsp. *carolinianus* rather than subsp. *trifidus*, though some of them are not very typical. The question, whether any of the Canadian *J. trifidus* are fully identical with subsp. *trifidus*, needs further study, as will the exact location of the northern limit of subsp. *carolinianus*.



Figure 2. *Juncus trifidus* subsp. *carolinianus* (*Saxifraga michauxii* in the left). North Carolina, Blue Ridge Mts., Craggy Garden (apparently the type locality), August 1978 (photo L. HäMET-AHTI).

Acknowledgements

I am very grateful to all the organizers, leaders and guides of the 16th IPE for the very interesting excursion and especially to Dr. Jan Pittillo who acquainted me with *Juncus trifidus* in the Blue Ridge Mountains. I am also obliged to the curators of the following herbaria for loaning specimens: BERN, GH, NCU, NY, S, US, and to Mr. Trevor Goward, Vancouver, B.C., for revising the English language.

Summary

The Appalachian taxon *Juncus trifidus* L., usually included in the European var. *monanthos* (Jacq.) Bluff & Fingerh., is recognized as a new subspecies, subsp. *carolinianus* Hämet-Ahti.

Zusammenfassung

Die Pflanzen von *Juncus trifidus* L. aus den Appalachen, bisher meistens in die europäische var. *monanthos* (Jacq.) Bluff & Fingerh. eingeschlossen, wurden als neue Unterart, subsp. *carolinianus* Hämet-Ahti, eingestuft.

References

- BÖCHER, T.W., 1972: Evolutionary problems in the Arctic flora. In: VALENTINE, D.H. (ed.), Taxonomy, phytogeography and evolution. Acad. Press London, New York. 101-113.
- BUCHENAU, F., 1906: Juncaceae. In: ENGLER, A., Das Pflanzenreich IV, 36, 1-284.
- FERNALD, M.L., 1950: Gray's manual of botany. American Book Company, New York etc. 1632 pp.
- HESS, H.E., LANDOLT, E. and HIRZEL, R., 1967: Flora der Schweiz I. Basel, 858 pp.
- HULTÉN, E., 1958: The Amphi-Atlantic plants and their phytogeographical connections. Kungl. Svenska Vetenskapsakad. Handl. (Ser. IV), 7 (1), 1-340.
- JANCHEN, E., 1960: Monocotyledones. Catalogus Floraes Austriae I, 4, 711-881.
- LÖVE, A. and LÖVE, D., 1965: Taxonomic remarks on some American alpine plants. Univ. Colorado Studies, Ser. Biol. 17, 1-43.
- 1966: Cytotaxonomy of the alpine vascular plants of Mount Washington. Univ. Colorado Studies, Ser. Biol. 24, 1-74.

- NILSSON, O. and SNOGERUP, S., 1972: Drawings of Scandinavian plants. 65.
Juncus trifidus L. 1753. Bot. Not. 125, 1-2.
- RADFORD, A.E., AHLES, H.E. and BELL, C.R., 1968: Manual of the vascular flora
of the Carolinas. Chapel Hill, N.C., 1183 pp.
- ROBINSON, B.L. and FERNALD, M.L., 1908: Gray's manual of botany. American
Book Company, New York etc. 926 pp.

Address of the author: Dr. Leena HäMET-AHTI
Department of Botany
University of Helsinki
Unioninkatu 44
SF-00170 Helsinki 17, SUOMI-FINLAND