

Zeitschrift: Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübél, in Zürich

Herausgeber: Geobotanisches Institut, Stiftung Rübél (Zürich)

Band: 75 (1981)

Artikel: Phenological methods in permanent plot research : the indicator value of phenological phenomena : a study in limestone grassland in Northern Switzerland = Phänologische Methoden bei Dauerquadratuntersuchungen : der prognostische Wert phänologischer Phänomene : eine Untersuchung in Halbtrockenrasen (Mesobrometum) in der Nordschweiz

Autor: Krüsi, Bertil

Kapitel: 1: Introduction

DOI: <https://doi.org/10.5169/seals-308658>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 26.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

1. Introduction

Changes in plant communities due to successional development or alterations in the management can be studied by short-term research, communities representing various stages of succession or different management forms being compared; alternatively, a long-term programme dealing with permanent plots can be established. Either of these approaches has its disadvantages: the predictive value of short-term investigations is not always infallible, whereas long-range studies are time-consuming and highly influenced by weather conditions.

Additional information thus being required to predict the future of the vegetation within a given site preferably after only a few years, phenological data seems to be helpful. Modifications of the physical environment due to a successional development or alterations in management may render the site favourable for some species yet disadvantageous for others. In earlier stages of succession, species not adapted to the new conditions most frequently show reduced vegetative vigour, blooming and fruiting (HARPER and OGDEN 1970, HARPER 1977) but they still persist, actual changes in the composition of the vegetation only later being observed. Phenological data is thus expected to reflect environmental changes earlier than the traditional relevés which are used in phytosociology (e.g. those of the Braun-Blanquet type).

Without doubt, man has always paid attention to the phenology of some particularly important species. However, symphenological records comprising the phenology of all or most species within a given plant community date back solely to the beginning of the present century (SALISBURY 1916, 1918, GAMS 1918, ALECHIN 1925, SCHENNIKOW 1927, 1932). Since then, phenological development of whole communities and their individual components has been described by numerous authors both verbally (e.g. BECKER 1941, WALTER 1968, BYKOW 1974) as well as diagrammatically (e.g. ELLENBERG 1939, FÜLLEKRUG 1967, 1969, BALÁ-TOWÁ-TULÁČKOWÁ 1970b, 1971, DIERSCHKE 1972, 1974, 1977, FALIŃSKA 1972, 1973a, b, 1975, 1976, KRÜSI 1977, 1980). Various ways of recording and presenting phenological data were reviewed by BALÁ-TOWÁ-TULÁČKOWÁ (1970b) and DIERSCHKE (1972, 1977). The bibliography of symphenological diagrams was compiled by

BALÁTOWÁ-TULÁCKOVÁ (1970a) and later by TÜXEN and WOJTERSKA (1977).

It should be noted, however, that phenology is obviously not restricted to merely describing the developmental rhythm of plants or plant communities, but represents an important auxiliary science in various research fields, viz. climatology (e.g. SCHNELLE 1955, ELLENBERG 1956a, SCHREIBER 1968, 1977, HEGG 1967, 1977, LIETH 1974), taxonomy (e.g. MARCET 1956, FALIŇSKA 1974, 1978) or ecosystem analysis (e.g. ELLENBERG 1956b, FALIŇSKA 1978). As far as vegetation science is concerned, phenological observations are useful in delineating phytosociological units (e.g. ZOLLER 1954, HEJNÝ 1978) or revealing some developmental trends (e.g. WELLS 1971). The latter possibility is well known amongst students of vegetation. According to BRAUN-BLANQUET (1964), the direction of development of communities is often first heralded by changes in vigour of particular species. RABOTNOV (1969) considered the decrease in vigour of mature plants and not the actual changes in the number of individuals of a given species as an infallible indicator of its deteriorating life conditions. This aspect has also been emphasized by other authors (e.g. BOTTLÍKOVÁ 1973, FALIŇSKA 1975).

Save for the Russian school (see HARPER and WHITE 1974 as well as GATSUK et al . 1980 for a bibliography), little work has hitherto been undertaken using phenological information to predict developmental trends in plant communities, one of the rare exceptions being the investigation of WELLS (1971). A study was therefore undertaken to examine the possible indicative value of short-term and mid-term phenological observations carried out in limestone grassland ecosystems in northern Switzerland. On one hand, phenological responses of some species towards environmental changes were studied, and on the other hand, phenological behaviour of whole communities was observed in this respect. The present paper deals with the first results obtained in the course of these investigations.

Acknowledgements

It is a pleasure to acknowledge the help of a large number of people. Thanks are due to Professor Dr E. LANDOLT who stimulated me to undertake this study. Very special thanks are addressed to Professor Dr K. Urbanska who revised the text and offered constructive criticism. Professor Dr F. Klötzli and Dr W. DIETL (Swiss Federal Research Station for Agronomy, Zürich-Reckenholz)

advised on phytosociological problems, Dr O. WILDI (Swiss Federal Institute of Forestry Research SFIFR) helped with the mathematical analysis and Dr H. FUNK (Geological Institute ETH) with geological nomenclature. The advice of Ms C. BROWN, who made many valuable suggestions on style and use of the English language is greatly appreciated. I have had useful discussions on various points with PD Dr A. GIGON. Mr E. SCHÄFFER assisted in the field, and Ms A. HONEGGER typed the manuscript; my sincere thanks are addressed to all these persons as well as to numerous colleagues from the Geobotanical Institute who occasionally helped throughout the course of the work. The financial support of the Swiss Federal Institute of Technology (SFIT), Zürich, Switzerland, is gratefully acknowledged.

2. Description of the study areas

The four study areas are localized in northern Switzerland, within the Jurassic mountains belonging to the community of Merishausen 7.5 km NNW of Schaffhausen (National Grid Reference 688 500 / 291 000, Fig. 1). The substratum consists of Upper Jurassic limestone. The soils are of a mull-like rendzina type; the content of calcium carbonate within the uppermost 5 cm of soil ranging from 29 to 60 per cent, the corresponding pH values vary from 7.6 to 8.0. Climatic conditions are diagrammatically presented in Fig. 2.

The vegetation within all study areas corresponds to grassland of the *Mesobrometum* type. In the region of Schaffhausen this meadow type is usually cut once a year in mid June and very rarely or not at all fertilized. Prior to experimental management, two study areas had been used for hay-making (study areas 1 and 2), two others having been abandoned for ten and twenty years, respectively (study areas 3 and 4). One of the areas used until experimental management was started was drier and poorer in nutrients (study area 1) than the other (study area 2).

The phytosociological classification of the study areas offers some problems as far as nomenclature is concerned. According to ZOLLER (1954), who studied the dry grasslands in this region, our study area 1 should be considered as a *Medicago falcatae-Mesobrometum*, whereas study area 2 should correspond to a *Dauco-Salvio-Mesobrometum*; the study areas 3 and 4 abandoned for ten and 20 years respectively would represent the *Seselio libanotidis-Mesobrometum*.